

Solar modulation of the galactic cosmic rays

Kazuoki Munakata[1]

[1] Physics Department, Shinshu Univ

The galactic cosmic rays penetrating deep into the heliosphere interact with the solar magnetic field being carried radially outward by the solar wind. The interaction with the large-scale ordered field causes the gradient- and curvature-drift motions of cosmic-rays in the inner heliosphere, while the interaction with the irregular (or disordered) field component results in the pitch angle scattering of cosmic-rays. The pitch angle scattering by the magnetic irregularities embedded in the radially expanding solar wind causes the deceleration (called the adiabatic cooling) and the outward convection of cosmic-ray. This process called the solar modulation of the galactic cosmic rays will be discussed with an emphasis on the charge dependent features of the effect.

The galactic cosmic rays penetrating deep into the heliosphere interact with the solar magnetic field being carried radially outward by the solar wind. The interaction with the large-scale ordered field causes the gradient- and curvature-drift motions of cosmic-rays in the inner heliosphere, while the interaction with the irregular (or disordered) field component results in the pitch angle scattering of cosmic-rays. The pitch angle scattering by the magnetic irregularities embedded in the radially expanding solar wind causes the deceleration (called the adiabatic cooling) and the outward convection of cosmic-ray. This process called the solar modulation of the galactic cosmic rays will be discussed with an emphasis on the charge dependent features of the effect.