

Isotopic analysis of seasonal change of nitrogen cycling in Lake Kizaki.

Yuji Sasaki[1]; Keisuke Koba[2]; Muneoki Yoh[3]; Akiko Makabe[4]; Masamichi Yamamoto[5]; Yuichiro Ueno[6]; Mayuko Nakagawa[7]; Naohiro Yoshida[8]; Sakae Toyoda[9]

[1] Environmental Science on Biosphere, TUAT; [2] Tokyo University Agric Tech; [3] Tokyo Univ. Agri. Tech.; [4] Environmental Sci and Tech, Tokyo Institute of Tech; [5] Research and Education Center for Inlandwater Environ., Shinshu Univ.; [6] Global Edge Inst., Tokyo Tech.; [7] Applied chemistry, TITech; [8] IGSSE, Tokyo Institute of Technology; [9] Environmental Chemistry and Engineering, Tokyo Tech

The stable isotope composition of nitrate and nitrite was measured during summer to winter 2007 in Lake Kizaki. In epilimnion, enrichment of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ was not observed with decrease of nitrate. In autumn, isotopic signal was not shown in all layers. This suggest that nitrification did not occur in Lake Kizaki in autumn.