

AAS021-22

会場:102

時間:5月23日 15:30-15:45

## Variations of atmospheric radon-222 at Rishiri Island, Japan and traced fetch regions Variations of atmospheric radon-222 at Rishiri Island, Japan and traced fetch regions

Chunmao Zhu<sup>1\*</sup>, Hisayuki Yoshikawa<sup>1</sup>, Hidekadzu Matsueda<sup>2</sup>, Yosuke Sawa<sup>2</sup>, Yosuke Niwa<sup>2</sup>, akira WADA<sup>3</sup>, Hiroshi Tanimoto<sup>4</sup>  
Chunmao Zhu<sup>1\*</sup>, Hisayuki Yoshikawa<sup>1</sup>, Hidekadzu Matsueda<sup>2</sup>, Yosuke Sawa<sup>2</sup>, Yosuke Niwa<sup>2</sup>, akira WADA<sup>3</sup>, Hiroshi Tanimoto<sup>4</sup>

<sup>1</sup>Grad. School Env. Sci., Hokkaido Univ., <sup>2</sup>Geochem. Res. Div., Meteor. Res. Inst., <sup>3</sup>Meteoro. College, <sup>4</sup>National Inst. Env. Studies

<sup>1</sup>Grad. School Env. Sci., Hokkaido Univ., <sup>2</sup>Geochem. Res. Div., Meteor. Res. Inst., <sup>3</sup>Meteoro. College, <sup>4</sup>National Inst. Env. Studies

Measurements of atmospheric <sup>222</sup>Rn were made on Rishiri Island (45.1°N, 141.2°E) over the period from December 2008 to December 2010, in order to discuss the fetch regions affecting atmospheric <sup>222</sup>Rn concentration and some other gases related to climate change. Atmospheric <sup>222</sup>Rn data showed variability of a diurnal, synoptic, and seasonal time scales, which will allow us to evaluate transport and mixing schemes of atmospheric and chemical transport models.

Atmospheric <sup>222</sup>Rn concentration indicated a clear diurnal variation in summer, which was characterized of a maximum appearing before the dawn and a minimum in the afternoon, with amplitude of 0.64 Bq m<sup>-3</sup>. This could be caused by the accumulation of <sup>222</sup>Rn emitted from the soil (Rishiri Island) in the stable nocturnal boundary layer, and vertical mixing of surface air with upper air due to convection (Zahorowski *et al.*, 2008). The amplitude of diurnal variation is relatively lower than that of the synoptic influence. No clear diurnal variation occurred during the remaining seasons.

Atmospheric <sup>222</sup>Rn concentration in daytime, selected to remove effects of local source on <sup>222</sup>Rn concentration, showed a broad minimum in winter and a maximum in summer, with an accompanying significant short-term variability. In February monthly mean of atmospheric <sup>222</sup>Rn concentration was 3.23 Bq m<sup>-3</sup>, and in July 0.95 Bq m<sup>-3</sup>. In Rishiri Island, short-term variations corresponding to the synoptic influence were large in February as compared with those in July. The amplitude of the seasonal variation (2.28 Bq m<sup>-3</sup>) was somewhat larger than that of Sado Island located in the Sea of Japan. At Sado Island, maximum <sup>222</sup>Rn concentration occurred in winter and minimum in summer, which was the pattern attributable to the onshore-offshore pattern of the Asian monsoons (Chambers *et al.*, 2009).

Backward trajectory analysis of air fetch regions was conducted using extremely low and high radon events. In the annual basis, most (76.7%) of high radon events was observed in winter, of which the air masses originated predominately from 40°N to 60°N of Eurasian continent. 41.7% of low radon events were observed in summer, of which air masses usually originated from the relatively lower latitude of the western North Pacific, and 48.3% of low radon events were observed in spring, most in May. In May, back trajectory analysis showed a pattern that the origin of air masses was in the west North Pacific, and air mass moved westward en route the southernmost of Sea of Okhotsk to the RIO.

キーワード: atmospheric tracer, back trajectory analysis, fetch regions, radon

Keywords: atmospheric tracer, back trajectory analysis, fetch regions, radon