

APE031-07

会場:104

時間:5月25日 15:45-16:00

## 瀬戸内海別府湾の過去100年にわたる植物プランクトン動態：陸域・外洋からの栄養塩供給の影響

### Phytoplankton dynamics during the 20th century in Beppu Bay: effects of nutrient supply from land and shelf waters

槻木 玲美<sup>1\*</sup>, 加三千宣<sup>1</sup>, 谷幸則<sup>2</sup>, 守屋和佳<sup>3</sup>, 郭新宇<sup>4</sup>, 國弘忠生<sup>4</sup>, 齊藤光代<sup>4</sup>, 藤井直樹<sup>4</sup>, 武岡英隆<sup>4</sup>

Narumi Tsugeki<sup>1\*</sup>, Michinobu Kuwae<sup>1</sup>, Yukinori Tani<sup>2</sup>, Kazuyoshi Moriya<sup>3</sup>, Xinyu Guo<sup>4</sup>, Tadao Kunihiro<sup>4</sup>, Mitsuyo Saito<sup>4</sup>, Naoki Fujii<sup>4</sup>, Hidetaka Takeoka<sup>4</sup>

<sup>1</sup> 愛媛大学・上級研究員センター, <sup>2</sup> 静岡県大・環境, <sup>3</sup> 早稲田・教育・地球科学, <sup>4</sup> 愛媛大学・沿岸環境科学研究センター  
<sup>1</sup>SRFC, Ehime Univ., <sup>2</sup>Environmental Science, Shizuoka Univ., <sup>3</sup>Earth Sciences, Waseda Univ., <sup>4</sup>CMES, Ehime Univ.

Anthropogenic activities during the 20th century have greatly increased the flux of nutrients to aquatic ecosystems worldwide, with an accelerating rate of increase. However, in recent years there have been signs of oligotrophication in some estuaries of developed countries due to an excess removal of nutrients from the discharged water of sewage treatment plants or the impoundment of rivers by dams. In the Seto Inland Sea, which plays a major role in coastal fishing part of Japan, after the progress of eutrophication the hasty reduction of phosphorus loading could have lowered the phytoplankton primary production and also caused a detrimental effect on the fishery production. On the other hand, the Seto Inland is also likely to be affected by the variation of Kuroshio Current axis. To establish an ecosystem management plan for a estuarine ecosystem, it is essential to understand how the phytoplankton community has been altered in response to these environmental changes. In this study, we examined diatom remains and fossil pigments in Beppu Bay sediment to reveal historical changes in the phytoplankton community structure over the past 100 years in the Seto Inland Sea, and identify environmental factors that caused such changes.

Sedimentary analysis showed that the abundance of phytoplankton in Beppu Bay drastically increased since the 1960s when the fluxes of TN and TP rapidly increased. Afterwards, that of phytoplankton decreased, but however again increased greatly in the early 1990s. These sedimentary records were almost consistent with the monitoring data of chlorophyll a in this bay, although monitoring data are sporadic. We report and discuss the causal factors driving the changes in phytoplankton community structure focused on the effects of nutrient loading from the watershed and year-to-year variation of Kuroshio axis.

キーワード: 近過去, 瀬戸内海, 黒潮, 富栄養化, 20世紀, 植物プランクトン

Keywords: 20th century, the Seto Inland Sea, Kuroshio Current, eutrophication, paleoceanography, phytoplankton