

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



MIS036-P18

会場: コンベンションホール

時間: 5月26日 14:15-16:15

GPS データから推定した 2011 年東北地方太平洋沖地震の地震時すべりと余効すべり

Coseismic and postseismic slip for the 2011 Tohoku earthquake inferred from inversion of GPS data

福田 淳一^{1*}, 青木 陽介¹, 加藤 照之¹

Jun'ichi Fukuda^{1*}, Yosuke Aoki¹, Teruyuki Kato¹

¹ 東京大学地震研究所

¹ERI, University of Tokyo

We used GPS data recorded by the continuous GPS network, GEONET, to estimate spatial distribution of coseismic slip and afterslip for the 2011 M=9.0 Tohoku earthquake. We analyzed the GPS data on March 11, 2011, using the GIPSY-OASIS software to estimate station coordinates every 300 seconds. This analysis allows us to separate coseismic displacements due to the mainshock from coseismic displacements due to large aftershocks and early postseismic deformation. The coseismic displacements due to the mainshock were inverted for spatial distribution of slip on the plate interface. Our result shows that the mainshock ruptured primarily the shallower part of the plate interface than asperities of historical M=7-8 earthquakes, where interseismic slip deficit rate was inferred to be low. The estimated maximum slip is 33.6 m and the moment magnitude from the estimated slip distribution is Mw=8.9.

We also analyzed GPS data after March 11 with the GIPSY-OASIS software to estimate daily station coordinates. The daily GPS time series show significant postseismic deformation. We inverted the data for distribution of afterslip on the plate interface. The inferred afterslip is mainly located in the deeper and southern parts of the coseismic rupture region. As of April 2, 2011, the estimated maximum afterslip is 0.9 m and the moment magnitude from the estimated slip distribution is Mw=8.2.