

MIS036-P61

会場: コンベンションホール

時間: 5月26日 14:15-16:15

反射法地震探査と海底調査による2011年東北地方太平洋沖地震の断層システム Fault System of the 2011 Off-Tohoku Earthquake; Insight from Seismic Reflection Data and Seafloor Observations

辻 健^{1*}, 伊藤喜宏², 木戸元之², 長田幸仁², 藤本博巳², 芦寿一郎⁴, 木下正高³, 松岡俊文¹

Takeshi Tsuji^{1*}, Yoshihiro Ito², Motoyuki Kido², Yukihiro Osada², Hiromi Fujimoto², Juichiro Ashi⁴, Masataka Kinoshita³, Toshifumi Matsuoka¹

¹ 京都大学, ² 東北大学, ³ 海洋研究開発機構, ⁴ 東京大学

¹ Kyoto University, ² Tohoku University, ³ JAMSTEC, ⁴ University of Tokyo

Faults related to the tsunamigenic 2011 off-Tohoku earthquake (Mw 9.0) were investigated by using multi-channel seismic reflection data acquired in 1999 and submersible seafloor observations from 2008. The shallow fault geometries as well as submarine slide are important to reveal mechanisms of the huge tsunami generation. On the reflection seismic profile, we identified three predominant faults; (A) backstop reverse fault working as a boundary between seaward accreted sequence and landward consolidated sequence, (B) a branch fault constructing the seafloor slope break, and (C) vertical fault extending toward a seafloor ridge. Several imbricate thrusts are developed within the accreted sequence seaward of (A) backstop reverse fault. Between (A) the backstop reverse fault and (B) the branch fault, underplating structures are identified on the profile. Displacement along (C) the vertical fault has offset a Cretaceous sequence surface by ~800 m. The location of these faults interpreted in the seismic reflection profile is distributed around the area with largest slip and tsunami induction of the 2011 event. Cold-seep communities and a high scarp observed at these fault traces suggest current activity on these faults. We interpret the fault system in the seismic profile as a shallow extension of the seismogenic fault that may have contributed to induction of the huge tsunami.