

静止軌道およびISSによる対流圏NO₂観測における地表面BRDFの影響 The effect of the surface BRDF on the measurement of tropospheric NO₂ from a geostationary orbit and ISS

野口 克行^{1*}, アンドレアス・リヒター², ジョン・パローズ², 入江 仁士³, 北 和之⁴
NOGUCHI, Katsuyuki^{1*}, Andreas Richter², John P. Burrows², IRIE, Hitoshi³, KITA, Kazuyuki⁴

¹ 奈良女子大学, ² ブレーメン大学, ³ 海洋研究開発機構, ⁴ 茨城大学

¹Nara Women's University, ²University of Bremen, ³JAMSTEC, ⁴Ibaraki University

BRDF (Bidirectional Reflectance Distribution Function) is the dependency of the surface reflectance on incident and output directions. Zhou et al. [2010] showed that the surface BRDF affects the retrieval of the tropospheric NO₂ column density with sun-synchronous low Earth orbit (LEO) satellites, as line-of-sight angles largely change in the sun-synchronous LEO measurements. In the present study, we have estimated the influence of BRDF on the tropospheric NO₂ DOAS retrieval supposing the observations of air pollution over Tokyo from a geostationary orbit (GEO) and ISS. In the analysis, we used the results of actual surface measurements: the MODIS BRDF product released by NASA. The result showed that the use of the assumption of Lambertian surface instead of BRDF would cause up to a 30% difference in the reflectance value. We also show the influence of such a difference on the tropospheric NO₂ column density measurements.

キーワード: 静止衛星, 国際宇宙ステーション, 対流圏化学, 二酸化窒素, 双方向反射率分布関数
Keywords: Geostationary orbit, ISS, tropospheric chemistry, NO₂, BRDF