Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



AAS22-04

会場:201B

時間:5月21日09:45-10:00

SMILES で観測された成層圏における反応 ClO + HO2 -> HOCl + O2 について On the reaction ClO + HO2 -> HOCl + O2 based upon SMILES observation

眞子 直弘 ^{1*}, 鈴木 睦 ¹, 佐野 琢己 ¹, 光田 千紘 ², 今井 弘二 ³, 高橋 けんし ⁴, 塩谷 雅人 ⁴ MANAGO, Naohiro^{1*}, SUZUKI, Makoto¹, SANO, Takuki¹, MITSUDA, Chihiro², IMAI, Koji³, TAKAHASHI, Kenshi⁴, SH-IOTANI, Masato⁴

1 宇宙航空研究開発機構宇宙科学研究所, 2 富士通 FIP, 3 とめ研究所, 4 京大生存圏研

SMILES (Superconducting Submillimeter-Wave Limb Emission Sounder) is an instrument to measure global distribution of minor species in the middle atmosphere by limb observation. It was attached to the Japanese Experiment Module (JEM) on the International Space Station (ISS) and obtained a half year's worth of data between mid October, 2009 and mid April, 2010. SMILES has an advantage in low system noise realized by cooling the receiver to 4 degrees kelvin with a mechanical cooler, and it enables to measure distribution of trace gases such as O3, HCl, ClO, HO2 and HOCl with high sensitivity.

It is a well-known fact that inorganic chlorine play an important role in the stratospheric chemistry. However, it is not fully understood quantitatively due to the limited precision of parameters such as the abundance of HCl, total abundance of inorganic chlorine, the ratio between HCl and other inorganic chlorine, the ratio between ClO and HOCl, and so on.

In this research, we estimated the reaction rate of CIO + HO2 -> HOC1 + O2 with steady-state approximation. By using SMILES L2 ver.2.1 for the HOCl concentration and MODTRAN5 to calculate the photodissociation rate of HOCl, we obtained reaction rates similar to the JPL 2006 value at the altitude range of 30 - 40 km.

キーワード: 成層圏, 無機塩素系化学種 Keywords: stratosphere, inorganic chlorine

¹ISAS/JAXA, ²Fujitsu FIP Corp., ³TOME R&D Inc., ⁴RISH/Kyoto Univ.