

日本の首都圏で日最高気温 40.9℃ を記録した極端な猛暑の形成メカニズム Formation Mechanisms of the Extreme High Surface Air Temperature of 40.9°C Observed in the Tokyo Metropolitan Area

高根 雄也^{1*}, 日下 博幸²

TAKANE, Yuya^{1*}, Hiroyuki Kusaka²

¹ 筑波大学大学院生命環境科学研究科, ² 筑波大学計算科学研究センター

¹Graduate School of Life and Environmental Sciences, University of Tsukuba, ²Center for Computational Sciences, University of Tsukuba

A record-breaking high surface air temperature in Japan of 40.9°C was observed on 16 August 2007 in Kumagaya, located 60 km northwest of central Tokyo. In this study, the formation mechanisms of this extreme high temperature event are statistically and numerically investigated using observational data and the Weather Research and Forecasting (WRF) model. The extreme event is caused by a combination of two particular factors: 1) Persistent sunshine and a lack of precipitation during the seven consecutive days preceding 16 August 2007 were seen in Kumagaya. This was the 12th-longest stretch of clear-sky days in July and August from 1998 up to 2008. Persistent clear-sky days allow the ground surface to dry out, which produces an increase in sensible heat flux from the ground surface. This contributes to the extreme event, and its mechanism is qualitatively supported by the results of sensitivity experiments of soil moisture on surface air temperature. 2) A foehnlike wind (Fig.1) appears in the numerical simulation, which is caused by diabatic heating with subgrid-scale turbulent diffusion and sensible heat flux from the ground surface when this airflow passes in the mixed layer over the Chubu Mountains and the inland of the Tokyo metropolitan area. Backward trajectory analysis and Lagrangian energy budget analysis show that the foehnlike wind plays a more important role in the extreme event than the adiabatic dynamic foehn pointed out by previous studies.

キーワード: 極端な高温, フェーン, 連続晴天, WRF, 熊谷市

Keywords: Extreme high temperature, Foehn, Persistent clear-sky days, WRF, Kumagaya city

