

Puの地表出現を遅らせるフェリハイドライト・コロイド Colloidal ferrihydrite retards the appearance of Pu on the Earth's surface

月村 勝宏^{1*}, 鈴木 正哉¹, 鈴木 庸平², 村上 隆²

TSUKIMURA, Katsuhiko^{1*}, SUZUKI, Masaya¹, SUZUKI, Yohey², MURAKAMI, Takashi²

¹ 産業技術総合研究所, ² 東京大学

¹ AIST, ² The University of Tokyo

Radioactive wastes from nuclear power plants must be disposed of safely. A large amount of Pu in the wastes is sorbed to colloidal ferrihydrite and moves in groundwater. This paper describes the behavior of Pu sorbed to colloidal ferrihydrite in nuclear waste disposal sites. When the concentration of ferrihydrite is higher than 10-5 mol Fe /L, most Pu is sorbed to the ferrihydrite. Ferrihydrite persists in groundwater as long as nitrate is present. When the filtration effect of geologic media is small, the gravity moves colloidal ferrihydrite to deep underground; the velocity is 0.12 m/year when the size of ferrihydrite is 70 nm. When the filtration effect is large, ferrihydrite is not transported further. In both the cases, ferrihydrite retards the appearance of Pu on the Earth's surface.

キーワード: コロイド, フェリハイドライト, プルトニウム, 核種移行, 重力効果, 吸着

Keywords: colloid, ferrihydrite, plutonium, transport of radionuclides, gravity effect, sorption