

ハマサンゴ骨格における Mg 同位体分別 : 飼育実験による低緯度域の新規古気候プロキシの評価

Mg isotope fractionation in Porites coral skeletons: evaluation of a new climate proxy from culture experiments

吉村 寿紘^{1*}, 谷水 雅治², 鈴木 淳³, 井上 麻夕里⁴, 古賀 奏子⁵, 酒井 一彦⁵, 川幡 穂高⁴

YOSHIMURA, Toshihiro^{1*}, Masaharu Tanimizu², SUZUKI, Atsushi³, INOUE, Mayuri⁴, Kaneko Koga⁵, Kazuhiko Sakai⁵, KAWAHATA, hodaka⁴

¹ 東京大学・新領域創成科学研究科, ² 海洋研究開発機構・高知コア, ³ 産業技術総合研究所・地質情報研究部門, ⁴ 東京大学・大気海洋研究所, ⁵ 琉球大学・熱帯生物圏研究センター

¹graduate School of Frontier Sciences, The University of Tokyo, ²KCC, JAMSTEC, ³geological Survey of Japan, National Institute of Advanced Industrial Science and Technology, ⁴atmosphere and Ocean Research Institute, The University of Tokyo, ⁵Tropical Biosphere Research Center, University of the Ryukyus

Recent developments in inorganic mass spectrometry have allowed new stable isotope systems to be explored by paleoceanographers. In the field of low-latitude climate reconstruction, proxy developments in massive hermatypic corals are still highly desirable for more accurate temperature reconstructions. Stable isotope compositions of the major and minor constituents, such as Ca and Sr, of coral aragonite are potentially new proxies for environmental studies (Boehm et al., 2006; Fietzke and Eisenhauer, 2006). These studies have focused on the temperature dependency of isotope fractionation during the precipitation of biogenic carbonates, because water temperature is the most fundamental parameter controlling the earth's surface environment. Although Sr/Ca is regarded as useful temperature proxy, Mg/Ca is thought to mainly reflect the growth rate (e.g., Inoue et al., 2007). Evaluation of Mg isotope fractionation process in hermatypic coral aragonite must be addressed in order to understand a coral biomineralization and develop a new palaeoceanographic proxy. Many researchers have analyzed Ca isotopes in biogenic and inorganic calcium carbonates, with the first objective being to evaluate them as a potential new paleothermometer. They have found a small but resolvable temperature dependence of Ca isotope fractionation (e.g., Gussone et al., 2009). Strontium is important minor element in CaCO₃, and in inorganic aragonite and cultured hermatypic corals, it displays similar isotope fractionation mechanisms to those of Ca (Fietzke and Eisenhauer, 2006). Previous studies have reported some steeper temperature dependence (slopes) in Ca and Sr isotope fractionation of biogenic CaCO₃ and the differences in temperature slopes are interpreted as a large kinetic isotopic effects depending on how fast the crystal growing rate and relative mass difference (rmd) among isotopes. Growth effects on Mg isotope are expected to be highly concerned since Mg/Ca is known to clearly dependent on growth rates (Inoue et al., 2007). In this study, we performed high-precision Mg isotope measurements in Porites coral cultured in thermostated tanks. We here evaluate the plausible factors controlling Mg isotope fractionation in coral CaCO₃ skeletons and present the relationships between Mg isotope values and e.g., water temperature, minor elemental concentrations, growth rates.

キーワード: 造礁サンゴ, マグネシウム同位体, MC-ICP-MS, 飼育実験

Keywords: hermatypic coral, magnesium isotope, MC-ICP-MS, culture experiment