

電離圏観測による津波速報 Early warning system with GPS-TEC observation

鴨川 仁^{1*}, 柿並 義宏²

KAMOGAWA, Masashi^{1*}, KAKINAMI, Yoshihiro²

¹ 東京学芸大学物理学科, ² 北海道大学大学院理学研究院附属地震火山研究観測センター

¹Dpt. of Phys., Tokyo Gakugei Univ., ²Institute of Seismology and Volcanology, Hokkaido University

Traveling ionospheric disturbances generated by an epicentral ground/sea surface motion, ionospheric disturbances associated with Rayleigh-waves as well as post-seismic 4-minute monoperiodic atmospheric resonances and other-period atmospheric oscillations have been observed in large earthquakes. In addition, a giant tsunami after the subduction earthquake produces an ionospheric hole which is widely a sudden depletion of ionospheric total electron content (TEC) in the hundred kilometer scale and lasts for a few tens of minutes. The tsunamigenic ionospheric hole detected by the TEC measurement with Global Position System (GPS) was found only in huge subduction earthquakes. This occurs because plasma is descending at the lower thermosphere where the recombination of ions and electrons is high through the meter-scale downwelling of sea surface at the tsunami source area, and is highly depleted due to the chemical processes. The results imply that magnitude of the tsunamigenic ionospheric hole is related to that of the tsunami. It means that we can directly observe the tsunami several minutes after the subduction earthquake occurs.

キーワード: 津波速報, 津波電離圏ホール, GPS

Keywords: Early warning system, Tsunamigenic ionospheric disturbance, GPS