

領域大気輸送モデルを用いた地表および海洋への放射性物質沈着量分布の推定 Estimation of radionuclides' deposition over the land and ocean using a regional chemical transport model

滝川 雅之^{1*}, 渡邊 明²

TAKIGAWA, Masayuki^{1*}, WATANABE, Akira²

¹ 独立行政法人海洋研究開発機構, ² 福島大学

¹Japan Agency for Marine-Earth Science and Technology, ²Fukushima University

To reveal the transport and deposition processes of radionuclides that have been emitted from the Fukushima Daiichi nuclear power plant after the nuclear accident caused by the Tohoku earthquake and tsunami, we have conducted numerical simulation of radionuclides (¹³¹I and ¹³⁷Cs) using a regional chemical transport model. Data assimilation using nudging for wind field and relative humidity was applied with the 10-minutes' interval ground-based observation, and the model-calculated meteorological fields with the ground-based observations in Fukushima prefecture showed that the model was capable to reproduce the change of wind direction following to the passage of low-pressure systems, and precipitation periods in March 2011. The transport and deposition process of two major discharges on March 15 was estimated using tag-tracer methods. The air mass from the first discharge released during 7-10 JST was transported southward, and 13% of ¹³⁷Cs was deposited mainly over Kanto area via dry deposition process. In the afternoon, wind direction was changed following to the approach of low-pressure system, and the air mass from the first discharge was transported northward and 13% of ¹³⁷Cs was deposited mainly over Naka-dori area in Fukushima prefecture via wet deposition process corresponding with the precipitation since 15-16 JST. The model showed that the deposition over Naka-dori area occurred within 3 hours after the beginning of precipitation, although the intensity of precipitation was less than 0.5 mm/hr during that period. The air mass from the second discharge occurred at 13-17 JST was transported northwestward, and 47% of ¹³⁷Cs in second discharge were estimated to be deposited over the land. The comparison of accumulated deposition of ¹³⁷Cs at Azuma-yama mountains supposed that the radionuclides from the Fukushima Daiichi nuclear power plant were trapped within the boundary layer, and the highest concentration existed at around 1000-1500 m above the ground level.

キーワード: 放射性物質, 化学輸送モデル, 大気化学, 物質輸送

Keywords: radionuclides, chemical transport model, atmospheric chemistry, chemical transport