

古大気組成と太陽進化トレーサーとしての波長依存同位体効果 Wavelength dependent isotope effect as a tracer for paleoatmosphere and solar evolution?

上野 雄一郎^{1*}, ダニエラチェ セバスチアン¹, 遠藤 美朗¹
UENO, Yuichiro^{1*}, DANIELACHE, Sebastian¹, ENDO, Yoshiaki¹

¹ 東京工業大学

¹Tokyo Institute of Technology

Discovery of sulfur isotope anomaly in Archean sedimentary rocks demonstrated that Earth's early atmosphere was virtually oxygen-free (Farquhar et al., 2000). The anomalous fractionation is produced by photolysis of SO₂. The fractionation factors are sensitive to the wavelength of irradiated UV (Danielache et al., 2008), thus is a potential tracer to understand chemistry of atmosphere and possibly change in solar spectrum. We newly determined higher resolution UV absorption cross sections of not only ³²SO₂, ³³SO₂ and ³⁴SO₂ but also ³⁶SO₂ within the two absorption bands: (1) 190 ? 220 nm and (2) 250 ? 320 nm. These data together with atmospheric reaction model allow us to predict isotopic compositions of photochemical product. The calculated photochemical fractionation pattern assuming broadband solar UV flux reproduce our previous work (Danielache et al., 2008), though the effect of UV shielding by each atmospheric species including SO₂ itself differ from previously estimated trend. Nonetheless, almost all of the simulations result in D³⁶S/D³³S ratio of -0.9 ~ -1.1, generally reproducing those observed in Archean sedimentary rocks. Thus, we conclude that photodissociation of SO₂ was a primary MIF-yielding reaction in the Archean atmosphere. Our simulation predict, however, the remaining SO₂ after UV photolysis acquires positive D³³S as opposed to widely-accepted previous model where H₂SO₄ (-D³³S) and S₈ (+D³³S) aerosols carried opposite MIF signals into ocean and sediment (Pavlov and Kasting, 2002; Ono et al., 2003). We speculate the possibility that almost Archean sulfide deposits were produced by sulfate reduction. The new model requires relatively inert reducing form of sulfur reservoir. If the atmosphere was strongly reducing and contained high level of CO or CH₄, photolytically produced SO was finally transferred into OCS (Ueno et al., 2009) or organo-sulfur compounds (Dewitt et al., 2010), respectively, which remained in the atmosphere and were not readily converted into sulfide. Occasional oxidation of the reducing sulfur pool enhanced sulfate concentration and deposited rare sulfate minerals with negative D³³S. The new dynamical sulfur cycle model may explain observed heterogeneity of S-MIF records in the basin to microscopic scale. This scenario is tested by our ongoing photochemical chamber experiment. If correct, this implies more reducing Archean atmosphere than previously thought.

キーワード: 太古代, 大気, 太陽

Keywords: Archean, atmosphere, sun