

レッドスプライトによる VLF 帯電波擾乱についての数値解析 Numerical simulations VLF signal perturbations due to red sprites

岩本 真彦^{1*}, 芳原 容英¹, 成毛亮介¹, 太田 健次², 廣木 一 亮³, 湊原 哲也³, 早川 正士¹

IWAMOTO, Masahiko^{1*}, HOBARA, Yasuhide¹, Ryosuke Naruke¹, Kenji Ohta², Kazuaki Hiroki³, Tetsuya Minatohara³, HAYAKAWA, Masashi¹

¹ 電気通信大学 情報理工学研究所 情報・通信工学専攻, 東京, 日本, ² 中部大学, 春日井, 日本, ³ 津山高専, 津山, 日本
¹Graduate School of Informatics and Engineering, UEC, Tokyo, Japan, ²Chubu Univ, Kasugai, Japan, ³Tsuyama National College of Technology, Tsuyama, Japan

In this paper we perform 3D finite-difference time-domain (FDTD) method to compute the subionospheric VLF signal perturbations due to the ionization from mesospheric transient event such as red sprites. Spatial scales of columns are determined by the sprite images obtained from our optical observations during winter lightning activities over the sea of Japan. Numerical results indicate that the multiple sprites generate the complicated scattering pattern of the VLF transmitter waves depending on special orientation and extent of sprite ionization columns. Spatial dependence of the scattered amplitude are compared with those from the experimental results of VLF observation network.