

Large scale influence on precipitation propagation over Indonesia Large scale influence on precipitation propagation over Indonesia

Marzuki Marzuki^{1*}, Hiroyuki Hashiguchi¹, Masayuki Yamamoto¹, Manabu D. Yamanaka², Shuichi Mori²
MARZUKI, Marzuki^{1*}, HASHIGUCHI, Hiroyuki¹, YAMAMOTO, Masayuki¹, YAMANAKA, Manabu D.², MORI, Shuichi²

¹Research Institute for Sustainable Humanosphere, Kyoto University, Japan, ²Japan Agency for Marine-Earth Science and Technology (JAMSTEC), ³Department of Physics, Andalas University, Indonesia

¹Research Institute for Sustainable Humanosphere, Kyoto University, Japan, ²Japan Agency for Marine-Earth Science and Technology (JAMSTEC), ³Department of Physics, Andalas University, Indonesia

Mechanism of organization of cloud clusters (CCs) over Indonesian Maritime Continent (IMC) is linkage between its complex geographical variation and large-scale atmospheric structure/circulation. Super cloud cluster (SCC), which is recognized as eastward-propagating envelopes of convection, composed of westward-propagating CCs in mesoscale [1]. In this study, the relationship between statistical properties (zonal span, duration, and propagation speed) of cloud episodes/streaks in Hovmoller space and vertical shear of horizontal wind, convectively coupled Kelvin waves and Madden-Julian oscillation (MJO) is investigated.

Ten years of hourly infrared (IR) brightness temperature (T_b) are used to study the cloud episodes/streaks over the IMC. To estimate the statistics of cloud streak, a 2D-autocorrelation function is applied to the data in the Hovmoller space [2]. Daily interpolated OLR data are used to diagnose the MJO and Kelvin wave during the interest period. The MJO is diagnosed using a 30-96 days bandpass Lanczos filter on daily OLR anomalies following [3]. The Kelvin wave filtering retains eastward-propagating OLR signals within the 2.5-20 day period and wave numbers 1-14 [4].

Table 1 shows a summary of all streak characteristics in the 10° S- 10° N band for four classified months. In general, westward moving streaks are dominant, longer span and move faster than eastward moving streaks. Seasonal variation is also observed. The relationship between statistical properties of cloud episodes and large scale influence (e.g., MJO, Kelvin wave) will be presented in the meeting.

[1] T. Nakazawa, Tropical super cloud clusters within intraseasonal variations over the western Pacific. *J. Meteor. Soc. Japan*, 66, 823-839, 1988.

[2] Carbone, R. E., J. D. Tuttle, D. A. Ahijevych, S. B. Trier, Inferences of Predictability Associated with Warm Season Precipitation Episodes, *J. Atmos. Sci.*, 59, 2002.

[3] Kiladis, G. N., K. H. Straub, P. T. Haertel, Zonal and Vertical Structure of the Madden-Julian Oscillation. *J. Atmos. Sci.*, 62, 2790-2809, 2005.

[4] Wheeler, M. C., and G. N. Kiladis, Convectively coupled equatorial waves: Analysis of clouds in the wavenumber-frequency domain. *J. Atmos. Sci.*, 56, 374-399, 1999.

キーワード: Precipitation propagation, Large scale, Indonesia

Keywords: Precipitation propagation, Large scale, Indonesia

MIS29-P10

会場:コンベンションホール

時間:5月24日 17:15-18:30

Table 1: Summary of all streak characteristics in the 10°S-10°N band for four classified months.

Characteristics	DJF	MAM	JJA	SON
<i>Westward</i>				
No.	2839	2576	2383	3033
Mean speed (m/s)	-16.8	-16.4	-17.1	-16.6
Mean duration (h)	9.3	9.0	9.6	10.1
Mean span (km)	507.8	482.5	538.5	546.5
<i>Eastward</i>				
No.	1147	1058	635	733
Mean speed (m/s)	16.0	14.8	16.1	15.3
Mean duration (h)	7.4	7.9	7.4	7.3
Mean span (km)	381.6	373.4	396.5	365.3
Ratio	2.5	2.4	3.8	4.1