

平均 Dst 指数を用いた電子放射線帯モデル

The electron radiation belt model depending on the mean Dst index

中村 雅夫^{1*}, 谷岡俊彦¹, 高田 拓², 浅井 佳子³, 松岡 彩子⁴, 長井 嗣信³

NAKAMURA, Masao^{1*}, Toshihiko Tanioka¹, TAKADA, Taku², ASAI, Keiko T.³, MATSUOKA, Ayako⁴, NAGAI, Tsugunobu³

¹ 大阪府立大学, ² 高知工業高等専門学校, ³ 東京工業大学, ⁴ 宇宙航空研究開発機構 宇宙科学研究所

¹Osaka Prefecture University, ²Kochi National College of Technology, ³Tokyo Institute of Technology, ⁴Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency

Electron radiation belt models are important for spacecraft designs to evaluate the total radiation dose. The Akebono satellite has been in the highly elliptical orbit since 1989 and measured almost whole region of the inner radiation belt and the high latitude region of the outer radiation belt over a period of 22 years which is the 2 solar cycles. From the long-term Akebono satellite observation, we find that the logarithm of the annual mean high energy (>2.5MeV) electron flux shows a good correlation with the annual mean Dst index in the slot region and the outer belt. We propose the electron radiation model depending on the mean Dst index.

キーワード: 電子放射線帯モデル, あけぼの衛星, Dst 指数

Keywords: Electron radiation belt model, Akebono satellite, Dst index