

Enceladus 起源水分子による土星内部磁気圏内電子散乱のテスト粒子解析 Test-particle analysis of electron scattering in the Saturn's inner magnetosphere by neutral H₂O from Enceladus

田所 裕康^{1*}, 加藤 雄人¹

TADOKORO, Hiroyasu^{1*}, KATOH, Yuto¹

¹ 東北大学大学院理学研究科

¹ Graduate School of Science, Tohoku University

Cassini observations revealed that Saturn's moon Enceladus ejects neutral H₂O from its southern pole with temporal variability [e.g., Hansen et al., 2006]. This volcanic activity, so-called 'plumes', leads to the electromagnetic coupling between Saturn's ionosphere and plasmas around Enceladus. The coupling causes auroral activities around the footprint of Enceladus [Pryor et al., 2011]. They reported that observed field aligned fluxes of electrons and ions are sufficient to brighten the footprint aurora observed by EUV onboard Cassini. On the other hand, an electron precipitation into the atmosphere through pitch-angle scattering also causes auroral emissions. The dominant physical process controlling the activity of the footprint aurora is still controversial.

In the present study, for the quantitative evaluation of auroral emissions caused by the pitch-angle scattering through elastic collisions between magnetospheric electrons and H₂O particles, we have developed a spatially one dimensional test-particle simulation code along a dipole magnetic field at Enceladus ($L = 3.95$). We assume that the initial velocity distribution of energetic electrons at the magnetic equator forms a velocity distribution with a loss-cone. An interaction between an electron and a background neutral cloud is solved by the Monte-Carlo method using differential cross sections of elastic collisions for H₂O. We show a result of the variability of precipitating electrons and estimation of the expected brightness of auroral emissions.

キーワード: テスト粒子シミュレーション, エンケラドス・トーラス, 土星, 弾性衝突, 水分子プルーム, ピッチ角散乱
Keywords: test particle simulation, Enceladus torus, Saturn, elastic collision, H₂O plume, pitch angle scattering