

サイクル23全周期及び23/24極小期における太陽風加速モデルの検証 Relationship between solar wind speed and coronal magnetic field parameter through solar cycle 23 and 23/24 minimum

藤木 謙一^{1*}, 野田 桃太郎², 徳丸 宗利¹

FUJIKI, Ken'ichi^{1*}, Momotaro Noda², TOKUMARU, Munetoshi¹

¹ 名古屋大学太陽地球環境研究所, ² 名古屋大学大学院理学研究科素粒子宇宙物理学専攻

¹Solar-Terrestrial Environment Laboratory, Nagoya University, ²Division of Particle and Astrophysical Science, Graduate School of Science, Nagoya University

Solar wind acceleration mechanism is one of the most important issues to be solved in solar wind physics. We have been studied the relationship between velocity of solar wind which comes from coronal holes and coronal magnetic field parameter. Precedence researches in our group showed that solar wind speed, V had been in proportion to a parameter Bp/f from solar minimum in cycle 22/23 to solar maximum in cycle 23, where Bp and f are magnetic field strength and expansion factor of the magnetic flux tube, respectively [1]. In solar minimum in the cycle 23/24, however, polar magnetic field gets weaker and solar wind has lower density, lower temperature, and lower mass flux than the previous minimum [2]. For this reason, we examined the relationship between V and Bp/f from 22/23 to 23/24 solar minima. In this analysis we used following data set. Solar wind velocity map in each Carrington rotation was derived from the interplanetary scintillation measurements at Solar-Terrestrial Environment Laboratory. Coronal magnetic field in each Carrington rotation was calculated by potential field source surface model using synoptic photospheric magnetic field data observed at Kitt Peak National Solar Observatory. As results, we found that V always correlates positively Bp/f over the solar cycle. Then we focused on the difference of slopes of the regression lines between two solar minima. The slope in the 23/24 minimum becomes larger, in other words, the fast solar wind in the 23/24 minimum has smaller Bp/f compared to that in the 22/23 minimum. This result was compared with a theoretical model of solar wind acceleration [3] by taking account of the declining of solar wind parameters. We confirmed that the difference of regression lines in two solar minima is consistent with global trend of solar and solar wind variations.

Reference

[1] Fujiki et al., Adv. Space Res., Volume 35, Issue 12, 2185, (2005)

[2] McComas et al., Geophys. Res. Lett., 35, 18103, (2008)

[3] Suzuki, T. K., Astrophys. J. Lett., 640, 75, (2006)

キーワード: 太陽風, 惑星間空間シンチレーション, 太陽活動周期

Keywords: solar wind, interplanetary scintillation, solar cycle