

SUBARU/IRCS と補償光学を用いた木星近赤外オーロラ発光の高度分布観測 The first observation of the altitude distribution of Jovian near-IR auroral emission using SUBARU/IRCS with Adaptive Op

宇野 健^{1*}, 坂野井 健¹, 埜 千尋², 笠羽 康正¹, 佐藤 毅彦²

UNO, Takeru^{1*}, SAKANOI, Takeshi¹, TAO, Chihiro², KASABA, Yasumasa¹, SATOH, Takehiko²

¹ 東北大学・理・地球物理, ² 宇宙航空研究開発機構宇宙科学研究所

¹Tohoku Univ., ²ISAS/JAXA

The altitude emission profile is very important to understand that why the spatial distribution of the IR emission from H₂ and H₃⁺ are morphologically different (e.g., Raynaud et al., 2004). The origin of this morphological difference is still unknown. It may be caused by the difference of heating altitude and/or difference of precipitation energy.

Although the altitude distribution of IR auroral emission of H₂ and H₃⁺ is well discussed by the theoretical model (e.g., Kim et al., 1990; Grodent et al., 2001), observational study is limited. The observation of vertical distribution of H₃⁺ column density and vibrational-rotational temperature are only reported by Lystrup et al., 2008. And there is no vertical-resolved observation of H₂ emission.

Based on the model calculation, it is thought that the difference of IR emission altitude between H₂ and H₃⁺ is about 500-1000 km (Grodent et al., 2001). It is impossible to detect this vertical difference by ground-based observation, because the typical seeing of 0.6 arcsec is corresponding to the vertical resolution of about 1800 km at the Jupiter. The recent technique of Adaptive Optics (AO) makes it possible to get the high spatial resolved data about 0.1 arcsec, corresponding to the vertical resolution of about 300 km.

Simultaneous H₂ and H₃⁺ observation near 2.1 μm took place on 30 Nov. 2011 using the SUBARU/IRCS with AO188 system. The slit is set along rotational axis (vertical to the equator) at northern pole. Using Europa for the guide star for AO system, we succeeded the first limb observation of Jupiter H₂ and H₃⁺ IR auroral emissions.

In the polar region, H₂ emission lines S1(0), S1(1), and S1(2) at the wavelengths of 2.22, 2.12, 2.03 μm and several H₃⁺ emission lines are detected.

We will report the difference in the spatial and vertical distributions of those emissions and temperatures, derived from the observation.

Acknowledgement: Based on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

Keywords: Jupiter, ionosphere, Infrared, spectroscopy, thermosphere, aurora