

## CALCIUM-ALUMINIUM INCLUSION IN THE KABA METEORITE AND ITS APPLICATION TO ASTROMINERALOGY

GUCSIK, Arnold<sup>1\*</sup>, NISHIDO, Hirotsugu<sup>2</sup>, NINAGAWA, Kiyotaka<sup>3</sup>, Szaniszló Berczi<sup>4</sup>, Szabolcs Nagy<sup>4</sup>, Peter Abraham<sup>5</sup>, Ildiko Gyollai<sup>6</sup>, Akira Tsuchiyama<sup>1</sup>, Irakli Simonia<sup>7</sup>, Peter Rozsa<sup>8</sup>, Daniel Apai<sup>9</sup>, Krisztian Mihalyi<sup>8</sup>, Mihaly Nagy<sup>10</sup>, Jozsef Posta<sup>11</sup>

<sup>1</sup>Department of Earth and Planetary Sciences, Graduate School of Science, Osaka University,, <sup>2</sup>Research Institute of Natural Sciences, Okayama University of Science, Okayama, Japan, <sup>3</sup>Department of Applied Physics, Okayama University of Science, Okayama, Japa, <sup>4</sup>Institute of Physics, Department of Material Physics, Eotvos University, Budapest, Hungary, <sup>5</sup>Konkoly Observatory of Hungarian Academy of Sciences, Budapest, <sup>6</sup>Department of Lithospheric Research Center for Earth Sciences, University of Vienna, Austria, <sup>7</sup>Graduate Studies of Ilia State University, Tbilisi, Georgia, <sup>8</sup>University of Debrecen, Debrecen, Hungary, <sup>9</sup>Lunar and Planetary Laboratory, Department of Planetary Sciences, The Arizona University, <sup>10</sup>Reformed College of Debrecen, Debrecen, Hungary, <sup>11</sup>Debrecen University, Dept Chemistry

CAI of the Kaba meteorite has a complex texture and consists of spinel, anorthite and augite (fassaite), where spinel grains (up to 10 micron in size) are surrounded by anorthite and augite grains. CAIs are observed and maximum 1.8-2.0 mm in size. The composition of anorthite is An<sub>95.6</sub>Ab<sub>4.4</sub>Or<sub>0</sub>. Augite has a composition of En<sub>45.5-55.1</sub>Wo<sub>44.0-53.9</sub>Fs<sub>0.6-0.9</sub>.

The age of Kaba ? as determined from Mn-Cr dating ? is thought to be between 4,562 and 4,563 Ma (Hua et al. 2005). It is instructive to attempt to place the formation and properties of Kaba in the context of protoplanetary disk evolution as observed around other stars. Any such comparison relies on the zero points of the astronomical and cosmochemical timescales, i.e. the time of the protostellar collapse and the time of the CAI formation. While these zero points are likely to be slightly shifted, detailed comparisons of protoplanetary disk evolution and events in the proto-solar nebula suggest that they could not differ by more than 1 Myr, if the proto-solar nebula was a typical disk (Pascucci & Tachibana 2010). Consistent with the above description we assume that CAIs have formed at the time of or very shortly after the protostellar collapse.

In contrast, the younger disks in Cha I and Taurus frequently display disks with flaring geometry (disk opening angle increasing with radius, see e.g., Sz?cs et al. 2010, Ciesla and Dullemond 2010). These disks also commonly display sharp and prominent crystalline silicate peaks, revealing the presence of sub-micron-sized forsterite and enstatite grains (e.g. Apai et al. 2005) with a few disks showing amorphous silicate emission features. The observed evolution of the small, initially amorphous dust grains into larger, crystalline grains is poorly understood, but it is often thought that grain-grain collisions and destructive planetesimals collisions will replenish and gradually replace the dust population. In this context, Kaba grains could provide an insight into the dust population of a disk halfway between a young protoplanetary disk and a debris disk: if so, a substantial amount of the building blocks of Kaba may have been recycled material from previous generation of small bodies. Furthermore, a systematic Micro-Raman spectral study (as future work) of an interaction between the organic compounds and CAIs in Kaba meteorite can provide us better understanding of the evolution of organic matter in the early Solar System.

### References

- Apai D., Pascucci I., Bouwman J., Natta A., Henning Th. and Dullemond C.P. 2005: The Onset of Planet Formation in Brown Dwarf Disks. *Science* 310: 834-836.
- Ciesla F.J. and Dullemond C.P. 2010: Evolution of Protoplanetary Disk Structures. In *Protoplanetary Dust: Astrophysical and Cosmochemical Perspectives*, Eds.: Apai D. and Lauretta D.S., Cambridge University Press, p. 66-96.
- Hua X., Huss G. R., Tachibana S. and Sharp T. G. 2005: Oxygen, silicon, and Mn-Cr isotopes of fayalite in the Kaba oxidized CV3 chondrite: Constraints for its formation history. *Geochimica et Cosmochimica Acta* 69: 1333-1348.
- Pascucci I. and Tachibana Sh. 2010: The Clearing of Protoplanetary Disks and of the Protosolar Nebula. In *Protoplanetary Dust: Astrophysical and Cosmochemical Perspectives*, Eds.: Apai D. and Lauretta D.S., Cambridge University Press, p. 263-298
- Sz?cs L., Apai D., Pascucci I. and Dullemond C.P. 2010: Stellar-mass-dependent Disk Structure in Coeval Planet-forming Disks. *The Astrophysical Journal* 720: 1668-1673.

Keywords: meteorite, Early Solar System, CAI, astromineralogy