

分子動力学シミュレーションにおける気相からの核生成過程：付着確率の効果 Nucleation process from vapor due to molecular dynamics simulations: effect of the sticking probability

田中 今日子^{1*}, 河野 明男², 田中 秀和¹

TANAKA, Kyoko^{1*}, KAWANO, Akio², TANAKA, Hidekazu¹

¹ 北海道大学低温科学研究所, ² 独立行政法人海洋研究開発機構 地球内部ダイナミクス領域

¹Institute of Low Temperature Science, Hokkaido University, ²Institute For Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology

In the previous studies (Tanaka et al. 2005, 2011), we performed molecular dynamics (MD) simulations of nucleation from vapor of Lennard-Jones (L-J) type molecules and found that the semi-phenomenological (SP) model reproduces very well the nucleation rates obtained from the MD simulations. In this study, we performed MD simulations of nucleation from vapor for systems of 4000 water molecules to test nucleation theories. Simulations were done for wide ranges of the initial supersaturation ratio ($S=4-400$) and temperature ($T=250-375$ K). Through comparison with the nucleation rates and the cluster size distributions obtained from our MD simulations, we investigated the validity of the SP model. Our results show that the semi-phenomenological model reproduces well the size distributions of the clusters and the nucleation rates. Furthermore, the sticking probability of vapor molecules onto clusters was examined in MD simulations, by observing the growth rate of stable clusters larger than the critical size. In all runs in the present study, the values of the sticking probability are larger than 0.1. Our results show that the obtained sticking probability depends on the supersaturation ratio.

K. K. Tanaka, K. Kawamura, H. Tanaka, and K. Nakazawa, J. Chem. Phys. 122, 184514, 2005

K. K. Tanaka, H. Tanaka, T. Yamamoto, and K. Kawamura, J. Chem. Phys. 134, 204313, 2011

キーワード: 核生成過程, 水, 凝縮, MD シミュレーション

Keywords: nucleation process, water, condensation, MD simulation