

54 クロムの異常と隕石母天体の集積年代 Cr-54 anomalies and accretion ages of meteorite parent bodies

杉浦 直治^{1*}, 藤谷 渉¹
SUGIURA, naoji^{1*}, FUJIYA Wataru¹

¹ 東京大学理学系研究科

¹Department of Earth and Planetary Sci. University of Tokyo, Tokyo, Japan

A positive correlation between ⁵⁴Cr excesses and accretion ages is observed among meteorites including iron meteorites, palasites, mesosiderites, aubrites, HED meteorites, angrites, ureilites, acapulcoites and chondrites (including E, O, R, CK, CO, CV, CH-CB, CR, CM and CI) [1]. This suggests that ⁵⁴Cr carriers were injected into the forming solar nebula. We could constrain the solar system evolution based on this observation. However, there are still many unsettled issues concerning the ⁵⁴Cr anomalies, the accretion ages and the interpretation of the correlation. Here, we examine some of the most important issues.

26Al Heterogeneity: Homogeneous distribution of ²⁶Al is assumed for calculating accretion ages of chondrites parent bodies. It is also assumed for estimating accretion ages of differentiated meteorite parent bodies. But, at present heterogeneous distribution of ²⁶Al [2] cannot be ruled out. Comparison of precisely determined Al-Mg ages and other ages is needed to solve this problem.

Exceptions: The NWA011 group (basaltic achondrites) and Tafassasset (primitive achondrite) do not fit the correlation. They both have high ⁵⁴Cr excesses [3,4] similar to that of CR chondrites and yet apparently formed early when there was enough ²⁶Al. A possible explanation may be that early-formed planetesimals in the terrestrial-planet formation region were gravitationally scattered into the far end of the asteroidal belt, capturing CR-like materials. This is an ad hoc explanation but is shown to be possible by numerical simulations [5].

CAIs: CAIs have ⁵⁴Cr, ⁵⁰Ti and ⁴⁸Ca isotope anomalies which are larger than those found in bulk meteorites. ⁵⁴Cr and ⁵⁰Ti anomalies in CAIs and bulk meteorites appear to be well correlated with each other [6] but ⁴⁸Ca anomalies are not so well correlated with them [7]. Since CAIs formed early, they do not fit the trend formed by various meteorites on the ⁵⁴Cr vs. accretion age diagram. If we consider that the isotope anomalies of neutron-rich isotopes in CAIs and bulk meteorites originated from a similar source, then, a kind of chemical fractionation that enriched carriers of the neutron-rich isotopes must have operated during CAI formation. Otherwise, the anomalies in CAIs may have originated from a totally different source.

Other issues such as the way to estimate accretion ages of differentiated meteorite parent bodies will also be discussed at the meeting.

References

[1] Sugiura N. and Fujiya W. 2011. MAPS. 46. abstract. 5105.pdf. [2] Larsen K. K. et al. 2011. *Astrophys. J. Lett.* 735. L37-L43. [3] Bogdanovski O. and Lugmair G.W. 2004. 35th LPSC Abstract #1715. [4] Gopel C. et al. 2009. MAPS. 44. abstract 5267.pdf. [5] Bottke W. F. et al. 2006. *Nature* 439. 821-824. [6] Trinquier A. et al. 2009. *Sci.* 324. 374-376. [7] Chen H-W. et al. 2011. *Astrophys. J. Lett.* 743. L23.

キーワード: 54 クロム, 集積年代, 隕石母天体

Keywords: ⁵⁴Cr, accretion age, meteorite parent bodies