

東南極セール・ロンダーネ山地におけるリソスフェアの進化 Evolution of continental lithosphere in the Sor Rondane Mountains, East Antarctica

大和田 正明^{1*}, 亀井淳志², 堀江憲路³, 馬場壮太郎⁴, 小山内康人⁵

OWADA, Masaaki^{1*}, KAMEI, Atsushi², HORIE, Kenji³, BABA, Sotaro⁴, OSANAI, Yasuhito⁵

¹ 山口大学大学院理工学研究科, ² 島根大学総合理工学部, ³ 国立極地研究所, ⁴ 琉球大学教育学部, ⁵ 九州大学比較文化研究院

¹ Yamaguchi University, ² Shimane University, ³ National Institute of Polar Research, ⁴ University of Ryukyus, ⁵ Kyushu University

The Sor Rondane Mountains is situated within the collision zone between the West and East Gondwana and the time of collision is regarded as the late Proterozoic (650-600 Ma). The mountains consist of greenschist- to granulite-facies metamorphic rocks and various kinds of intrusive rocks. The tonalite complex is exposed in the southern part of the mountains and its magmatic age is considered to be at the middle Proterozoic (990-920 Ma). This tonalite would originally be formed at the subduction related tectonic setting prior to the collision event. Large amounts of microgabbro occur as mafic magmatic enclaves (MMEs; 990 Ma) and dikes (950 Ma) in the tonalite complex. Unmetamorphosed lamprophyre dikes intrude the tonalite complex and gneisses during the late- to post-collisional stages. The intrusive age of the lamprophyre is of 560 Ma. The magma processes of the tonalite complex together with the late- to post collisional lamprophyre dikes, therefore, provide us useful information of the evolution of continental lithosphere during the formation of Gondwana supercontinent.

The microgabbro represents the low-K and tholeiitic series, and is geochemically classified into Low-Ti and High-Ti microgabbros. The MMEs and dikes of microgabbros are equivalent to the Low-Ti and the High-Ti microgabbros, respectively. The Low-Ti and High-Ti microgabbros show geochemical signature similar to the Oceanic Arc Basalts and the Back-Arc Basin Basalts, respectively. The middle Proterozoic magma processes would, therefore, proceed at a subduction zone with back arc spreading in an oceanic arc environment. The lamprophyre corresponds to alkaline rocks in the TAS diagram, and are characterized by high abundances of LILE elements and REE, especially Rb, Ba, Sr and LREE. The trace element abundances normalized to primitive mantle display enrichment of LILE and depression of HFSE with Nb and Ta negative anomalies. The lamprophyre is plotted in the within-plate field and a part of the island arc field that is close to the within-plate field on some discrimination diagrams. Considering the geochemical features, the lamprophyre was formed in a within-plate tectonic setting by the mixing of subduction-related materials. The initial Sr isotopic ratios (SrI) range from 0.7022 to 0.7040 (epsilon SrI = -14 to 12) for the Low-Ti microgabbro and from 0.7024 to 0.7030 (epsilon SrI = -14 to 1) for the High-Ti microgabbro. The initial epsilon Nd values for the Low- and High-Ti microgabbros are calculated within the same range (epsilon NdI = -0.1 to +0.5). On the other hand, the isotopic compositions of the lamprophyre show SrI = 0.7043 - 0.7044 (epsilon SrI = +7.6 to +9.2) and epsilon NdI = -0.62 to -0.34.

The geochemical studies including Sr-Nd compositions reveal that the microgabbros have been originated from a depleted source, whereas chemical compositions of the lamprophyre is more enrichment rather than those of the microgabbros. Consequently, the magma processes in the Sor Rondane Mountains reflect the evolution of lithosphere from the middle Proterozoic to the early Paleozoic; the depleted mantle at the initial subduction stage then changing to the enriched mantle at the continental collision stage. This lithospheric evolution can be explained by interaction between the depleted mantle and the enriched materials (e.g., slab-derived fluids, melting product of subducted crustal rocks, or reaction with fossil wedge mantle) during closure of the Mozambique Ocean.

キーワード: ゴンドワナ超大陸, 南極, セール・ロンダーネ山地, トーナル岩, 細粒斑れい岩, ランプロファイアー

Keywords: Gondwana supercontinent, Antarctica, Sor Rondane Mountains, tonalite, microgabbro, lamprophyre