

日本列島に分布する熱水性鉱床のPb同位体比組成 Pb isotopic compositions of hydrothermal deposits in the Japanese island arc

藤永 公一郎^{1*}, 加藤 泰浩¹, 稗田 裕樹¹, 高谷 雄太郎¹, 谷水 雅治², 清水 徹⁴, 中村 仁美³, 岩森 光³
FUJINAGA, Koichiro^{1*}, KATO, Yasuhiro¹, HIEDA, Yuki¹, TAKAYA, Yutarō¹, Masaharu Tanimizu², SHIMIZU, Toru⁴, NAKA-MURA, Hitomi³, IWAMORI, Hikaru³

¹ 東京大学大学院工学系研究科, ² 独立行政法人海洋研究開発機構, ³ 東京工業大学大学院理工学研究科, ⁴ 産業技術総合研究所

¹University of Tokyo, ²JAMSTEC, ³Tokyo Institute of Technology, ⁴Geological Survey of Japan

Quite recently, it has been pointed out that "geofluids" released from the subducting plates may be involved in various phenomena in subduction zone, such as young volcanic rocks, deep-seated hot springs and hydrothermal deposits. Systematical investigations of these various materials are needed for identifying the geochemical characteristics of the geofluids. Nakamura et al. (2008) revealed that the slab-fluids derived from two subducted plates (the Pacific plate and the Philippine Sea plate) contribute largely to the genesis of arc magmas in the Central Japan. Here we focus on hydrothermal deposits (vein-type and skarn-type) in the Japanese island arc. Hydrothermal fluids that formed sulphide mineral (galena, pyrite, chalcopyrite, sphalerite etc.) deposits are generally considered to have been derived from magmatic and/or meteoric waters based on H, C, O, and S isotopes in the deposit materials. However, ore fluids may be derived from deep slab-fluids. We report Pb isotopic compositions of hydrothermal deposits in the Japanese island arc and discuss about the origin of ore fluids.

Keywords: Pb isotopic composition, hydrothermal deposit, slab-fluid