

一ノ目淵のかんらん岩捕獲岩中のClを含むCO₂-H₂O流体包有物 Cl-bearing CO₂-H₂O fluid-inclusions of peridotite xenoliths from Ichinomegata

熊谷 仁孝^{1*}, 川本 竜彦¹, 山本 順司¹

KUMAGAI, yoshitaka^{1*}, Tatsuhiko Kawamoto¹, Junji Yamamoto¹

¹ 京都大学 理学研究科 地球熱学

¹Inst. Geothermal Sci., Kyoto Univ.

Hydrous minerals in a subducting slab carry OH- and H₂O into the Earth's interior, and at points beyond their stability conditions they release H₂O to the overriding mantle wedge (Tatsumi and Eggin 1995). The H₂O fluids transport materials from the slab to the mantle wedge. Recently, analyses of halogen elements of high-pressure metamorphic rocks suggest that saline fluids are preserved in the subducting slab as marine pore-fluids until the depths of at least 100 km (Sumino et al., 2010, EPSL). Salinity of H₂O fluids affects dissolution properties of metal ions (Keppler, 1996, Nature). It is, therefore, important to understand the salinity of the H₂O fluids in the mantle wedge in terms of subduction system of metal.

Fluid inclusions in mantle xenoliths preserve direct information of the fluids in the mantle. Mantle xenoliths from the Ichinomegata volcano, located in back-arc side in the northeast Japan arc, have CO₂-H₂O fluid inclusions (Roedder, 1965, Am Mineral). In the present study, we report salinity of the CO₂-H₂O fluid inclusions in the mantle xenoliths from the Ichinomegata volcano.

All mantle xenoliths studied are porphyroclastic lherzolite, composed of olivine, orthopyroxene, clinopyroxene, spinel and hornblende. The CO₂-H₂O fluid inclusions are occasionally present in orthopyroxene porphyroclasts. The fluid inclusions have not reacted with host orthopyroxene crystals after the formation. We suppose, therefore, that the salinity of the fluid inclusions represents the original value in the mantle. Formation depths of the fluid inclusions are estimated by the following steps: (1) estimating the bulk mole volume of CO₂-H₂O fluid inclusion using homogenization temperatures of CO₂ liquid-vapor and CO₂-H₂O (Bakker and Diamond, 2000, *Geochem. Cosmochim. Acta*), (2) calculating pressure of the formation of the fluid inclusion using equilibrium temperature estimated by a pyroxene geothermometer (Wells 1977, *Contrib. Mineral. Petrol.*) and isochore of CO₂-H₂O system (Loner AP, from Software Package FLUIDS, v.2, Bakker), (3) converting the pressure to depth by assuming densities of crust and mantle are 2.85 and 3.3 g/cm³, respectively, and Mohorovicic discontinuity is 27 km. Salinities of fluid inclusions are determined using melting temperature of clathrate (Darling, 1991, *Geochem. Cosmochim. Acta*). The depth is estimated to be about 30 km, which is consistent with the following petrographical feature. Some xenoliths have plagioclase and symplectites formed by reaction of plagioclase and olivine. This indicates that the xenoliths were from the boundary between plagioclase-peridotite and spinel-peridotite. The salinity of fluid inclusions is 3.93 ± 0.55 wt %. Using relationship between the molality of Cl and the fluid/melt partition coefficients (Zajacz et al., 2008, *Geochem. Cosmochim. Acta*), for example, the fluid/melt partition coefficients of Pb and Zn under this salinity are 7.8 and 18.6, respectively (those of Cl-free hydrous fluid are almost 0 and 8.2, respectively).

キーワード: 塩濃度, 流体包有物, 物質輸送, 沈み込み帯, マントル捕獲岩, 一ノ目淵

Keywords: salinity, fluid inclusion, material transport, subduction zone, mantle xenolith, Ichinomegata