Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



SCG69-04

会場:102B

時間:5月24日09:45-10:00

A statistical study of ULF seismo-magnetic phenomena in Kanto, Japan during 2000-2010

A statistical study of ULF seismo-magnetic phenomena in Kanto, Japan during 2000-2010

Peng Han^{1*}, Katsumi Hattori¹, Febty Febriani¹, Yuki Ishiguro¹, Chie Yoshino¹ HAN, Peng^{1*}, Katsumi Hattori¹, Febty Febriani¹, Yuki Ishiguro¹, Chie Yoshino¹

Recently electromagnetic phenomena have been considered as a promising candidate for short-term earthquake prediction. And especially passive ground-based observation of ULF (ultra low frequency) geomagnetic signatures is considered to be the most promising method for seismo-magnetic phenomena study due to deeper skin depth. In order to clarify the earthquake-related ULF magnetic phenomena, a sensitive geomagnetic network has been installed in Japan and plenty of data associated with moderate-large earthquakes have been accumulated. In this study, we have analyzed geomagnetic data observed during the past decade in Kanto area, Japan.

First, the ULF magnetic signals at the frequency 0.01Hz have been investigated. We have applied wavelet transform analysis to the 1Hz sampling data observed at three magnetic observatories in Boso Peninsula (Kiyosumi, Uchiura, and Fudago) and Izu Peninsula (Seikoshi, Mochikoshi, and Kamo), respectively. The signature at the 0.01Hz frequency band has been revealed and daily average energy has been computed. In order to minimum artificial noise, we only use the midnight time data (LT 1:00~4:00). And to remove influences of global magnetic perturbations, we have developed another method to obtain reliable background based on principal component analysis (PCA). Three standard geomagnetic stations (Memambetsu, Kakioka, and Kanoya) operated by the Japan Meteorological Agency have been selected as reference stations and PCA method has been applied to the yearly energy variation of the 0.01Hz signals at the three stations. The first principal component which contains more than 95% energy is considered to be global background.

After comparing the results at the stations in Boso and Izu Peninsula with global background, it is found that there are several local energy enhancements which only appear in Boso or Izu area. Especially for the case studies of the 2000 Izu Island earthquake swarm and the 2005 Boso M6.1 earthquake, significant anomalous behaviors have been detected in Z components.

Finally, we have applied superposed epoch analysis (SEA) to the above results and make a statistical study. The statistical results have indicated that before an earthquake there are clearly larger probabilities of anomalies than that after the earthquake. For Izu area, three weeks and few days before statistical value of anomalies is significant; for Boso region, around ten and few days before it is significant.

キーワード: ULF seismo-magnetic phenomena, statistical study Keywords: ULF seismo-magnetic phenomena, statistical study

¹Graduate School of Science, Chiba University

¹Graduate School of Science, Chiba University