

## 貫入岩体周辺の面構造ダイヤグラムに基づく東ピルバラの花崗岩複合岩体の貫入様式について

### Type of granite complex intrusion in East Pilbara terrane based on foliation pattern diagram around intrusion body

松村 太郎次郎<sup>1\*</sup>, 増田 俊明<sup>2</sup>

MATUMURA, Taroujiro<sup>1\*</sup>, MASUDA, Toshiaki<sup>2</sup>

<sup>1</sup> 静岡大学大学院理学研究科, <sup>2</sup> 静岡大学理学部

<sup>1</sup> Graduate School of Science, Shizuoka university, <sup>2</sup> Faculty of Science, Shizuoka University

Theoretical analysis of stressed two-dimensional elastic material with a circular hole filled with a viscous material revealed that three types (TT, TX and XX types) of distribution pattern of maximum principal stress orientation around the hole can be produced as a function of far-field stress  $S_1$ ,  $S_2$  and internal pressure of viscous material  $p$ . The TT type is characterized by tangential orientation of  $S_2$  axis in all directions around the hole, while the TH is characterized by  $S_2$  axis of tangential and normal orientations to the circular hole in orthogonal orientations. The XX type is characterized by normal orientation of foliation all around the hole. Assuming that  $S_2$  axis is parallel to the foliation, we consider the stress state for actual foliation patterns in aureoles of granitic intrusions. We can find some natural examples of TT and TX type patterns, whereas we have never found the XX type patterns in nature. In this poster we present TT a type foliation pattern around the Mount Edgar Batholith in Pilbara area, Western Australia, and discuss how the TT type pattern foliation can be produced as a function of internal pressure of granitic body.

キーワード: ピルバラ, バソリス, 太古代, 貫入イベント

Keywords: Pilbara, batholith, Archean, intrusion event