

スタグナントスラブの融解：中国東部新生代アルカリ玄武岩からの制約 Melting of a stagnant slab in the mantle transition zone: Constraints from Cenozoic alkaline basalts in eastern China

柵山 徹也^{1*}, Tian Wei², 木村 純一¹, 深尾 良夫¹, 平原 由香¹, 高橋 俊郎¹, 仙田 量子¹, 常 青¹, 宮崎 隆¹, 大林 政行¹, 川畑 博¹, 巽 好幸¹
SAKUYAMA, Tetsuya^{1*}, Tian Wei², KIMURA, Jun-Ichi¹, FUKAO, Yoshio¹, HIRAHARA, Yuka¹, TAKAHASHI, Toshiro¹, SENDA, Ryoko¹, CHANG, Qing¹, MIYAZAKI, Takashi¹, OBAYASHI, Masayuki¹, KAWABATA, Hiroshi¹, TATSUMI, Yoshiyuki¹

¹ 海洋研究開発機構, ² 北京大学

¹JAMSTEC, ²Peking Univ.

The feasibility of the melting of oceanic igneous crust in stagnant slabs has been proposed by studies on experimental petrology, however, relevant geochemical evidence of melting has not yet been found from igneous rocks. We present evidence that proves that melts from the igneous layer in the stagnant Pacific slab have contributed to the source composition of basalts from eruption in eastern China. Fe-rich (>13 wt%), Si-poor (<43 wt%) basalts only occur above the leading edge of the stagnant Pacific slab in eastern China. Their source has Nd-Hf isotope compositions akin to the igneous layer in the Pacific slab, while they have Sr-Nd-Pb isotope compositions similar to those of mid-oceanic-ridge basalt. The extremely low Rb and Pb (Ce/Pb > 30) contents of these basalts suggest that this source material was modified by a subduction process. Together, these geochemical characteristics help us to conclude that these basalts have received a significant contribution from the melts derived from dehydrated carbonate-bearing oceanic crust, without a long time-integrated ingrowth of Sr-Nd-Hf-Pb isotope systems at the leading edge of the stagnant Pacific slab.

キーワード: プレート内アルカリ玄武岩, 中国東部, 山東半島, HIMU 玄武岩, スタグナント太平洋スラブ, 海洋地殻リサイクル

Keywords: Intraplate alkaline basalt, eastern China, Shandong Peninsula, HIMU basalt, stagnant Pacific slab, oceanic crust recycling