

X線吸収法を用いた高温高圧下における Fe-O 融体の密度測定 Density measurement of liquid Fe-O at high pressure and high temperature using an X-ray absorption method

田窪 勇作^{1*}, 下山 裕太², 寺崎 英紀¹, 浦川 啓³, 西田 圭佑², 鹿室 僚太³, 岸本 俊八³, 近藤 忠¹, 大谷 栄治², 片山 芳則⁴
TAKUBO, Yusaku^{1*}, SHIMOYAMA, Yuta², TERASAKI, Hidenori¹, URAKAWA, Satoru³, NISHIDA, Keisuke², KAMURO, Ryota³, Shunpachi Kishimoto³, KONDO, Tadashi¹, OHTANI, Eiji², KATAYAMA, Yoshinori⁴

¹ 大阪大学, ² 東北大学, ³ 岡山大学, ⁴ 日本原子力研究開発機構

¹Osaka University, ²Tohoku University, ³Okayama University, ⁴Japan Atomic Energy Agency

The Earth's outer core is thought to be composed of liquid iron alloys with a small amount of light elements, such as sulfur, oxygen and silicon. Existence of a liquid core is also suggested to other terrestrial planets (Mars and Mercury). Thus the effect of light elements on the density of liquid iron is fundamental to understand the composition and structure of the planetary cores.

The densities of liquid Fe-S, Fe-Si, and Fe-C have been reported using X-ray absorption method (Nishida et al., 2011; Sanloup et al., 2004; Terasaki et al., 2010). As a result, it was revealed that the rate of density decrease is quite different depending on the dissolving light element. Hence, it is important to figure out the effects on liquid iron by individual light elements. Although oxygen is one of the most popular candidates of the light elements in the Earth's outer core, the effect of oxygen on the density of liquid iron has never been reported to date. In this study, we have measured the density of liquid Fe-O (O = 0.5 wt%) up to 3 GPa and 2250 K using X-ray absorption method at BL22XU, SPring-8 synchrotron facility. The obtained density of this study is 6.65(3) g/cm³ at 3 GPa and 2005 K. Compared to the density of pure liquid iron at the present experimental condition, the density of liquid Fe-O is about 7% smaller than that of liquid iron and thermal expansion coefficient of liquid Fe-O is similar to that of liquid iron.

キーワード: 地球核, 酸素, 密度, 高温高圧, 放射光

Keywords: core, oxygen, density, high pressure and high temperature, synchrotron