

## モンゴル中央部, バヤンホンゴル地域に産出する変成岩類の岩石学的・年代学的解析

### Petrology and geochronology of metamorphic rocks in the Bayankhongor area, the central part of Mongolia.

足立 達朗<sup>1\*</sup>, 小山内 康人<sup>1</sup>, 中野 伸彦<sup>1</sup>, 大和田 正明<sup>2</sup>, Madhusoodhan Satish-Kumar<sup>3</sup>, Sereenen Jargalan<sup>4</sup>, Boldbaatar Chimedtseye<sup>5</sup>, 米村 和紘<sup>1</sup>, 吉本 紋<sup>1</sup>

ADACHI, Tatsuro<sup>1\*</sup>, OSANAI, Yasuhito<sup>1</sup>, NAKANO, Nobuhiko<sup>1</sup>, OWADA, Masaaki<sup>2</sup>, MADHUSOODHAN, Satish-Kumar<sup>3</sup>, SEREENEN, Jargalan<sup>4</sup>, Chimedtseye Boldbaatar<sup>5</sup>, Kazuhiro Yonemura<sup>1</sup>, Aya Yoshimoto<sup>1</sup>

<sup>1</sup>九州大学, <sup>2</sup>山口大学, <sup>3</sup>静岡大学, <sup>4</sup>モンゴル科学技術大学, <sup>5</sup>モンゴル鉱物資源石油管理庁

<sup>1</sup>Kyushu University, <sup>2</sup>Yamaguchi University, <sup>3</sup>Shizuoka University, <sup>4</sup>Mongolian Univ. Sci. Tech, <sup>5</sup>Miner. Res. Auth. Mongolia

Mongolia is situated in the Central Asian Orogenic Belt (e.g. Mossakovsky et al., 1993), which extends from the Siberian craton, Baltica craton, North China craton and Tarim craton. It is composed of subduction-accretion complexes and developed from c. 1000 Ma to c. 250 Ma (Windley et al., 2007). Metamorphic rocks are partly distributed in the central and western part of Mongolia. Metamorphic history would be a key for understanding the tectonic evolution of this area, however detailed study on metamorphic rocks has not been well-documented, including geochronology. In this study, we report petrology and U-Pb monazite and zircon ages on metamorphic rocks occurring in the Bayankhongor area.

The Bayankhongor area is situated between Gobi-Altai range and Hangay range in the central part of Mongolia. The area is mainly composed of granitic gneisses (biotite gneiss and clinopyroxene-biotite gneiss), amphibolites and pelitic gneisses. Pelitic gneisses are classified into garnet-biotite gneiss, garnet-sillimanite-biotite gneiss and garnet-cordierite-biotite gneiss. Garnet in garnet-cordierite-biotite gneiss is replaced by cordierite and biotite, indicating decompression process. Sillimanite in garnet-sillimanite-biotite gneiss is replaced by andalusite, which also indicates later stage metamorphism under the lower pressure condition. Amphibolites are classified into garnet-amphibolite, clinopyroxene-amphibolite and amphibolite. Garnet in garnet-amphibolite is replaced by plagioclase and hornblende, which qualitatively imply decompression as well as pelitic gneisses. Thus, metamorphic rocks in the Bayankhongor area widely recorded amphibolite- to upper amphibolite-facies metamorphism with subsequent decompression.

In the presentation, we show additional data including mineral chemistry, P-T estimations and U-Th-Pb in situ dating and discuss metamorphic evolution of the Bayankhongor area. Based on the interpretation of those data, we suggest its implication for the tectonic process related to the development of the Asia continent.

#### References

Mossakovsky et al. (1993) Central Asian fold belt: geodynamic evolution and history of formation. *Geotectonics*, 6, 3-33.

Nakano et al. (2010) Metamorphic rocks from the Mongol-Altai: A first report of Japan-Mongolia Joint Geological Research (2009-2013). *Journal of Geological Society of Japan*, XI-XII.

Windley et al. (2007) Tectonic models for accretion of the Central Asian Orogenic Belt. *Journal of the Geological Society*, London, 164, 31-47.

キーワード: 変成岩, 温度圧力経路, ウラン 鉛年代, バヤンホンゴル地域, モンゴル

Keywords: metamorphic rock, P-T condition, U-Pb dating, Bayankhongor Area, Mongolia