

## 第一原理電子状態計算による含水リングウッドイトの振動特性 Vibrational properties of hydrous ringwoodite, first principles investigation

土屋 旬<sup>1\*</sup>

TSUCHIYA, Jun<sup>1\*</sup>

<sup>1</sup> 愛媛大学上級研究員センター

<sup>1</sup>Senior Research Fellow Center, Ehime University

Wadsleyite and ringwoodite are primary constituent minerals in the Earth's transition zone. These phases can contain up to a few wt% H<sub>2</sub>O in the crystal structure and are thought to be the most important water reservoirs in the Earth. We have investigated the high pressure protonation sites in hydrous wadsleyite using first principles calculation and found that the oxygen O<sub>1</sub> site is the most favorable for protonation in wadsleyite because of the electric imbalance of this site. On the other hand, the crystal structure of ringwoodite does not have such peculiar protonation sites and the reason of such high retention of water in ringwoodite has been unclear so far. In present study, I have calculated the vibrational properties of hydrous ringwoodite under pressure with various protonation models by first principles. Comparing with the IR and Raman measurements, I will discuss the protonation sites in hydrous ringwoodite.

Research supported in part by special coordination funds for promoting science and technology (Supporting Young Researchers with Fixed-term Appointments) and Grants-In-Aid for Scientific Research from the Japan Society for the Promotion of Science 20103005 and 21740380.

キーワード: 含水リングウッドイト, 第一原理計算, 振動特性

Keywords: hydrous ringwoodite, first principles, vibrational property