

## 化石分岐断層から得られた連続的コア・検層データ：延岡衝上断層掘削速報 Continuous coring and logging dataset obtained from fossilized megasplay fault

山口飛鳥<sup>1\*</sup>, 木村学<sup>1</sup>, 浜橋真理<sup>1</sup>, 福地里菜<sup>2</sup>, 亀田純<sup>1</sup>, 濱田洋平<sup>1</sup>, 藤本光一郎<sup>2</sup>, 橋本善孝<sup>3</sup>, 比名祥子<sup>1</sup>, 栄田美緒<sup>3</sup>, 斎藤実篤<sup>4</sup>, 北村有迅<sup>4</sup>, 水落幸広<sup>5</sup>  
YAMAGUCHI, Asuka<sup>1\*</sup>, KIMURA, Gaku<sup>1</sup>, HAMAHASHI, Mari<sup>1</sup>, FUKUCHI, Rina<sup>2</sup>, KAMEDA, Jun<sup>1</sup>, HAMADA, Yohei<sup>1</sup>, FUJIMOTO, Koichiro<sup>2</sup>, HASHIMOTO, Yoshitaka<sup>3</sup>, HINA, Shoko<sup>1</sup>, EIDA, Mio<sup>3</sup>, SAITO, Saneatsu<sup>4</sup>, KITAMURA, Yujin<sup>4</sup>, MIZUOCHI, Yukihiko<sup>5</sup>

<sup>1</sup> 東京大学大学院理学系研究科地球惑星科学専攻, <sup>2</sup> 東京教育大学自然科学系, <sup>3</sup> 高知大学大学院理学研究科応用理学専攻, <sup>4</sup> 海洋研究開発機構地球内部ダイナミクス領域, <sup>5</sup> 住鉱資源開発株式会社  
<sup>1</sup> Department of Earth and Planetary Science, the University of Tokyo, <sup>2</sup> Faculty of Education, Tokyo Gakugei University, <sup>3</sup> Department of applied science, Faculty of Science, Kochi University, <sup>4</sup> Institute For Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology, <sup>5</sup> Sumiko Resources Exploration & Development, Co., Ltd.

Mechanics and evolution of large thrust faults along subduction plate boundaries are one of the essential topics in earth sciences because of their potential for causing catastrophic geohazards. Megasplay faults, large landward-dipping thrust fault branching from plate boundary megathrusts, are one of the candidates for the source of large tsunamis. Deep portion (~5200 mbsf) of the megasplay fault in the Nankai Trough is listed for the next drilling target of the NanTroSEIZE project. In such depths, the megasplay fault is recognized as strong reflector of seismic profiles, however, the thickness, architecture, deformation styles of the fault zone is still ambiguous. To evaluate the status of modern megasplay fault before drilling, we projected drilling, coring and logging to the Nobeoka thrust, Japan, a fossilized on-land analog of such megasplay fault and its basic setting has been constrained (Kondo et al., 2005): Nobeoka Thrust Drilling Project (NOBELL).

Drilling operation continued from July to September, 2011. Coring was operated up to 255 m depth with excellent recovery (99.82%). Visual core descriptions including detail sketch, lithological and structural characterization, measurements on 3,787 structural elements, were performed on the whole core. Subsequently, geophysical logging (temperature, spontaneous potential logging, natural gamma-ray, resistivity, P/S-wave velocity, neutron porosity, caliper) and borehole imaging (optical and ultrasonic wave) were operated and continuous dataset from 12 to 252 m-depth were obtained. Cores were stored in Kochi Core Center (KCC) at Kochi University, and gamma-ray density, magnetic susceptibility measurements were performed by a multi-sensor core logger.

Although analyses of core description and logging dataset are now in progress, various cataclasites and slip zones possibly reflecting protolith type and deformation mechanisms, and many spikes on logging data have been recognized. The results of NOBELL would provide new insights on not only the architecture but mechanics and evolution of ancient and modern megasplay faults.