

## 鮮新世北大西洋堆積物の年代決定に関わる新手法

### New method for the chronostratigraphic reconstruction of Pliocene marine sediments at the subpolar North Atlantic

林 辰弥<sup>1\*</sup>, 大野正夫<sup>2</sup>, 桑原義博<sup>2</sup>, 山下剛史<sup>2</sup>, 水田麻美<sup>2</sup>, 北逸郎<sup>2</sup>

HAYASHI, Tatsuya<sup>1\*</sup>, Masao Ohno<sup>2</sup>, Yoshihiro Kuwahara<sup>2</sup>, Takafumi Yamashita<sup>2</sup>, Asami Mizuta<sup>2</sup>, Itsuro Kita<sup>2</sup>

<sup>1</sup> 国立科学博物館, <sup>2</sup> 九州大学

<sup>1</sup>The National Museum of Nature and Science, <sup>2</sup>Kyushu University

We established a new method to reconstruct chronostratigraphy of subpolar North Atlantic sediments during the Pliocene by using a new hybrid environmental proxy. Pliocene sediments recovered at IODP Site U1314 in the Gardar Drift (North Atlantic) are characterized by the paucity of foraminiferal tests, and this hampered the establishment of an age-depth model of the sediments by the orbital tuning method using an oxygen isotope record. Therefore, we created the hybrid environmental proxy by combining magnetic susceptibility and natural gamma radiation as a tuning material. In the hybrid proxy record, glacial-interglacial climate variations are extracted and the small-scale variations (attributed to ice-rafted debris) are eliminated. We established a high-resolution age model for the sediments at IODP Site U1314 during 2.76-2.10 Ma by tuning the hybrid proxy record to the global standard oxygen isotope record (LR04 stack). We would point out the possibility that our tuning method may be applicable to other Pliocene sediments in subpolar North Atlantic regions.

キーワード: 年代層序, 鮮新世, 海底堆積物, 北大西洋, 帯磁率, 自然ガンマ線

Keywords: chronostratigraphy, Pliocene, marine sediments, North Atlantic, magnetic susceptibility, natural gamma radiation