

## 高温高圧実験による富士火山マグマ溜りの研究 Experimental study on magma plumbing system beneath Fuji volcano

浅野 健太<sup>1\*</sup>, 高橋 栄一<sup>1</sup>, 浜田 盛久<sup>1</sup>, 潮田 雅司<sup>1</sup>, 鈴木 敏弘<sup>2</sup>

ASANO, Kenta<sup>1\*</sup>, TAKAHASHI, Eiichi<sup>1</sup>, HAMADA, Morihisa<sup>1</sup>, USHIODA, Masashi<sup>1</sup>, SUZUKI, Toshihiro<sup>2</sup>

<sup>1</sup> 東京工業大学大学院理工学研究科地球惑星科学専攻, <sup>2</sup> 海洋研究開発機構 地球内部変動研究センター

<sup>1</sup>Department of Earth and Planetary Sciences, Tokyo Institute of Technology, <sup>2</sup>Institute for Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology

Fuji volcano, largest in volume and eruption rate in Japan, is located at the center of Honshu, where North America, Eurasia and Philippine Sea plates meet. Beneath Fuji volcano, subduction of both Pacific and Philippine Sea plates are ongoing. Eruption of Fuji volcano may be related to large magnitude interplate earthquakes at least in some cases. Magma chamber beneath Fuji volcano is considered to be unusually deep compared with other volcanoes in Izu-Mariana arc. Fujii (2007) interpreted that unusual depth of Fuji magma chamber is due to thickened low density granitic crust by collision of Izu peninsula. Because of the significance of Fuji volcano both in tectonic settings and potential volcanic hazard, there are a great number of studies on Fuji volcano. However, studies focused on magma plumbing system beneath Fuji volcano are limited and there are no high-pressure experiments on Fuji basalt so far. The purpose of this study is to determine the conditions of the magma chamber (P, T, fO<sub>2</sub>, etc) of Fuji volcano through high pressure melting experiments.

Basalt scoria Tr-1 which represents the final ejecta of Hoei eruption in 1707, was adopted as a starting material. Experiments at 4kbar were carried out using an internally heated pressure vessel (HIP-5000) at the Magma Factory. Temperature conditions were 1050, 1100 and 1150°C, and H<sub>2</sub>O contents were 1.3, 2.7 and 4.7wt.%, respectively. The fO<sub>2</sub> was controlled at NNO-buffer. At 4kbar, magnetite is the first liquidus phase and plagioclase is the second liquidus phase and is followed by clinopyroxene and orthopyroxene. Compositions of melts at 4 kbar were determined by EPMA analysis of quenched run products. SiO<sub>2</sub> content of melt increases with crystallization and is different from silica non-enrichment compositional trend of Fuji basalt.

In order to explain silica non-enrichment differentiation trend of Fuji volcano, Fujii(2007) suggested that orthopyroxene may play important role at the deep magma chamber. Experiments at 7 kbar are in progress using another internally heated pressure vessel (HIP-8600) at the Magma Factory. Phase relations and melt compositional trend at 7 kbar will be reported. Based on high-pressure melting experiments and petrologic study, mechanism of Hoei sub-plinian eruption and origin of the dacite which was erupted at the initial stage of Hoei eruption will be discussed.

キーワード: 富士火山, マグマ, 実験岩石学, 沈み込み帯, 高圧実験

Keywords: Fuji volcano, magma, Experimental petrology, Subduction zone, High-pressure experiment