Japan Geoscience Union Meeting 2012

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.

MIS23-05



時間:5月21日14:45-15:00

## ラマン分光分析および熱分析によるオホーツク海サハリン島沖ガスハイドレートの 結晶学的特性 Crystallographic properties of gas hydrates off Sakhalin Island (Sea of Okhotsk) using

Crystallographic properties of gas hydrates off Sakhalin Island (Sea of Okhotsk) using Raman spectroscopy & calorimetry

八久保 晶弘  $^{1*}$ , 坂上 寛敏  $^1$ , 南 尚嗣  $^1$ , 山下 聡  $^1$ , 高橋 信夫  $^1$ , 庄子 仁  $^1$ , ジン ヤン  $^2$ , オブジロフ アナトリー  $^3$ , ヴェレシャー ギナ オルガ  $^3$ 

HACHIKUBO, Akihiro<sup>1\*</sup>, Hirotoshi Sakagami<sup>1</sup>, MINAMI, Hirotsugu<sup>1</sup>, YAMASHITA, Satoshi<sup>1</sup>, TAKAHASHI, Nobuo<sup>1</sup>, Hitoshi Shoji<sup>1</sup>, Young K. Jin<sup>2</sup>, Anatoly Obshirov<sup>3</sup>, Olga Vereshchagina<sup>3</sup>

<sup>1</sup> 北見工業大学,<sup>2</sup> 韓国極地研究所,<sup>3</sup>V.I. イリチェフ太平洋海洋学研究所

<sup>1</sup>Kitami Institute of Technology, <sup>2</sup>Korea Polar Research Institute, <sup>3</sup>Pacific Oceanological Institute, FEB RAS

Natural gas hydrates have been studied in the gas seep sites off Sakhalin Island, the Sea of Okhotsk. More than ten gas seep sites in the north area of the Lavrentyev seabed Fault (LVF) form gas hydrates in the sea-bottom sediments. In the framework of SSGH (Sakhalin Slope Gas Hydrate) project, hydrate-bearing sediments are also discovered in the south area of LVF, where more than 400 acoustic anomaries exist but the size of seepage structures are relatively small. We obtained samples of natural gas hydrate in the CHAOS project (2003-2006) and the SSGH project (2007-2011) and analyzed their crystallographic properties using a calorimeter and a Raman spectrometer. Their gas composition were mainly methane, suggested the cubic structure I of these crystals. The hydration number, cage occupancies of both large and small cages, and dissociation heat of the sample were almost same as those of pure methane hydrate. Raman spectra showed that hydrogen sulfide were encaged in both large and small cages. The peak ratio of large to small cages were about 2.4 and 3.5 for hydrogen sulfide and methane, respectively.

キーワード: ガスハイドレート, 解離熱, ケージ占有率, 水和数, オホーツク海 Keywords: gas hydrate, dissociation heat, cage occupancy, hydration number, Sea of Okhotsk