## **Japan Geoscience Union Meeting 2012**

(May 20-25 2012 at Makuhari, Chiba, Japan)

©2012. Japan Geoscience Union. All Rights Reserved.



MTT06-05 Room:301B Time:May 24 10:00-10:20

## Geomagnetic detection of the sectorial solar magnetic field and the historical peculiarity of minimum 23-24

LOVE, Jeffrey<sup>1\*</sup>, E. Joshua Rigler<sup>1</sup>, Sarah E. Gibson<sup>2</sup>

Analysis is made of the geomagnetic-activity as index covering solar cycle 11 to the beginning of 24, 1868-2011. Autocorrelation shows 27.0-d recurrent geomagnetic activity that is well-known to be prominent during solar-cycle minima; some minima also exhibit a smaller amount of 13.5-d recurrence. Previous work has shown that the recent solar minimum 23-24 exhibited 9.0 and 6.7-d recurrence in geomagnetic and heliospheric data, but those recurrence intervals were not prominently present during the preceeding minima 21-22 and 22-23. Using annual-averages and solar-cycle averages of autocorrelations of the historical as data, we put these observations into a long-term perspective: none of the 12 minima preceeding 23-24 exhibited prominent 9.0 and 6.7-d as recurrence. We show that the detection of these recurrence intervals can be traced to an unusual combination of sectorial spherical-harmonic structure in the solar magnetic field and anomalously low sunspot number. We speculate that 9.0 and 6.7-d recurrence is related to transient large-scale, low-latitude organization of the solar dynamo, such as seen in some numerical simulations.

Keywords: Geomagnetism, Magnetic observatory, Recurrent geomagnetic activity, Solar-terrestrial interaction, Solar wind, Solar dynamo

<sup>&</sup>lt;sup>1</sup>USGS Geomagnetism Program, <sup>2</sup>NCAR High Altitutde Observatory