

Carbon dioxide dynamics in coastal regions of Osaka Bay

FUJII, Tomoyasu^{1*}, FUJIWARA Tateki², KOMAI Yukio³

¹Nara University of Education, ²Kyoto University, ³Osaka Institute of Technology

Measurement technique of CO₂ for coastal seas is developed and applied to elucidate CO₂ dynamics in the coastal regions, where the photosynthetic rate is far larger than that in the open sea and the short-term change is significant. Continuous measurements of salinity, pH and DO were conducted at three stations in Osaka Bay. The values of CO₂ related terms were calculated using a classical method that uses pH and total alkalinity. Dissolved inorganic carbon (DIC) and DO fluctuated with high correlation ($R^2 = 0.97$). This suggests that CO₂ system can be measured by this method in coastal regions. DO and pCO₂ (CO₂ partial pressure) records in the eastern Osaka Bay, where primary production is significant, indicated prominent diurnal variations which correspond to diurnal irradiation variations. In contrast, magnitudes of DO and pCO₂ variations were smaller in a well mixed region in the western Osaka Bay.

Sediment phosphorus content and its potential release in the coastal area of Seto Inland Sea Japan

JIN, Guangzhe^{1*}, ONODERA, Shin-ichi¹, AMANO, Atsuko², SHIMIZU, Yuta¹, SATO, Takaharu¹, JIGE, Mayumi³

¹Hiroshima University, faculty of Integrated arts and science, ²Geological Survey of Japan, AIST, ³Chiba Institute of Science Faculty of Risk and Crisis Management

Eutrophication is an important world-wide problem and became a heated debate recent years. In many coastal sea areas around the world, such as Tokyo bay and Baltic Sea, the Phosphorus (P) plays a key role in this process; the Kojima bay is located in Okayama prefecture and is an important water flow to the Seto inland sea. The P load to the Seto inland sea appears to have an important effect on the eutrophication in this area. Kojima Lake is formed by enclosing the dike in 1959, so research on the effect of P formation on the environment is important and interesting. Our study is mainly focused on the effect of phosphorus in sediment and the overlying water samples in Kojima bay and Kojima Lake.

Surface and core sediment samples were collected both in Kojima bay and Kojima Lake in this study. The surface sediment samples were collected by box sampler, the core samples in Kojima lake were taken by piston core sampler while the cores in Kojima bay were taken by diver using acrylic tubes (7-8 cm diameter). Pore water samples were also extracted by centrifuge and the nutrient in pore water, near bottom and surface water samples were determined in the laboratory with a spectrophotometer (Bltec Swaat autoanalyser). We use the ²¹⁰Pb activity and ¹³⁷Cs activity to determine the sedimentation and dating data of the core samples. In this study, a six-step extraction method of P in sediment was used to describe the chemical species of P. By dividing the P into active forms (loosely sorbed P, redox sensitive P) and immobile forms (oxide metal bound P, apatite P and residue P),

The sediment phosphorus content in surface sediment samples is higher in the lake samples (average 27 μmol/g in 7 sites) than in the bay samples (average 14 μmol/g in 20 sites), while the higher pore water samples and water samples both showed higher in bay samples than in lake. It may indicate that the higher stabilization form of phosphorus in Kojima Lake surface sediment with lower possibility of transportation in releasing to pore water and overlying water. P fractionation results show that redox sensitive P forms are the critical P forms leading to the variation of phosphorus which is related to the iron content in sediment, dissolved iron and manganese showed lower content in lake water volume. The core samples showed that phosphorus content decreased after it was deposited with the increasing of P content in pore water. The redox sensitive phosphorus content decreased sharply with the increasing of loosely sorbed phosphorus, pore water phosphorus and salinity at the down core. This represents the releasing of phosphorus content from sediment deposited with the low oxygen condition and higher salinity in the deep layer of the sediment. The relatively high salinity with pH in Kojima bay will inhibit phosphate adsorption onto Fe oxides/hydroxides. Also, the concentration of Fe oxides/hydroxides is reduced in sulfide environments by the formation of solid Fe sulfides and if sulfate-reduction rates are controlled by sulfate concentrations. This may be one of the main reasons for the variations of phosphorus content in coastal bay and lake sediment. The relatively high concentrations of dissolved P associated with riverine inputs in Kojima Bay are to some extent buffered by the relatively high concentrations of suspended sediments resulting from tidal flows. Phosphorus may be released during transport to the sea due to decreases in the active phosphorus forms, increases in salinity and release from bottom sediments as a result of low oxygen conditions.

Keywords: Sediment, Kojima Bay, Phosphorus, Fractionation, Release

Impacts of the 3D nutrient-transport by the Kuroshio on the land-sea biogeochemical interaction in the western North Pac

KOMATSU, Kosei^{1*}

¹Graduate School of Frontier Sciences, University of Tokyo, ²Atmosphere Ocean Research Institute, University of Tokyo

The Kuroshio, the western boundary current in the North Pacific, plays major roles in transporting heat and organic/inorganic materials from the subtropical region to the subarctic one, and moreover from the coastal region to the offshore one. The Kuroshio undoubtedly must impact on the ecosystem in its neighboring and downstream regions, however it is generally recognized as a mere boundary between the oligotrophic Subtropical waters to the south and the more productive coastal or subarctic waters to the north. Surprisingly neither quantitative nor qualitative researches have advanced to clarify the actual distribution of nutrients in the Kuroshio region with focus on the impacts of the jet, the core of the current maximum. The transport of nutrient and its impacts on the ecosystem have been still unknown mainly because of lack of simultaneous measurement of both horizontal and vertical fluxes of nutrients around the jet.

In order to clarify the 3D distribution of the water properties in the Kuroshio region and to estimate horizontal and vertical fluxes of nutrients and their impacts on the productivity in the surrounding and downstream regions, an intensive observation was conducted in Apr. 2009 and historical hydrographic data were analyzed. The observation was carried out by the R/V Tansei-maru at intervals of 10 miles along the 5 lines crossing the Kuroshio south of Japan. It obtained the 3D distribution of the water properties by CTD with multi-profilers and bottle-samplings, the horizontal velocity by the shipboard and lowered ADCP, and the vertical turbulent diffusivity by the microstructure profiler. As a result we detected that maxima of nitrate, silicate, phosphate and AOU were located along the jet on the isopycnal surface of 24.5-26.0sigma-theta. It is the first detection of the nutrient/AOU maximum along the Kuroshio jet, and the structure is analogous to the characteristic one well-known as Nutrient Stream found in the Gulf Stream region. Moreover, the nutrient/AOU maximum along the Kuroshio jet was also detected on the 24.5-25.5sigma-theta surface in spring in the whole region of the Kuroshio, by analyzing the historical data of JODC.

It should be emphasized that the nutrient concentration on the isopycnal surface of 24.5-25.5sigma-theta gradually decreases along the jet toward the downstream region. It implies that the high nutrient water is originated from the upstream and its adjacent coastal regions and transported downstream epipycnally along the Kuroshio as is the case with the Gulf Stream. Our observation estimated the maximum of the epipycnal nitrate flux at $10\text{mmolNm}^{-2}\text{s}^{-1}$ around the 26.0sigma-theta surface just beneath the current maximum of the Kuroshio jet. A part of the flux is served out to both the northern and southern sides of the jet due to eddy diffusivity, and especially on the northern side the nutrient transport is important for the new production under sufficient irradiance.

Moreover our observation clarified quantitatively that nutrient is supplied upward more intensively on the jet and its inshore side than the offshore side due to higher diapycnal mixing observed by direct measurement of microstructure. The diapycnal flux of nitrate amounts to $3.0\times 10^{-6}\text{mmolNm}^{-2}\text{s}^{-1}$ at the 25.0-25.5sigma-theta just above the core of the epipycnal flux, indicating that the high nutrient transported epipycnally along the jet is supplied efficiently upward by the strong diapycnal mixing and that it contributes significantly to the spring new production around the Kuroshio.

Importantly, on the northern side of the Kuroshio Extension the water mass of 25.0-25.5sigma-theta is distributed at the upper part of the euphotic layer in spring, as a result the nutrient flux via the Kuroshio jet contributes the high productivity around the region, where enhanced concentration of chlorophyll can be seen from the ocean-color map and favorable habitats are formed for various pelagic fishes.

Keywords: Kuroshio, Nutrient Stream, diapycnal flux, epipycnal flux

Dynamic response of a sporadically opened lagoon to land and sea

CHIKITA, Kazuhisa^{1*}, IWASAKA, Wataru², Kazuhiro Ohmori³, MAMUN, Abdullah²

¹Faculty of Science, Hokkaido University, ²Graduate School of Science, Hokkaido University, ³Faculty of Science, Hokkaido University

Four coastal lagoons in Tokachi, Hokkaido, are sporadically opened to the Pacific Ocean by breaking gravel bars due to overflow. One of the lagoons, Oikamanai Pond, is opened to the ocean about four times per year. The water budget of the lagoon was estimated by hydrological and meteorological observations. As a result, during non-rainfall periods, the net groundwater output to the surrounding marsh was evaluated, of which the magnitude exhibited the linear relationship with the water level (m asl) of the pond. Hence, it is suggested that the groundwater output to the sea through the gravelly confined aquifer prevails rather than the unconfined groundwater output to the marsh.

Keywords: coastal lagoon, sporadic opening, water budget, marsh, Pacific Ocean

Variabilities of isotope ratios and turbidity of the Yangtze River water and their impacts on the East China Sea

IRINO, Tomohisa^{1*}, SAITO, Keita², TADA, Ryuji², LUO Chao³, ZHENG Hongbo³

¹Hokkaido University, ²University of Tokyo, ³Nanjing University

Water discharge and suspension load of a river are potentially recorded in sediments in the drainage and / or the river mouth. Isotope composition of fossil calcareous skeletons and detrital provenance and flux reconstructed from the sediment samples could provide us useful proxies for paleoclimatic study. Sediment load from the Yangtze River to the East China Sea (ECS) from the delta to the Okinawa Trough have been widely used to reconstruct the East Asian summer monsoon (EASM) in the past since the water discharge from the Yangtze would be highly affected by monsoon rain, which could deliver much fresh water and sediment to the ECS. The past impact of fresh water from the Yangtze could be reconstructed from stable oxygen isotope signal recorded in the fossil calcareous skeletons found in the ECS sediments, which has also been used as proxy for EASM.

Theoretically, sediment provenance and its yield could be changed from time to time depending on the distribution of precipitation which would control the balance of water discharges from the tributaries. Change in the precipitation distribution also affects the water isotopic composition of each tributary and then the main stream of the Yangtze. Although such variability could change the end-member composition and concentration of the fresh water and sediment load provided to the ECS, paleoceanographic studies in this region have not considered well about the potential change in the basic condition. Therefore, we need to know the water isotope and sediment budget along the Yangtze main stream with regards to the inputs from its major tributaries in order to understand the potential effects from the change in the distribution of the EASM precipitation.

For this purpose, we have started a systematic sampling of the Yangtze River water to determine the stable oxygen and hydrogen isotope ratios and suspension loads as well as the ECS surface water since summer in 2011. We will report the seasonal variations of isotope and turbidity of the River water in comparison with the distribution of surface water mass in the ECS.

Keywords: Yangtze, Oxygen isotope, Hydrogen isotope, Suspended matter, East China Sea, East Asian monsoon

The mechanism of suspended sediment load from a forested drainage basin

IWASAKA, Wataru^{1*}, CHIKITA, Kazuhisa², Kazuhiro OMORI¹, MAMUN, Abdullah¹

¹Graduate School of Science, Hokkaido Univ., ²Faculty of Science, Hokkaido Univ.

The sediment load of a river is related to the transport, erosion and deposition of nutrient and organic matter. Thus, up to date, many researchers have been exploring the relationship between sediment load and ecosystem. In order to clarify the mechanism of suspended sediment load, this study focused on the hysteresis between the time series of the discharge and suspended sediment concentration in the Oikamanai River, Tokachi, Hokkaido. As a result, the "later type", where a peak suspended sediment concentration precedes a peak discharge, and the "simultaneous type", where both the peaks appear simultaneously, were observed during rainfall runoffs of the Oikamanai River. The later type was observed in 2009 and 2010 with three rainfall runoffs of 8-10 m³/s, while the simultaneous type was seen in 2011 with small rainfall runoffs of less than 5 m³/s except for two typhoon events. The later type is possibly due to the soil erosion on the basin slope by throughflow, while the simultaneous type is probably caused by the erosion of sediment accumulated in the river channel during non-rainfalls.

Keywords: forested drainage basin, suspended sediment load, hysteresis, throughflow, typhoon event

Discharge and sediment load from a subarctic river basin: the Tanana River, Alaska

WADA, Tomoyuki¹, CHIKITA, Kazuhisa^{1*}, KIM, Yongwon², KUDO, Isao³

¹Faculty of Science, Hokkaido University, ²International Arctic Research Center, University of Alaska Fairbanks, ³Faculty of Fisheries Sciences, Hokkaido University

The time series of discharge and sediment load from the subarctic Tanana River basin, Alaska, were simulated by a tank model. As a result, the simulations are reasonable, and hence, the contributions of glacier-melt runoff and glacier-melt sediment runoff to the Tanana discharge and sediment load were quantified at 26 - 57 % and 76 - 94 %, respectively.

Keywords: Glacier-melt runoff, sediment load, POC, PON, Permafrost, Snowmelt runoff

Contribution of detrital materials from the Yukon River to the continental shelf sediments of the Bering Sea

NAGASHIMA, Kana^{1*}, ASAHARA, YOSHIHIRO², TOYODA, Shin³, WADA, Tomoyuki⁴, CHIKITA, Kazuhisa⁴, HARADA, Naomi¹

¹JAMSTEC, RIGC, ²Graduate School of Environmental Studies, Nagoya University, ³Faculty of Science, Okayama University of Science, ⁴Graduate School of Science, Hokkaido University

Bering Sea sediments contain detrital materials from the Yukon River. These materials may contain records of past climate changes in the Arctic area, such as the melting of glaciers around the drainage basin of the Yukon River, which help to understand hydrological cycle in this area. In the Bering Sea, however, the spatial extent and pattern of the detrital materials supplied from the Yukon River is not yet fully understood due to the lack of proxy. For better discrimination of the detrital materials from the Yukon Rivers, we focused on quartz, because it is the major component of both the silt- and sand-sized populations of detrital materials from the Yukon River (Eberl, 2004) and because it is resistant to chemical alteration and physical ablation by weathering, transport, and diagenesis. We adopted two parameters of quartz, the electron spin resonance signal intensity of its E1' center (Toyoda and Hattori, 2000) and its crystallinity index (Murata and Norman, 1976), and characterized the quartz in different sizes derived from the Yukon River. We then estimated the spatial pattern of quartz contributed by the Yukon River on the Bering Sea shelf by applying the method to core-top samples from the continental shelf and slope of the eastern Bering Sea.

The results showed a large contribution of sand-sized quartz from the Yukon River to wide areas of the continental shelf and slope, whereas contributions of clay- to silt-sized quartz from the Yukon River were small, except on the northeastern shelf. These spatial distribution patterns suggest that sand-sized quartz was repeatedly reworked and transported by processes such as storm surges to the outer continental shelf, whereas the clay- to silt-sized quartz on the northeastern shelf was supplied, as suspended materials, directly from the Yukon River. Therefore, the clay- to silt-sized quartz on the northeastern continental shelf probably records past climate changes related to the amount and intensity of the Yukon River discharge, whereas the sand-sized quartz in the eastern Bering Sea probably records variations of stormy weather.

Keywords: Yukon River, Bering Sea, Quartz, Electron Spin Resonance, Crystallinity

Treatment Facilities for Hygienic Fish Market Wastewater

Yustiani Yonik Meilawati^{1*}, Anni Rochaeni¹

¹Dept. of Environmental Engineering Pasundan University

Hygienic fish market in Indonesia is now being developed to have better products and healthy condition in the fishermen and vendors community. Inefficiency and inappropriate of wastewater treatment facilities can cause the water quality of the coastal nearby becomes deteriorated. In this research, we investigate the fish market activities condition in related to the wastewater generated. Samples were taken from wastewater generated by washing activities in Cituis Fish Market; thus the dominant parameters need to be treated can be determined. Data of wastewater quality was collected also from previous researches. The results show that BOD, COD, TSS, E.Coli, oil, and grease concentration have to be reduced to meet the government standard. Determination on treatment unit was conducted by considering the availability of land, user-friendly technology, treatment capacity, low-cost maintenance, and side-product potency. The recommended wastewater treatment unit is called up-flow modified septic tank. Trickling filter is installed to modify the conventional septic tank in order to increase the biological processes inside the tank. Sedimented waste in septic tank chamber can be used for other purposes such as manure and pellet.

Keywords: wastewater treatment facility, fish market wastewater, modified septic tank

Estimating phosphate supply via submarine groundwater discharge in Osaka bay, Japan.

ONODERA, Shin-ichi^{1*}, SHIMIZU, Yuta², SAITO, Mitsuyo³, Masato Fukuoka²

¹Graduate school of Integrated Arts and Sciences, University of Hiroshima, ²Hiroshima University, ³Center for marine environmental studies, Ehime University

Nutrient condition in water environment controls the ecosystem. Groundwater discharge to the oceans is significant as nutrient supply (Slomp et al, 2004 etc). Especially, phosphorus and silica concentration generally are relatively high in coastal area. However, it has not been enough to confirm the source of phosphorus in coastal groundwater in previous studies. This study aims to confirm hydrogeological properties in coastal groundwaters, and estimate the possible phosphorus sources of groundwater.

The study areas are Osaka, Marugame, Okayama, and Fukuyama alluvial plains and small island groundwaters in Hiroshima prefecture. We arranged hydrogeological and groundwater quality data sets in previous studies of Hiroshima University. The phosphorus concentrations were high in anoxic condition. In addition, shallow aquifers around alluvial clay had high concentrations. The phosphorus contents in alluvial sediments of Okayama plain were relatively high around alluvial clay. These results suggest the contribution of phosphorus from alluvial sediment to groundwater.

Keywords: submarine groundwater discharge, phosphate, semi-enclosed bay