## Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



AAS22-12

会場:301B

時間:5月20日12:00-12:15

## JEM/SMILES による O3, ClO, HOCl, BrO の観測について Observation of O3, ClO, HOCl, HO2, and BrO by JEM/SMILES

鈴木 睦  $^{1*}$ , 眞子 直弘  $^2$ , 今井 弘二  $^1$ , 佐野 琢己  $^1$ , 光田 千紘  $^3$ , 内藤 陽子  $^4$ , 秋吉 英治  $^5$ , 尾関 博之  $^7$ , 西本 絵梨子  $^6$ , 塩谷 雅 人  $^6$ 

Makoto Suzuki<sup>1\*</sup>, Naohiro Manago<sup>2</sup>, Koji Imai<sup>1</sup>, Takuki Sano<sup>1</sup>, Chihiro Mitsuda<sup>3</sup>, Yoko Naito<sup>4</sup>, Hideharu Akiyoshi<sup>5</sup>, Hiroyuki Ozeki<sup>7</sup>, Eriko Nishimoto<sup>6</sup>, Masato Shiotani<sup>6</sup>

 $^1$  宇宙研,  $^2$  千葉大/CeRES,  $^3$  富士通 FIP,  $^4$  京大,  $^5$  環境研,  $^6$  京大生存圏研,  $^7$  東邦大  $^1$ ISAS,  $^2$ Chiba U./CeRES,  $^3$ FIP,  $^4$ Kyoto U.,  $^5$ NIES,  $^6$ Kyoto U./RISH,  $^7$ Toho U.

SMILES; Superconducting Submillimeter-Wave Limb Emission Sounder is a 4 K cooled 625-650 GHz limb sounder to observe O3, HCl, ClO, HO2, HOCl, BrO, HNO3, and O3 isotopes. SMILES had been operated on the International Space Station from Oct. 12, 2009 to Apr. 21, 2010. ?Since ISS is 51 degree inclined orbit, 30-45 days SMILES zonal mean could provide diurnal variation of chemical species in the stratosphere and mesosphere. Diurnal variation of O3, ClO, HOCl, HO2, and BrO are compared with two nudged CGCM calculations (SD-WACCM; Specified-dynamics WACCM, and MIROC) and satellite observations.

Diurnal variation of O3 agreed with SD-WACCM over 50-82 km, but small peak in the morning (7 am local time) is apparent for the SMILES but not for the SD-WACCM at 70 km.

Diurnal variation of ClO agreed quite well between SMILES L2 ver. 2.2 and SD-WACCM from 19 to 76 km altitude region. But nighttime ClO value of SMILES L2 ver. 2,2 above 50 km is less than SD-WACCM (70%), which is not clearly explained by the SMILES retrieval issue or our current knowledge of chemical kinetics.

Diurnal variation of HOCl also agreed quite nicely from 31 to 76 km. The nighttime build up of HOCl observed SMILES at 44-68 km are nicely reproduced by the SD-WACCM calculation using JPL2006 chemical kinetics dataset.

HO2 diurnal variation also agreed with SD-WACCM from 24 to 72 km. Above 76 km, SMILES L2 ver. 2.2 needs modification of a priori and its co-variance, and we will get better agreement with model calculations. SMILES L2 ver. 2.2 also shows night time bias due to AOS (Acousto-Optics Spectrometer) characteristics.

キーワード: オゾン層, 塩素系化学種, 臭素系化学種, サブミリ波, 国際宇宙ステーション, 中間圏 Keywords: Ozone layer, Chlorine species, Bromine species, sub-mm, International Space Station, mesosphere