

数理モデルによる全球凍結直後の酸素濃度上昇過程の復元：マンガン及び鉄鉱床形成への示唆

Modeling the rise of oxygen after the Snowball earth: implications for the Paleoproterozoic manganese and iron formation

原田 真理子^{1*}, 田近 英一², 関根 康人², 尾崎 和海³

Mariko Harada^{1*}, Eiichi Tajika², Yasuhito Sekine², Kazumi Ozaki³

¹ 東京大学大学院地球惑星科学専攻, ² 東京大学大学院複雑理工学専攻, ³ 東京大学大気海洋研究所

¹Dept. of Earth and Planetary Science, The University of Tokyo, ²Dept. of Complexity Science & Engineering, The University of Tokyo, ³Atmosphere and Ocean Research Institute, The University of Tokyo

Earth's atmosphere and oceans are considered to have experienced stepwise and irreversible oxidation through its history (e.g. [1]), which may have paved the way to the complicated life such as eukaryotes and metazoans [2]. It is interesting to note that geological evidence suggests that a remarkable increase of oxygen concentration has occurred shortly after the Paleoproterozoic Snowball Earth event, based on widespread depositions of manganese and iron oxides immediately above the glacial diamictites found in the Transvaal Supergroup, South Africa, and the Huronian Supergroup, Canada [3,4]. Carbonate precipitation occurs above the deposition of manganese and iron in both the Transvaal and Huronian Supergroups [3,5], which may represent a climate recovery from the greenhouse condition in the Snowball earth aftermath [3]. In this study we numerically investigated the linkage between a global-scale glaciation and an oxygenation of the atmosphere-ocean system, in the aim of comparing our results to the geological records.

The results of our numerical experiments with a biogeochemical cycle model indicate that the super greenhouse conditions ($p\text{CO}_2 \sim 0.7 \text{ atm}$ and $T \sim 320 \text{ K}$) in the aftermath of the Paleoproterozoic Snowball Earth event significantly enhance the chemical weathering of continents, causing ~ 10 times as high as the present levels of nutrient input and the biological productivity. In the consequence of high levels of biological productivity together with a positive feedback in the atmosphere among a rise in oxygen, ozone formation, and UV shielding of methane, the atmospheric oxygen levels rapidly rise from $< 10^{-5} \text{ PAL}$ to 0.01 PAL (PAL: the present atmospheric level) after the glaciation. The oxygen levels then overshoot to $\sim 1 \text{ PAL}$ in $\sim 10^6$ years after the glaciation due to the high levels of biological productivity sustained by a long-term global warming. Atmospheric oxygen then gradually decreases by oxidizing reducing materials from Earth's interior. Eventually, a steady state of atmospheric oxygen of $\sim 0.01 \text{ PAL}$ is achieved in 10^8 years. Such an irreversible rise in atmospheric oxygen (i.e., from $< 10^{-5} \text{ PAL}$ to $\sim 0.01 \text{ PAL}$) is explained by a transition between different steady state of atmospheric oxygen levels [6] caused by a Snowball Earth glaciation and the subsequent perturbations of biogeochemical cycles.

We found that the rapid oxygenation causes the deposition of manganese and iron in the shallow marine environments. Manganese and ferrous iron in the anoxic deep water are driven by thermohaline circulation, immediately oxidized in the shallow marine water within 10^4 years after the glaciation. Assuming deep water is initially saturated with respect to Mn-carbonates, we derive 10^{15} mol of manganese deposition, which would be sufficient to form manganese ore in the Hotazel formation, Transvaal Supergroup ($\sim 8 \text{ Gt}$, $\sim 10^{14} \text{ mol}$ of manganese [7]). We also found that calcite precipitation is prevented in the ocean during 10^5 years after the glaciation owing to high atmospheric $p\text{CO}_2$. Our results imply that manganese and iron oxides deposition might precede the carbonate deposition, which is consistent with the geological records found both in the Transvaal and Huronian supergroups [3,4].

[1] Kasting (1993) *Science*, 259, 920-926. [2] Runnegar (1991) *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 97, 97-111 [3] Kirschvink et al. (2000) *Proc. Natl. Acad. Sci. USA*, 97, 1400-1405. [4] Sekine et al. (2011) *Earth Planet. Sci. Lett.*, 307, 201-210. [5] Bekker et al. (2006) *Precambrian Research*, 148, 145-180. [6] Goldblatt et al. (2006) *Nature*, 443, 683-686. [7] Tsicos (2003) *Economic Geology*, 98, 1449-1462