## Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



SMP44-07

会場:102A

時間:5月20日11:00-11:15

Ab initio computation on the Fe L-edge X-ray emission spectroscopy of Fe-bearing Mg-SiO3

Ab initio computation on the Fe L-edge X-ray emission spectroscopy of Fe-bearing Mg-

Ab initio computation on the Fe L-edge X-ray emission spectroscopy of Fe-bearing Mg-SiO3

Xianlong Wang<sup>1\*</sup>, Taku Tsuchiya<sup>1</sup> Xianlong Wang<sup>1\*</sup>, Taku Tsuchiya<sup>1</sup>

Behaviors of iron (Fe) in the minerals of the Earth's lower mantle (LM), including valence state, spin state, and chemical environments, at high pressures are important for more detailed understanding the LM properties. The pressure induced spin transition of Fe-bearing MgO and MgSiO<sub>3</sub> perovskite (Pv) were detected of usually by using high-resolution K-edge X-ray emission spectroscopy (XES) [1,2,3] and confirmed by theoretical simulations [4,5]. Since the Fe K-edge XES is associated to the 3p orbital, which is far from the valence orbitals (3d and 4s), it provides no information about Fe's coordination environments. However, Fe L-edge XES can directly determine the distribution and intensity of Fe-3d character. To identify the spin state, valence state and substitution site of Fe in Fe-bearing Pv at the LM pressure range, we systematically investigated the L-edge XES of Fe<sup>2+</sup>- and Fe<sup>3+</sup> (Al<sup>3+</sup>)-bearing Pv under high pressure by using the first-principles method combined with the slater-transition method. Our results show that the spin transition of Fe<sup>2+</sup> and Fe<sup>3+</sup> can be identified easily by the L-edge XES technique. The valence state of Fe can be furthermore certified, since the shift of the first main peak of Fe<sup>3+</sup> is about two times larger than that of Fe<sup>2+</sup> across the spin transition. The width of the L-edge XES of Fe<sup>3+</sup> is also sensitive to the substitution site, indicating that their coordination environments might also be distinguishable from the Fe L-edge XES spectra. These strong sensitivities to the Fe states suggest that the high-resolution Fe L-edge XES measurement would be a useful experimental technique to investigate Fe-bearing silicate minerals. Corresponding experiments are expected.

 $\pm - 7 - F$ : First-principles calculation, First-principles calculation, Fe L-edge XES, Mg perovskite Keywords: First-principles calculation, First-principles calculation, Fe L-edge XES, Mg perovskite

<sup>&</sup>lt;sup>1</sup>Geodynamics Research Center, Ehime University

<sup>&</sup>lt;sup>1</sup>Geodynamics Research Center, Ehime University