

DThe estimation for temporal and spatial fluctuations of forest fire hazard index

Koji Tamai^{1*}

¹Forestry & Forest Products Research Institute

Mapped estimates of the risk of forest fire would benefit forest management, and could be used to decide restrictions on the public use of forest areas. In this study, the litter moisture content ratios were predicted with this model and forest fire hazard were estimated the forested parks in Japan. Model was adapted to the around 40ha area in this park classified into 9 stands with tree height, tree species and slope direction. Fuel moisture decreased with each speed for each forest stand among simulated days depending the solar radiation on the each forest floor. Litter moisture was less than 0.2g g⁻¹ and fire risk is judged to be highest in 7 forest stands among 9 stands on the day after long drought period. On the otherhand, spatial variation of litter moisture was widest to be 0.198 - 0.811g g⁻¹ on the day during the drying process. This means that litter drying speed and fire risk is different between forest stands. Thus, it is significant to construct the forest fire warning system for each forest stand to manage the people activities in this public forested area to prevent the forest fire.

Keywords: Fuel moisture, Solar radiation on the forest floor

Developing Distributed Hydrologic-Vegetation Dynamics Coupling Model: A Case Study in African River Basins

Yohei Sawada^{1*}, Toshio Koike¹, Cho Thanda Nyunt¹, Akio Yamamoto², Masaru Kitsuregawa²

¹Department of Civil Engineering, The University of Tokyo, ²Institute of Industrial Science, The University of Tokyo

We develop an eco-hydrological model (WEB-DHM + DVM) that can simulate vegetation growth, soil moisture, river discharge, ground water dynamics and land surface fluxes at the same time in river basin scales. Modeling of vegetation dynamics in a hydrological model makes it possible to discuss the integrated management of water resources and bioresources in river basins. Moreover, dynamic vegetation modeling can contribute climate and weather prediction because we have a strong feedback of vegetation distribution to regional climate through land-atmosphere interactions especially in semi-arid regions. Considering landscape changes driven by human exploitation and climate change, we need to calculate vegetation dynamics as a diagnostic valuable in a basin-scale hydrological model. We apply the model to the Medjerda River Basin located in North Africa for a model confirmation study. The simulated river discharge has a good agreement with the in-situ observed river discharge. In addition, we show the model has adequate capacity for simulating vegetation dynamics in the semi-arid region by comparing simulated leaf area index (LAI) with the Moderate Resolution Imaging Spectroradiometer (MODIS) 8-daily LAI product. We also demonstrate the assessment of climate change impact on vegetation dynamics in the Volta River Basin located in West Africa by using WEB-DHM + DVM and multi General Circulation Models (GCMs) output. Our future projections and sensitivity analyses show that an extension of dry season duration and high land surface temperature produced by climate change may cause a dieback of vegetation in West Africa, while an increase of atmospheric humidity has a positive impact on vegetation growth. The negative impacts of certain climate forcings sometimes overwhelm the positive impacts of the other forcings, and positive and negative impacts sometimes cancel each other. Thus, there are different magnitudes of change in biomass amount in different GCMs, although we select three GCMs whose climatologies agree well with past climate. This approach demonstrates that multi-model climate change assessment is crucial, and the sensitivity analysis developed here is useful for extracting principal environmental drivers of terrestrial biomass under a changing climate. The method in this study makes it possible to address the impact of future change of terrestrial biomass on climate and water resources on a regional scale.

Keywords: vegetation dynamics, distributed hydrological model, climate change

Separation of ET in Nile delta into transpiration and evaporation (E) by canopy model and E reduction by windbreak trees

Tatsuki Shimizu^{1*}, Michiaki Sugita²

¹Environmental Science, Life & Environ. Sci., University of Tsukuba, ²Geoscience, Life & Environ. Sci., University of Tsukuba

Agriculture can be considered as human intervention to the regional ecosystems. Thus its influence to hydrologic cycle is one of the important contemporary issues. In the present study, evapotranspiration (ET) of cropland in Nile delta was determined by applying eddy correlation method. In addition, 2-layer canopy model was applied to separate ET into soil evaporation (E) and transpiration (T). It was found that in the furrow period in 2011, ET (=E) was 221 mm for 2 months, while in growing season of maize ET was some 554 mm with T=256 mm and E=298 mm. With introduction of

windbreak trees to the edge of cropland in the direction of prevailing wind direction, both E and T were reduced some 20-25% during growing season, and E was reduced by 30% during furrow period. Transpiration of wind break trees were found to be a minor component.

Keywords: Nile-delta, croplands, evapotranspiration, soil evaporation and transpiration, Windbreak trees

CO₂ flux over the Lake Kasumigaura and factors influencing its variation: a preliminary analysis of 5-year observation

Takanari Takahashi¹, Michiaki Sugita^{2*}

¹Geoscience, Life & Environ. Sci., University of Tsukuba, ²Life & Environ. Sci., University of Tsukuba

In this research, CO₂ flux over the Lake Kasumigaura was measured mainly with eddy correlation method at the center observatory from June in 2007 to November in 2012 for five years. It was found that CO₂ flux of hourly time scale changed in the range of about -0.5~0.5 mg/m²/s. Absorption and release of CO₂ occurred alternatively, and both processes occurred both in the daytime and the night. Comparison with previous studies indicated that the CO₂ flux over a unit area over the Lake Kasumigaura was slightly smaller than that of a forest, and larger than that of a prairie, and several orders of magnitude larger than that of the ocean. With regards CO₂ flux changes on seasonally time scale, it was found that the amount of daily CO₂ release differed greatly in the range of about -10~10 g/m²/day. The amount of the 10-day average CO₂ release changed in the range of about -6~2 g/m²/day regardless of the season through the whole period. However, negative flux was dominant in most of the time. A change of the dissolved CO₂ levels C_w of the lake was found to be a factor of CO₂ flux change on seasonally time scale. There are an internal factors and an external factors that affect C_w values. C_w change cannot be attributed to a single factor, but various factors work simultaneously and intricately.

Keywords: eddy correlation method, Lake Kasumigaura, CO₂ flux, photosynthesis, phytoplankton

Changes in watershed environments and water balance of Lake Kasumigaura from 1970s to present

Kazuya YAMAMOTO^{1*}, SUGITA, Michiaki²

¹Graduate School of Life & Environ Sci., University of Tsukuba, ²Faculty of Life & Environ Sci., University of Tsukuba

Water balance of Lake Kasumigaura was estimated by Muraoka (1981) for 1970s. Since then, watershed environments have been changed drastically; also human interventions to the water balance have been made in various forms. However, systematic and comprehensive estimation of water balance has not been carried out. Thus in the present study, first the same methods of Muraoka (1981) were applied to the present measurements to estimate water balance. Next, some of the water balance components which have become available with advent of better measurements technology were employed. Among them is the lake surface evaporation.

As a result, annual inflow and outflow of Lake Kasumigaura increased by about 30%. Main cause of this increase was the increase of artificial flow (both intake and outflow) such as service water, industrial water, and treated sewage. Natural hydrologic flow such as river and groundwater discharge also increased but by smaller amount of 5 - 35%.

Keywords: water balance, Lake Kasumigaura, water intake, hydrological cycle

Year-round observation of energy balance components within a permafrost black spruce forest in interior Alaska

Taro Nakai^{1*}, KIM, Yongwon¹, BUSEY, Robert C.¹, Rikie Suzuki², NAGAI, Shin², KOBAYASHI, Hideki², PARK, Hotaek², SUGIURA, Konosuke², ITO, Akihiko³

¹IARC, UAF, ²JAMSTEC, ³NIES

Black spruce is one of the most abundant forest types in the North American boreal forest, also occupying approximately 44 % of the forest cover in interior Alaska. In this study, energy balance components such as radiation and eddy covariance fluxes were observed year-round in 2011, within a black spruce forest underlain by permafrost at the University of Alaska Fairbanks Poker Flat Research Range, located in interior Alaska. This research was conducted under the JAMSTEC-IARC Collaboration Study, with funding provided by the Japan Agency for Marine-Earth Science and Technology (JAMSTEC) under a grant to the International Arctic Research Center (IARC).

Here, if ground heat flux was observed using the default value of thermal conductivity, the sum of sensible and latent heat flux exceeded the available energy in summer. By multiplying the correction factor of 0.74 by ground heat flux, the deficit of energy balance was minimal. This 0.74 value was consistent with the thermal conductivity of the heat flux plate (HFPO1SC) value of $0.8 \text{ W m}^{-1} \text{ K}^{-1}$, whereas the maximum Sphagnum thermal conductivity has been reported to have ranged from 0.5 to $0.6 \text{ W m}^{-1} \text{ K}^{-1}$ at high volumetric water content. As a result, the energy balance of this black spruce forest was almost closed in summer, and ground heat flux was proven to play an important role in energy balance, explaining 26.5 % of the net radiation in summer.

On the other hand, a large energy balance deficit was observed in spring. The energy balance deficit during the snowmelt season was mostly explained by the latent heat energy of fusion consumed by snowmelt, which was calculated from the observed snow water equivalent.

The mean daily evapotranspiration of this forest in summer was 1.37 mm day^{-1} , considered typical for boreal forests. Further, the annual evapotranspiration and sublimation amounted to $207.3 \text{ mm year}^{-1}$, which was much smaller than the annual precipitation, and sublimation accounted for 8.8 % ($18.2 \text{ mm year}^{-1}$) of this annual total value. Thus, sublimation is *not* negligible in the annual water balance for boreal forests.

Keywords: energy balance, black spruce, permafrost, ground heat flux, evapotranspiration, sublimation

Climate change and the ecohydrological feedback of Siberian permafrost ecosystems

Tetsuya Hiyama^{1*}, Shamil Maksyutov², Heonsook Kim², Takahiro Sasai³, Yasushi Yamaguchi³, Atsuko Sugimoto⁴, Hitoshi Yonenobu⁵, Takeshi Ohta⁶, Ayumi Kotani⁶, Kazukiyo Yamamoto⁶, Takeshi Yamazaki⁷, Kazuhiro Oshima¹, Hotaek Park⁸

¹Research Institute for Humanity and Nature, ²National Institute for Environmental Studies, ³Graduate School of Environmental Studies, Nagoya University, ⁴Graduate School of Environmental Science, Hokkaido University, ⁵Department of Health and Living Sciences Education, Naruto University of Education, ⁶Graduate School of Bioagricultural Sciences, Nagoya University, ⁷Graduate School of Science, Tohoku University, ⁸Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology

High levels of precipitation in the Lena River Basin, Siberia, from 2005 to 2008 led to tremendous changes in terrestrial environments. The changes observed include a deepening and moistening of the active layers, hindrance of tree growth, and the expansion of water surfaces due to floods. The anomalously wet condition of forest soils caused larch trees to wither at our forest monitoring site in the middle part of the basin. However an analysis of satellite data revealed that such tree withering occurred only at certain points. On the basis of our permafrost-ecosystem models, we have identified increases in thawing depth and surface soil moisture, and an increase in net primary production. The annual maximum thawing depth (AMTD) was revealed to have gradually increased (deepened) on a decadal scale. Increase in terrestrial water storage in the Lena River Basin generated increases in river base flows during the open water season. Our results also indicated that between 1950 and 2008 the basin-scale AMTD increased at an average rate of approximately 1 cm/year in the region. Moistening and warming of surface soil affect methane emissions from Siberian terrestrial ecosystems. Regional methane fluxes were estimated using an inversion model with data collected from aircraft and tower measurements in Siberia. In 2007 and 2008, enhanced methane fluxes from the wetlands in Western Siberia were estimated under relatively wet conditions with high temperatures. Interestingly, methane fluxes after 2008 have gradually decreased but those in Eastern Siberia have increased unsymmetrically. Such an unsymmetrical (seesaw) pattern between Western and Eastern Siberia has also been observed for carbon dioxide exchanges in terrestrial ecosystems. Gross primary production and ecosystem respiration in the 2000s were estimated using our permafrost-ecosystem models, which showed a decreasing trend in Western Siberia and an increasing trend in Eastern Siberia. These differences were primarily due to the differences in the trends of temperature and precipitation between the two regions.

Keywords: global warming, permafrost, annual maximum thawing depth (AMTD), methane, carbon dioxide

Methane flux observation over a forest ecosystem by micrometeorological and chamber methods using laser based analyzers

Masahito Ueyama^{1*}, Ryota TAKEUCHI¹, Yuriko TAKAI¹, Yoshiyuki Takahashi², Mioko ATAKA³, Kenshi Takahashi⁴, Yoshiko KOSUGI³, Ken HAMOTANI¹

¹Graduate School of Life and Environmental Sciences, Osaka Prefecture University, ²Center for Global Environmental Research, National Institute for Environmental Studies, ³Graduate School of Agriculture, Kyoto University, ⁴Research Institute for Sustainable Humanosphere, Kyoto University

Recent applicability of laser-based analyzers allows the FLUXNET community continuously measure methane flux over various terrestrial ecosystems with high precisions. Such measurements have mainly conducted at ecosystems, such as wetland and agricultural land, which are expected to be an obvious methane source. On the other hand, measurements at forest ecosystems are limited in part due to difficulty in measuring approximately 1-order smaller magnitude of fluxes than those of emission fluxes. In this study, we have conducted a continuous measurement of methane flux over a forest in order to quantify the spatially representative flux, understand processes, and improve our ability to measure the fluxes.

The measurement was conducted at Fujihokuroku Flux Research site in Yamanashi, Japan. A 50-year-old stand of planted Japanese larches dominates on volcanic ash soils, homogeneously. The hyperbolic relaxed eddy accumulation (HREA) method was applied to measure canopy scale fluxes since August, 2011. Five-height vertical methane concentration profile was also measured by same system switched flow lines. The robustness of the HREA method was confirmed by compared CO₂ fluxes with the eddy covariance and HREA method. The measured fluxes were corrected for storage terms. The automated dynamic closed chambers were installed at six plots in the forest soil since October 2012. Concentration of methane, CO₂, and water vapor were measured by flowing sample air from the HREA and chamber systems to respective laser-based analyzers (GGA-24r-EP and FGGA-24r-EP, Los Gatos Research Inc., USA) after dehydration. The analyzers were conducted two-point calibrations every-day, where there was no detectable change in the sensitivity.

Throughout the measurement period from August 2011 to December 2012, methane uptake was measured by the HREA method. The methane uptake showed a clear seasonal variation roughly coinciding with seasonality in soil temperature near the ground, where sensitivity to soil temperature (Q₁₀) was 1.8. The methane uptake decreased to near zero at winter when the forest floor was covered by snow (approximately 30 cm depth). As previously reported for CO₂ fluxes, the annual methane sink strongly changed with the friction velocity (u^*) filtering correction. The annual sink was 900 mg CH₄ m⁻² y⁻¹ with the u^* -filtering correction and 700 mg CH₄ m⁻² y⁻¹ without the u^* -filtering correction. Based on the profile system, methane concentration was most depleted near the soil surface, indicating that measured methane sink mostly associated with soil processes. This was also consistent with the chamber measurement, where measured fluxes by the HREA and chamber systems were similar range.

Keywords: Methane flux, HREA method, automated dynamic chamber method, forest, continuous observation, laser-based analyzer

Carbon dioxide balance of tropical peat ecosystems

Takashi Hirano^{1*}, SEGAH, Hendrik¹, KUSIN, Kitso², LIMIN, Suwido², TAKAHASHI, Hidenori¹, OSAKI, Mitsuru¹

¹Hokkaido University, ²University of Palangkaraya

Tropical peatlands, which are widely distributed on lowlands in Southeast Asia, have accumulated a huge amount of soil carbon under peat swamp forests over millennia. However, the carbon pool is presently disturbed on a large scale by land development and management, and consequently has become vulnerable. Peat degradation occurs most rapidly and massively in Indonesia, because of fires, drainage and deforestation of swamp forests. Peat fires release carbon dioxide (CO₂) through peat combustion intensively but occasionally, whereas drainage increases CO₂ emission steadily through the acceleration of oxidative peat decomposition. Therefore, tropical peatlands present the threat of switching from a carbon sink to a carbon source to the atmosphere. However, the ecosystem-scale CO₂ exchange is still unknown in tropical peatlands. A long-term field experiment in Central Kalimantan, Indonesia showed that tropical peat ecosystems, including a relatively intact peat swamp forest with little drainage (UF), a drained swamp forest (DF) and a drained burnt area (DB, a former swamp forest), functioned as net carbon sources. Mean annual net ecosystem CO₂ exchange (NEE) (with a standard deviation) for four years from July 2004 to July 2008 was 174 with 203, 328 with 204 and 499 with 72 gC m⁻² y⁻¹, respectively, for the UF, DF and DB sites. The carbon emissions increased according to disturbance degrees. We found that the carbon balance of each ecosystem was chiefly controlled by groundwater level (GWL). The NEE showed a linear relationship with GWL on an annual basis. The relationships suggest that GWL lowering of every 0.1 m causes additional net annual CO₂ emissions of 79-238 gC m⁻² probably because of the enhancement of oxidative peat decomposition. In addition, CO₂ uptake by vegetation photosynthesis was reduced by shading due to dense smoke from peat fires ignited accidentally or for agricultural practices. Our results may indicate that tropical peatland ecosystems are no longer a carbon sink under the pressure of human activities.

Keywords: drainage, draught, eddy covariance technique, ENSO, fire

Nutrient limitation and the responses of the carbon dynamics in forest ecosystems to global change

Kanehiro Kitayama^{1*}

¹Graduate School of Agriculture, Kyoto University

It is well known that net primary productivity and/or decomposition rate can be limited by nutrients when climate is mild. Particularly because phosphorus and nitrogen are directly involved in the carboxylation biochemical processes, the rate of carbon fixation in forest ecosystems can become lowered when the supply of these nutrients is low. The magnitude of the responses of forest ecosystems to global warming can be, thus, dependent on the strength of nutrient limitation. It is expected that the temperature dependency of the net primary productivity of a forest is low if the forest is strongly limited by nutrients. However, how/if such interactions of nutrients and temperature occur in natural forest ecosystems has not been elucidated well. In order to investigate such a research question, I and co-workers are comparing two altitudinal gradients (one phosphorus-impooverished and the other relatively phosphorus richer) on a humid tropical mountain. Our long-term observation indicates that trees adjust phosphorus-use efficiency when the demand for phosphorus co-varies with temperature and maintain net primary productivity despite the strong phosphorus limitation. Our results indicate that the temperature dependency of the net primary productivity under a strong nutrient limitation can be greater than expected earlier.

Keywords: Nutrients, P-use efficiency, Net primary productivity, Tropical rain forest, Altitude gradient, Carbon dynamics

^{13}C pulse labeling technique for study carbon allocation in tree

Masako Dannoura^{1*}

¹Graduate School of Global Environmental Studies

Carbon is fixed by photosynthesis of plant and used for growth or storage, then finally released to atmosphere by respiration. It is necessary for better understanding of future climate change to construct process based model and estimate which and how factors affect carbon allocation in forest ecosystems. As one of the methods to analyze carbon balance, pulse labeling technique will be introduced. It is a method to trace stable carbon isotope which is fixed by the tree to see when and where carbon goes. By this method, the velocities of carbon of different tree species and its seasonal dependency were observed. Some experiments conducted in both Japan and abroad will be shown.

Keywords: Stable carbon isotope, pulse labeling, carbon allocation, ^{13}C

Japan Long-Term Ecological Research Network (JaLTER)

Hiroyuki Muraoka^{1*}

¹Gifu University, JaLTER

JaLTER was established in November 2006 to provide scientific knowledge which contributes to conservation, advancement and sustainability of environment, ecosystem services, productivity and biodiversity for a society by conducting long-term and interdisciplinary research in ecological science including human dimensions. JaLTER is a formal member network of the International Long-Term Ecological Research Network (ILTER Network). JaLTER contains 56 sites including forests, grasslands, cropland, freshwater and marine. The goal of JaLTER are (1) Creation of general knowledge based on multidisciplinary long-term and large-scale research, (2) Creation of well-designed database to exchange and share original data to support scientific communities, general public people and policy makers, and to find better solutions for critical ecological and environmental problems, (3) Promotion of education regarding long-term and large-scale changes of ecosystem and environment. (4) Facilitation of collaboration and coordination among scientists of long-term ecological researches. To achieve these goals, we are promoting research theme such as (a) Response and feedback of biodiversity and ecosystem functioning under the climate changes, (b) Hydro-biogeochemical processes and ecosystem interaction from terrestrial to marine ecosystem, (c) Development and establishment of ecosystem monitoring network and techniques with multiple scales and dimensions.

In addition to the site-network based research on ecological theme, JaLTER has been promoting to link ecosystem research networks and institutes, so called "J-community", by JaLTER, JapanFlux (CO₂ flux network), JAXA (Japan Aero-Space Agency) and JAMSTEC (Japan Agency for Marine-Earth Science and Technology). The J-community was established to meet multidisciplinary research on ecosystem structure, function and biodiversity under climate change. As part of this activity, a joint research project to develop earth observation algorithm by a new earth observation satellite "GCOM-C" has been conducted since 2009. In addition, recently, new projects were established to clarify the relationship between biodiversity and ecosystem functions, and to evaluate their vulnerability to environmental changes.

Keywords: LTER, ecological research, ecosystem, biodiversity

Hydrologic Responses to the changes in watersheds environments from Meiji-era to present

Takuma Watanabe^{1*}, Michiaki Sugita²

¹College of Geoscience, School of Life and Environmental Sciences, University of Tsukuba, ²Doctral Program in Graduate School of Life and Environmental Sciences, University of Tsukuba

Two sub-basins were selected from within the Kasumigaura watershed for this study. Currently one basin (Sonobe) is mostly a rural area which was covered forests in Meiji period; thus changes are mainly from forests to agricultural fields. Another basin (Onogawa) is more of a sub-urban area which used to be forest/grass lands. A hydrologic model was applied to each of the basins; the model was calibrated against discharge and flux data in this area. After the calibration, two sets of land use were applied to the model, one representing Meiji period and one current condition, while the meteorological forcing data were kept the same by assuming no change of climate.

By comparing the two sets, decrease in evapotranspiration, and net radiation, and increase in discharge, and rise in water table were found until now in comparison to the case in Meiji at Onogawa. These changes can be summarized as follows: about 30% decrease in evapotranspiration, more than twice increase in discharge, about 15% decrease in net radiation, and about 30% decrease in latent heat flux. From inspection of the distribution maps of these variables in watershed, the above changes were identified to have occurred at the areas where forest changed into urban, or agricultural areas. It was considered that the main reason of the decrease in evapotranspiration was the decrease in canopy storage, which caused increase in discharge. Also, increase in surface runoff occurred due to increase in impermeable area; increase in sensible heat flux was due to glass, and urban area that has high albedo. On the other hand, in the comparison with the Sonobegawa results, it was clearly found that there is a difference of impact on water and heat balance due to difference of increase rate of urban, agricultural area.

Especially, in the Onogawa watershed, the rate of increase of urban area was larger in upstream area. Also agricultural area increase was most common in midstream area, and paddy field area was abundant in downstream areas. These land uses were distributed as a mass in Onogawa, while in Sonobegawa changes occurred in scatter, which impacted on the water and heat balance changes.

Keywords: semi-distributed hydrological model, Land use change, Lake Kasumigaura, catchment water budget

Developing and improving a simple runoff model of dissolved organic carbon considering soil infiltration and river runoff

Kazunori Ebata^{1*}, Kei Nishida²

¹UY, ²ICRE-UY

Estimation of dissolved organic carbon (DOC) runoff from forested watershed is important for carbon cycle researches and water quality control. Empirical regression models or process-based models have been proposed up to now, however, those are not yet ready for general use because of lack of internal processes as well as the complexity of parameterization. I proposed a simple DOC runoff model considering both soil processes, i.e. soil infiltration and seasonal changes, and river runoff processes assuming DOC source area. Two years' data for model development were collected from three different scale-stations in the Mizugaki research watershed. In soil sub model, dry-wet cycle in soil was successfully simulated by advection-diffusion and dissolution formulation and soil water movement was significantly responsible for overestimation especially for dry soil. In runoff sub model, the results from soil sub model being imported, DOC source area was assumed by hydrograph separation and wetland ratio. The model was highly applicable upstream, however, overestimated downstream due to error in quantifying riparian zone as DOC source area. In addition, hydrological model (TOPMODEL) was introduced for the further improvement.

Keywords: DOC, runoff model

Acidification of the soil by air pollution and the relation of withering of trees.

Teiko Omori^{1*}, YOSIIKE Yuzo², OKAMURA Sinobu³, IWASAKI Masato⁴

¹Department of Science, Toho University(former), ²Department of Science, Toho University(former), ³Department of Science, Toho University, ⁴A Comprehensive School affiliated with Ashikaga Institute of Technology

Trees wither by sulfuric acid of air pollutant. Sulfuric acid generates by combustion of fossil fuel. Sulfuric acid moves by wind and adheres to trees. Adhered sulfuric acid is added to the soil of root with rain. The soil dissolves the metal of ingredient by sulfuric acid. Dissolved metal ion is absorbed into water in trees. The absorbed metal ion combines with phosphoric acid in the trees. As the metal phosphoric acid compound which combined has very low solubility, the work as phosphoric acid becomes impossible. Trees become the same phenomenon as the shortage of phosphoric acid and decline. As for the tree which weakened, creation of the defense ingredient to an insect becomes impossible. As for a pine, the generation of resin decreases. Tannin combines with metal and detoxifies a Japanese oak. As a result, trees become the food and residence of an insect. Insects increase so much and trees wither. The main cause of withering is by the acidification of soil. Withering is saved by neutralizing the acidified soil. As for the method of neutralization, it is optimal to use charcoal. Trees include Na, K, Ca, Mg and P. Those remain as carbonate or oxide into charcoal and if rain is added to charcoal, it will become an alkaline solution and it neutralizes acid soil. The remaining elements are contained at the rate which the tree needs. As a result, they become an ideal nutrient. Charcoal becomes a moisture absorbent and the breeding place of soil bacteria. As a result, the soil is activated and can prevent withering of trees.

Keywords: Air pollution, Withering., Charcoal., Metal phosphate., Tannic acid iron, Acidification of soil.

Evaluation of the acid deposition on to Aya forest, the UNESCO biosphere reserve forest in Miyazaki, Japan

Koyomi Nakazawa^{1*}, Osamu Nagafuchi², Yuki Nishida², Seiichi Kanetani³

¹Osaka University, ²The University of Shiga Prefecture, ³Forestry and Forest Products Research Institute

The Aya forest is located at the northern limit of the evergreen broad-leaved forest zone in East Asia. In addition, it has various endemic species in Japan. People who live near Aya forest have been using these natural resources with a sustainable way for more than five decades until now. In July 2012, Aya forest was designated as a UNESCO biosphere reserve, the Man And the Biosphere programme (MAB). In order to preserve this area and its culture, we evaluate the effect of long-range transport air pollution on to the Aya forest ecosystem. The research was conducted in two years, from 2010 to 2012. We installed bulk deposit samplers and Ogawa passive samplers both in-canopy and the outside of forest experimental site. After collecting samples, we measured major ion concentration for wet deposition and NO, NO₂, NH₃, O₃ and SO₂ gaseous concentration for dry deposition, respectively. In addition, to evaluate the actual deposition amount on to Aya forest, we collected conifer needle tree leaves from six trees, which grow up in this area.

During this experimental campaign, we found following things. In 2010, the effect of eruption of Mt. Shin'moe was observed. However, we couldn't find the effect of long range transport of air pollution from East Asian continent which was shown in Yakushima Island experimental site. The result of conifer needle tree leaves provide us the different SO₄²⁻ concentration level between north and south side of leaves which were collected from the same tree. This tendency was observed in all collecting samples.

Keywords: Aya biosphere reserve, Atmospheric acid deposition, Long range transport, Man And the Biosphere, conifer needle tree

Toward the continuous development of hydrological and flux monitorings in terrestrial ecosystems

Makoto Tani^{1*}

¹Graduate School of Agriculture, Kyoto University

Interactions between water/chemical cycles and terrestrial ecosystems are the most important base supporting our irreplaceable planet. A close collaboration of the geoscience for climate and tectonic activities with ecosystem sciences for biological responses is needed for this interdisciplinary subject. This highly requires continuous monitorings in ecosystem study sites, but the organized system has not been established. The strategy is discussed.

Keywords: terrestrial ecosystem, hydrological monitoring, flux, continuous observation, gas exchange between land and the atmosphere

Characteristics of Water Chemistry in Degraded Peatland Groundwater and Riverwater in Indonesia

Masayuki Itoh^{1*}, MUHAMMAD, Ahmad², NEOH, Kok-Boon¹, WATANABE, Kazuo¹, GUNAWAN, Haris², SYAHRONI, Reza², FURNANDO, Edo², KOZAN, Osamu¹

¹Center for Southeast Asian Studies, Kyoto University, ²Faculty of Mathematics & Natural Sciences, Riau University

Tropical peatland forests in Southeast Asia are considered to be one of the most important parts of larger ecosystems due to the huge amount of carbon stock and biodiversity they contain. Yet, recent rapid and intensive deforestation to procure timber and land for commercial plants or crops (Oil palm or rubber plantations) must have induced fundamental changes in the material cycling.

We focused on the effects of human impacts such as deforestation, plantation, and manmade fires on groundwater chemistry in the degraded peatland area in East part of Sumatra Island, Indonesia.

We measured dissolved organic and inorganic matters in peatland groundwater both in wet and dry seasons by using piezometers which are installed in degraded peatland sites. We compared the results of both bare land site after the deforestation and the fires and oil palm site planted after the fire.

Keywords: Tropical Peatland, Groundwater, Biogeochemistry, Dissolved Organic Matter