

グリシン前駆体 CH_3NH_2 の推定存在量 Estimated Abundances of a Precursor of Glycine, CH_3NH_2

大石 雅寿^{1*}

Masatoshi Ohishi^{1*}

¹ 国立天文台

¹National Astronomical Observatory of Japan

It is widely accepted that prebiotic chemical evolution from small to large and complex molecules would have resulted in the Origin of Life. On the other hand there have been conflicting two views regarding where inorganic formation of organic molecules (hereafter OMs) occurred in the early Earth, in the Earth or out of the Earth. Ehrenfreund et al. (2002) indicated that exogenous delivery of OMs to the early Earth could be larger than their terrestrial formation by three orders of magnitude. If amino acids are formed in interstellar clouds, significant amount of them may be delivered by comets and/or asteroids to (extra-solar) planets. Detection of amino acids would accelerate the discussion concerning the universality of “life”.

So far, many trials to detect simplest amino acid, glycine ($\text{CH}_2\text{NH}_2\text{COOH}$), were made towards Sgr B2 and other high-mass forming regions, but none of them were successful due to insufficient sensitivities and spatial resolution of telescopes. This is the background that detection of amino acids and other prebiotic in the Universe is one of the key science targets for ALMA. However we need to have carefully selected good candidate sources for amino acids before conducting searches for amino acids by ALMA since lines could be contaminated by other molecular lines. One idea would be to survey precursors of amino acids; higher abundances sources of precursors would be amino acid rich sources.

Although the chemical evolution of interstellar N-bearing OMs is poorly known, methylamine (CH_3NH_2) is proposed as a precursor to glycine; theoretical and laboratory studies have indicated that glycine is formed on icy grain surface from methylamine and CO_2 through UV irradiation (Holtom et al. ApJ, 626, 940 (2005), Kim and Kaiser et al., ApJ, 729:68 (2011)). These studies also suggest that methylamine can be formed from abundant species, CH_4 and NH_3 , on icy dust surface. Although CO_2 is widely known in molecular clouds, distribution of methylamine is poorly known. Further methylene imine (CH_2NH) would be related to CH_3NH_2 . These species are thought to be formed through hydrogenation (addition of hydrogen) to HCN on dust surface (Dickens et al., 1997, Kim and Kaiser et al. 2011): $\text{HCN} \rightarrow \text{CH}_2\text{NH} \rightarrow \text{CH}_3\text{NH}_2$. This is similar to the hydrogenation of CO to form CH_3OH : $\text{CO} \rightarrow \text{H}_2\text{CO} \rightarrow \text{CH}_3\text{OH}$.

The first detection of methylamine was made by Kaifu et al. (1974). But even now, CH_3NH_2 is known toward two objects (Orion KL and Sgr B2) only. CH_2NH has been reported only in Sgr B2, W51, Orion KL, and G34.3 (Dickens et al. 1997).

Therefore it would be crucial to estimate abundances of CH_3NH_2 and CH_2NH by using a chemical reaction network, which may be compared with observations. In this paper we present preliminary results of estimated abundances of these species, which would provide good guidance in finding good glycine sources and in understanding poorly known chemistry of N-bearing organic molecules in the Universe.

キーワード: 宇宙有機分子, 化学進化, 惑星形成

Keywords: Organic Molecules in Space, Chemical Evolution, Planetary Formation

NGC 6334-V の近赤外円偏光撮像:アストロバイオロジーへの示唆
Near-Infrared Circular Polarization Images of NGC 6334-V: Implication for Astrobiology

権 静美^{1*}, 田村 元秀¹
Jungmi Kwon^{1*}, Motohide Tamura¹

¹ 総合研究大学院大学 / 国立天文台

¹The Graduate University for Advanced Studies, National Astronomical Observatory of Japan

We present results from deep imaging linear and circular polarimetry of the massive star-forming region NGC 6334-V. These observations show high degrees of circular polarization (CP), as much as 22% in the Ks band, in the infrared nebula associated with the outflow. The CP has an asymmetric positive/negative pattern and is very extended ($\sim 80''$ or 0.65 pc). Both the high CP and its extended size are larger than those seen in the Orion CP region. Three-dimensional Monte Carlo light-scattering models are used to show that the high CP may be produced by scattering from the infrared nebula followed by dichroic extinction by an optically thick foreground cloud containing aligned dust grains. Our results show not only the magnetic field orientation of around young stellar objects, but also the structure of circumstellar matter such as outflow regions and their parent molecular cloud along the line of sight. This is the second case to support the large CP in scattering protostellar nebulae as a possible explanation for the extraterrestrial origin of homochirality of life on Earth.

太陽系外惑星バイオマーカー検出に向けた光捕集機構のエネルギー転移計算 The energy transfer calculation of light harvesting systems for detecting biomarker on extrasolar planets

小松 勇^{1*}, 梅村 雅之¹, 庄司 光男¹, 矢花 一浩¹, 白石 賢二¹, 神谷 克政¹, 栢沼 愛¹, 田口 真彦¹, 佐藤 皓允¹
Yu Komatsu^{1*}, Masayuki Umemura¹, Mitsuo Shoji¹, Kazuhiro Yabana¹, Kenji Shiraishi¹, Katsumasa Kamiya¹, Megumi Kayanuma¹,
Masahiko Taguchi¹, Akimasa Sato¹

¹ 筑波大学

¹University of Tsukuba

A large number of extrasolar planet candidates have been detected by Kepler. Moreover the planets in habitable zone have already been detected, and ELP (Earth-like planet) is expected to be discovered. As detecting signs of life from the spectrum of ELPs in future, several indicators called as biomarkers or biosignatures were proposed [1]: (1) molecule absorption line, such as O₂, CH₄, and (2) red edge, which is a strong contrast in NIR (~700-750 nm) that derives from plant's feature. Red edge comes from absorption in visible by the chlorophyll, which is one of the photosynthetic pigments, and reflectance in NIR due to the structural features such as cell wall, air space of leaf and so on. The spectrum considered as red edge is observed by remote sensing or earthshine [2].

However, it is not guaranteed that red edge on extrasolar planets is detected as same wavelength on Earth. In fact, the photosynthetic organisms on Earth harvest light according to the surrounding environments to efficiently use light having a variety of wavelength that reaches, and of course their spectrum varies. In case of photosynthesis in extrasolar planets, photosynthetic organisms should evolve as optimized to utilize their principal star. We focus on the fundamental light harvesting mechanism and aim to propose how to detect the spectrum of the planet orbiting the different spectrum types of star otherwise Sun. At first, we adopted our models to photosynthetic organisms on Earth and compared with the experimental data. The light harvesting antenna in these organisms differs from kinds of pigments and their conformations.

We investigated the mechanism how the organisms harvest light by quantum mechanical calculation. However, because of the cost difficulty, we introduced an approximation instead of calculating all the electrons in the system. First, we calculated the excitation energies and the transition moments from the ground state to the excited states in the pigments by TDDFT (time-dependent density functional theory) [3]. Then, by introducing the transition moments of each pigment to the antenna, which consists of several kinds of pigments (other environment: proteins, solvent,...), we assume as an approximation that one pigment has the excited energy and interacts with the other pigments by the dipole-dipole interaction. When the light, as it seems to be considered as traveled from a star to a surface of an extrasolar planet, induces the system, we traced the time evolution of the energy transfer by solving Liouville equation. We dealt the light with an external potential. By this method, we can calculate the spectral intensity and energy efficiency.

In certain types of bacteria, the contrast like red edge can be detected, although the contrast is weaker than that of plants. In purple bacteria, red edge is not detected or emerges in longer wavelength (~1013-1025 nm). In addition, its structure of the antenna is simple so that we adopted easily our model to the bacteria. The calculated spectrum has a good agreement with the experimental result from purple bacteria. We will extend our model to the other species. By comparing the light harvesting mechanisms showing red edge and no red edge, we can examine how red edge emerges in photosynthetic organisms. For the other inducing light condition, we will survey the light harvesting mechanism optimized to extrasolar planets.

[1] N. Y. Kiang et al., *Astrobiology*, 7 (1), 252, 2007

[2] L. Arnold et al., *A&A*, 392, 231, 2002

[3] M. A. L. Marques and E. K. U. Gross, *Annu. Rev. Phys. Chem.*, 55, 427, 2004

Keywords: extrasolar planet, biomarker, quantum mechanical calculation

惑星間塵中の有機物とその生命起源との関連 Organic compounds in interplanetary dust particles and their relevance to origins of life

小林 憲正^{1*}, 川本 幸徳¹, 岡部 拓人¹, 金子 竹男¹, 大林 由美子¹, 神田 一浩², 三田 肇³, 藪田 ひかる⁴, 中川 和道⁵, 奥平 恭子⁶, 矢野 創⁷, 橋本 博文⁷, 横堀 伸一⁸, 山岸 明彦⁸

Kensei Kobayashi^{1*}, Yukinori Kawamoto¹, Takuto Okabe¹, Takeo Kaneko¹, Yumiko Obayashi¹, Kazuhiro Kanda², Hajime Mita³, Hikaru Yabuta⁴, Kazumichi Nakagawa⁵, Kyoko Okudaira⁶, Hajime Yano⁷, Hirofumi Hashimoto⁷, Shin-ichi Yokobori⁸, Akihiko Yamagishi⁸

¹ 横浜国立大学, ² 兵庫県立大学, ³ 福岡工業大学, ⁴ 大阪大学, ⁵ 神戸大学, ⁶ 会津大学, ⁷ JAXA 宇宙科学研究所, ⁸ 東京薬科大学

¹Yokohama National University, ²University of Hyogo, ³Fukuoka Institute of Technology, ⁴Osaka University, ⁵Kobe University, ⁶Aizu University, ⁷JAXA/ISAS, ⁸Tokyo University of Pharmacy and Life Science

原始および現在の太陽系環境下での惑星間塵中の有機物(アミノ酸・核酸塩基を含む)の安定性や変成に関して, シンクロトロン放射光等を用いた模擬実験を行った。その結果をもとに地球上での生命起源との関連を議論する。

キーワード: 惑星間塵, アミノ酸, 生命の起源, 複雑有機物, 軟 X 線, たんぽぽ計画

Keywords: interplanetary dust particles, amino acids, origins of life, complex organic compounds, soft X-rays, the Tanpopo Mission

Interdisciplinary Perspectives on Abiogenesis Interdisciplinary Perspectives on Abiogenesis

Hut Piet^{1*}

Piet Hut^{1*}

¹ プリンストン高等研究所, 地球生命研究所

¹Institute for Advanced Study, Earth-Life Science Institute, Tokyo, Japan

I will report on new interdisciplinary activities at the Institute for Advanced Study, in Princeton, NJ, USA, focused on the origins of life. We have recently started a series of weekly meetings, involving mathematicians, physicists, chemists, biologists, astrophysicists and others. Our discussions range from important technical details to more general meta-level discussions concerning the flow of information through biological networks and the characteristics of life in general, on Earth and possibly elsewhere in the Universe. We include scholars in humanities and linguistics, since there are interesting parallels between the origins of culture and the origins of life, and in general with the origins of "systems", where unreliable parts are used to create reliable systems.

My work is inspired by my affiliation with ELSI, the new Earth-Life Science Institute at Tokyo Tech, under the WPI program of MEXT.

キーワード: 生命の起源, 情報の流れ, 生物学的ネットワーク, 生命の特徴, システム

Keywords: origins of life, flow of information, biological networks, characteristics of life, systems

たんぽぽ計画における有機物曝露実験

Exposure experiments of organic compounds in space environments in the TANPOPO mission

三田 肇^{1*}, 横堀伸一², 橋本博文³, 東出真澄⁴, 今井栄一⁵, 河口優子², 河合秀幸⁶, 川本幸徳⁷, 神田一浩⁸, 小林憲正⁷, 中川和道⁹, 鳴海一成¹⁰, 奥平恭子¹¹, 田端誠³, 藪田ひかる¹², 山下雅道³, 矢野創³, 吉田聡¹³, 山岸明彦², たんぽぽ WG³
Hajime Mita^{1*}, Shin-ichi Yokobori², Hirofumi Hashimoto³, Masumi Higashide⁴, Eiichi Imai⁵, Yuko Kawaguchi², Hideyuki Kawai⁶, Yukinori Kawamoto⁷, Kazuhiro Knada⁸, Kensei Kobayashi⁷, Kazumichi Nakagawa⁹, Issay Narumi¹⁰, Kyoko Okudaira¹¹, Makoto Tabata³, Hikaru Yabuta¹², Masamichi Yamashita³, Hajime Yano³, Satoshi Yoshida¹³, Akihiko Yamagishi², Tanpopo WG³

¹ 福岡工大・工, ² 東京薬大・生命科学, ³ JAXA/ISAS, ⁴ JAXA・未踏技術研究セ, ⁵ 長岡技大・生物, ⁶ 千葉大・院理, ⁷ 横浜国大・院工, ⁸ 兵庫県立大・LASTI, ⁹ 神戸大・院人間発達環境, ¹⁰ 原子力機構・量子ビーム, ¹¹ 会津大, ¹² 阪大・院理, ¹³ 放医研
¹Fukuoka Inst. Technol., ²Sch. Life Sci., Tokyo Univ. Pharm. Life Sci., ³JAXA/ISAS, ⁴Innovative Tech. Res. Cen., JAXA, ⁵Nagaoka Univ. Tech., ⁶Grad. Sch. Sci., Chiba Univ., ⁷Grad. Sch. Eng., Yokohama Natl. Univ., ⁸LASTI, Univ. Hyogo, ⁹Grad. Sch. Human Develop. Environ., Kobe Univ., ¹⁰Quantum Beam, JAEA, ¹¹Univ. Aizu, ¹²Grad. Sch. Sci., Osaka Univ., ¹³NIRS

The Tanpopo mission is a Japanese astrobiological experiment which will be conducted on the Japanese Experiment Module (JEM) of the International Space Station (ISS). The Tanpopo mission consists of six subthemes: 1) capture of microbes in space, 2) exposure of microbes in space, 3) exposure of organic compounds in space, 4) capture of organic compounds (in micrometeoroids) in space, 5) evaluation of ultra low-density aerogel developed for the Tanpopo mission, and 6) capture of space debris at the ISS orbit (approximately 400 km altitude).

Here, we overview the exposure experiment of organic compound in space environment. Since many kinds of organic compounds, especially, amino acids which are ones of most important organic compounds in living organisms, are found from extraterrestrial materials, extraterrestrial and outer-solar environments are thought as the place for the prebiotic organic compound synthesis. Then, it is proposed that the first organisms on the earth was born from the prebiotic organic compounds delivered into the early earth on meteorites, micrometeorites and/or comets. In order to discuss the possibility of the hypothesis, alteration of prebiotic compounds in space environments should be clear. Therefore, we will expose some prebiotic organic compounds on the exposure facility at ISS-JEM.

Glycine, isovaline, hydantoin, ethylmethylhydantoin and complex organics (CAW) are chosen for the exposure. Hydantoin and ethylmethylhydantoin are plausible low molecular weight precursors for glycine and isovaline, respectively. CAW which is a simulated material of interstellar medium prepared by proton radiation into mixture of CO, NH₃ and H₂O is a different type of plausible precursors for amino acids. In the space environments, uv-light and cosmic rays (heavy ions and gamma-rays) will cause the alteration of organic compounds. Therefore, simulation experiments were studied using Xe-excimer lamp (uv 172 nm), synchrotron radiation at NewSUBARU BL06 (uv > 130 nm), 60Co gamma-ray radiation (JAEA Takasaki) and carbon ion beams (290MeV, NIRS). gamma-Ray and heavy ion beam irradiation with dose of ISS environment for one year induced little decomposition of organic compounds. However, uv irradiation was critical for organic compounds. Although almost all glycine and isovaline were decomposed, remains of hydantoin and ethylmethyl hydantoin were approximately 29% and 72%, respectively, with uv dose of ISS environment for one year. Furthermore, CAW was more stable than hydantoins. Amino acids precursors, especially, complex organics were more stable than free amino acids. Therefore, extraterrestrial amino acids precursors would be effective source for origins of life on the earth. We will demonstrate this conclusion on the ISS-JEM.

In addition, Nakagawa and his colleagues were found that dialanine was formed from alanine films by uv-irradiation. We will demonstrate a peptide synthesis with uv-irradiation in the space environment. Furthermore, piece of meteorite will be also exposed in order to examine the weathering effect in the ISS environment.

キーワード: たんぽぽ, 有機物曝露, 国際宇宙ステーション, アミノ酸

Keywords: Tanpopo, exposure of organics, ISS, amino acids

リボースの安定性に対するホウ酸の影響 Effects of borate on the stability of ribose

古川 善博^{1*}, 堀内 真愛¹, 掛川 武¹
Yoshihiro Furukawa^{1*}, Mana Horiuchi¹, Takeshi Kakegawa¹

¹ 東北大学理学研究科

¹ Graduate School of Science, Tohoku University

RNA is considered an important biomolecule for the origin of life, with its abilities as the carrier of genetic information and catalysts for biological reactions. RNA is composed of phosphoric acid, nucleic acid bases, and ribose. Among them, ribose is the most unstable component. Therefore, accumulation of ribose on the early Earth is an important step for the origin of RNA. Ribose as well as other aldopentoses can be abiotically synthesized by formose reaction in which formaldehyde oligomerize under alkaline conditions. Previous studies showed a stabilization of pentoses in the formose reaction by borate. In this study, we have investigated the effects of borate on the stability of aldopentoses.

Incubation experiments of each four aldopentoses were conducted at approximately 45 degree C with sodium tetraborate decahydrate of three concentrations. The pH of the experimental solution was buffered with calcium hydroxide. The experimental solution was collected at a fixed interval and analyzed with liquid chromatography-mass spectrometry.

In the borate-free experiments, all aldopentoses showed high decrease rates. In borate-containing experiments, formations of pentose-borate complexes were observed. The decrease rates for all aldopentoses were reduced with the concentration of borate. These results indicate that borate can stabilize all aldopentoses. Therefore, borate might have stabilized aldopentoses including ribose and contributed to the formation of primordial RNA on the early Earth.

キーワード: リボース, ホウ酸, RNA, 生命の起源

Keywords: ribose, RNA, borate, origin of life

オリゴペプチドによるホモキラリティー濃縮の経路 A Possible Pathway of Homochirality Accumulation by Oligopeptides

胸組 虎胤^{1*}
Toratane Munegumi^{1*}

¹ 鳴門教育大学大学院自然系コース(理科)

¹Department of Science Education, Naruto University of Education

多くの仮説が生体物質の片手構造を説明するために提案されてきたが、その多くはアミノ酸や単糖のような単量体の片手構造に関するものであった。しかしながら、単量体の片手構造は必ずしも、多量体中での片手構造を保証するものではない。なぜならば、原始地球および地球外には単量体を分解するような様々なエネルギー源が存在するからである。

本研究はオリゴペプチドによるポリペプチド中での片手構造の濃縮に焦点を当てている。ラセミ体のアミノ酸が2量化するとジアステレオマーが生成する。それがたとえば、アラニンであれば、鎖状、環状合わせて7種類のジアステレオマーが生じる。これらの化合物は化学構造ばかりでなく、疎水性、反応性、エピ化速度も異なる。一般に、ヘテロキラルペプチドはホモキラルペプチドよりも疎水性が高い。

発表では、オリゴペプチドによるポリペプチド中での片手構造の濃縮のスキームを、ジアステレオマーの分別、分解、立体特異的縮合に関連づけて示す。

キーワード: ホモキラリティー, 縮合反応, オリゴペプチド

Keywords: homochirality, condensation, oligopeptides

GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature
GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature

Kuhan Chandru^{1*}

Kuhan Chandru^{1*}

¹Faculty / Graduate School of Engineering Yokohama National University

¹Faculty / Graduate School of Engineering Yokohama National University

GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature

Kuhan Chandru, Eiichi Imai, Yumiko Obayashi, Takeo Kaneko and Kensei Kobayashi

GADV peptide hypothesis was proposed by Ikehara (2009) as a possible alternative that precedes the RNA world hypothesis (Gilbert, 1986) due to many limitations. The hypothesis suggest a very plausible explanation, but has only been experimentally tested by Oba and co workers (2012), in terms of possible peptide formation using heat-drying cycles and their catalytic activities. We would want to examine the formation of GADV aggregates (or peptides) in an simulated hydrothermal system to represent a more realistic prebiotic environment. We have used Gly, L-Asp, L-Asp and L-Val into our Supercritical Water Flow Reactor (SCWFR) at a temperature range of 100-300 Celsius . Our initial results suggest that no visible aggregates (bigger than 0.2micrometer) were seen after the heating of 2 min in the mentioned temperature range. Initial MALDI-TOF-MS are also suggesting that we only obtained a small peaks about m/z 410 within the spectrum lesser what Oba et al (2012) which was 525, 539, 657. Although many hydrothermal simulation experiments has shown the recovery of amino acids (Islam et al 2002; Kohara et al; Kobayashi et al, 1997) and the formation of oligomers (Imai, 1999; Goto et al , 2005) we only believe, that only small aggregation occur and cannot promote bigger oligomers or polymers due to heat and pressure stress. Hence, based on our initial findings, we are very uncertain about the formation of GADV aggregates or peptides in a hydrothermal system if we use free-form amino acids. It would be of interest to investigate the hypothesis by using bound amino acids or amino acid precursors.

References

Ikehara, K. (2009). Pseudo-Replication of [GADV]-Proteins and Origin of Life. *International Journal of Molecular Sciences*, 10(4), 1525-1537. doi:10.3390/ijms10041525

Gilbert, W (1986) The RNA world. *Nature* 319, 618

Oba, T., Fukushima, J., Maruyama, M., Iwamoto, R., & Ikehara, K. (2005). Catalytic activities of [GADV]-peptides. *Origins of Life and Evolution of the Biosphere : the Journal of the International Society for the Study of the Origin of Life*, 35(5), 447-460.

Kohara, M., Gamo, T., Yanagawa, H., & Kobayashi, K. (2003). Stability of amino acids in simulated hydrothermal vent environments. *Chemistry Letters*, 1-2.

Islam, M. N., Kaneko, T., & Kobayashi, K. (2003). Reaction of Amino Acids in a Supercritical Water-Flow Reactor Simulating Submarine Hydrothermal Systems. *Bulletin of the Chemical Society of Japan*, 76(6), 1171-1178. doi:10.1246/bcsj.76.1171

Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



BAO01-09

会場:301A

時間:5月21日 12:15-12:30

Kobayashi, K., Kohara., Dokiya, M and Yanagawa, H (1997) Formation and stability of amino acids in simulated hydrothermal vent environments. *Viva Origino* 25, 167 - 176

Imai, E.-I., Honda, H., Hatori, K., Brack, A and Matsuno, K (1999) Elongation of Oligopeptides in a Simulated Submarine Hydrothermal System. *Science*, 283(5403), 831-833. doi:10.1126/science.283.5403.831

Goto, T., Futamura, Y., Yamaguchi, Y., & Yamamoto, K. (2005). Condensation reactions of amino acids under hydrothermal conditions with adiabatic expansion cooling. *Journal of Chemical Engineering of Japan*, 38(4), 295-299.

キーワード: GADV, Hydrothermal, prebiotic

Keywords: GADV, Hydrothermal, prebiotic

数理モデルによる全球凍結直後の酸素濃度上昇過程の復元：マンガン及び鉄鉱床形成への示唆

Modeling the rise of oxygen after the Snowball earth: implications for the Paleoproterozoic manganese and iron formation

原田 真理子^{1*}, 田近 英一², 関根 康人², 尾崎 和海³

Mariko Harada^{1*}, Eiichi Tajika², Yasuhito Sekine², Kazumi Ozaki³

¹ 東京大学大学院地球惑星科学専攻, ² 東京大学大学院複雑理工学専攻, ³ 東京大学大気海洋研究所

¹Dept. of Earth and Planetary Science, The University of Tokyo, ²Dept. of Complexity Science & Engineering, The University of Tokyo, ³Atmosphere and Ocean Research Institute, The University of Tokyo

Earth's atmosphere and oceans are considered to have experienced stepwise and irreversible oxidation through its history (e.g. [1]), which may have paved the way to the complicated life such as eukaryotes and metazoans [2]. It is interesting to note that geological evidence suggests that a remarkable increase of oxygen concentration has occurred shortly after the Paleoproterozoic Snowball Earth event, based on widespread depositions of manganese and iron oxides immediately above the glacial diamictites found in the Transvaal Supergroup, South Africa, and the Huronian Supergroup, Canada [3,4]. Carbonate precipitation occurs above the deposition of manganese and iron in both the Transvaal and Huronian Supergroups [3,5], which may represent a climate recovery from the greenhouse condition in the Snowball earth aftermath [3]. In this study we numerically investigated the linkage between a global-scale glaciation and an oxygenation of the atmosphere-ocean system, in the aim of comparing our results to the geological records.

The results of our numerical experiments with a biogeochemical cycle model indicate that the super greenhouse conditions ($p\text{CO}_2 \sim 0.7 \text{ atm}$ and $T \sim 320 \text{ K}$) in the aftermath of the Paleoproterozoic Snowball Earth event significantly enhance the chemical weathering of continents, causing ~ 10 times as high as the present levels of nutrient input and the biological productivity. In the consequence of high levels of biological productivity together with a positive feedback in the atmosphere among a rise in oxygen, ozone formation, and UV shielding of methane, the atmospheric oxygen levels rapidly rise from $< 10^{-5} \text{ PAL}$ to 0.01 PAL (PAL: the present atmospheric level) after the glaciation. The oxygen levels then overshoot to $\sim 1 \text{ PAL}$ in $\sim 10^6$ years after the glaciation due to the high levels of biological productivity sustained by a long-term global warming. Atmospheric oxygen then gradually decreases by oxidizing reducing materials from Earth's interior. Eventually, a steady state of atmospheric oxygen of $\sim 0.01 \text{ PAL}$ is achieved in 10^8 years. Such an irreversible rise in atmospheric oxygen (i.e., from $< 10^{-5} \text{ PAL}$ to $\sim 0.01 \text{ PAL}$) is explained by a transition between different steady state of atmospheric oxygen levels [6] caused by a Snowball Earth glaciation and the subsequent perturbations of biogeochemical cycles.

We found that the rapid oxygenation causes the deposition of manganese and iron in the shallow marine environments. Manganese and ferrous iron in the anoxic deep water are driven by thermohaline circulation, immediately oxidized in the shallow marine water within 10^4 years after the glaciation. Assuming deep water is initially saturated with respect to Mn-carbonates, we derive 10^{15} mol of manganese deposition, which would be sufficient to form manganese ore in the Hotazel formation, Transvaal Supergroup ($\sim 8 \text{ Gt}$, $\sim 10^{14} \text{ mol}$ of manganese [7]). We also found that calcite precipitation is prevented in the ocean during 10^5 years after the glaciation owing to high atmospheric $p\text{CO}_2$. Our results imply that manganese and iron oxides deposition might precede the carbonate deposition, which is consistent with the geological records found both in the Transvaal and Huronian supergroups [3,4].

[1] Kasting (1993) *Science*, 259, 920-926. [2] Runnegar (1991) *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 97, 97-111 [3] Kirschvink et al. (2000) *Proc. Natl. Acad. Sci. USA*, 97, 1400-1405. [4] Sekine et al. (2011) *Earth Planet. Sci. Lett.*, 307, 201-210. [5] Bekker et al. (2006) *Precambrian Research*, 148, 145-180. [6] Goldblatt et al. (2006) *Nature*, 443, 683-686. [7] Tsicos (2003) *Economic Geology*, 98, 1449-1462

A Simple Astrobiological Scenario to go from the Great Oxygenation Event to the Origination of Eukaryotes

A Simple Astrobiological Scenario to go from the Great Oxygenation Event to the Origination of Eukaryotes

Joseph Kirschvink^{1*}

Joseph Kirschvink^{1*}

¹Geological and Planetary Sciences, California Institute of Technology, Pasadena CA, USA and, ²Earth/Life Science Institute, Tokyo Institute of Technology, Tokyo, Japan

¹Geological and Planetary Sciences, California Institute of Technology, Pasadena CA, USA and, ²Earth/Life Science Institute, Tokyo Institute of Technology, Tokyo, Japan

Taken in the context of Earth History, it is now clear that all previously reported Archean biomarkers indicative of oxygenic photosynthesis are most likely in error, a result of contamination during sampling. Reported geochemical whiffs of oxygen from ABDP-9 are also tightly associated with hydrothermal veins of highly crystalline clinocllore, and are thus metasomatic artifacts from hydrothermal vein deposits. Firm evidence for Archean oxygen simply does not exist.

A much simpler scenario will be presented here in which photosystem-II evolves at ~ 3.35 Ga, causing the radiation of the Cyanobacterial clade during the Gowganda glaciation. The resulting drawdown of CO₂, and destruction of CH₄, triggered the Makganyene global (snowball) glaciation, freezing the ocean surfaces for ~ 50 to 100 Myr, ending at 2.22 Ga. This is immediately followed by the deposition of the Kalahari Manganese Field, which is the oldest firm geological constraint on copious quantities of free O₂ in Earth's environment. As there must be a time gap between the evolution of Photosystem-II and the origin of oxidative phosphorylation (aerobic respiration), Earth's carbon cycle must have been extremely out of balance. We interpret the Lomagundi/Jatuli Carbon Isotope excursion (about 2.2 to 2.06 Ga) as precisely this imbalance, where the oxygen produced by photosynthesis, and organic matter produced, could not be easily recycled. The fraction of organic carbon buried increased from the long-term average of about 20% up to about ~ 75%, yielding isotopic compositions of carbonate as heavy as +15 per mil. Oxygen in the atmosphere is known to have increased to high levels ? perhaps several bars - resulting in the supergene alteration of the ~ 2.1 Ga iron ores in the Sishen deposit of South Africa during formation of the Hekpoort Paleosol. During this Lomagundi event, the high oxygen levels would be limited by the inability of cyanobacteria to release oxygen against high back pressures, and the remineralization of organic carbon would be accomplished by oxidative weathering of sulfide minerals releasing sulfate, and by the subsequent action of sulfate reducing bacteria to break down the organic carbon.

As the radiation of the Proteobacteria post-dates the radiation of the Cyanobacteria, and the ancestor of the Eukaryotic mitochondria was once a free-living alpha-Proteobacterium, it also follows that the first organelle-bearing eukaryotes must post-date the Makganyene Snowball Earth event. Numerous eukaryotic organelles have ancestry in this bacterial endosymbiont, as do cellular components such as the hydrogenosome in basal Protists.


A puzzling organelle present in both Bacteria and Eukaryotes (including many Protists) is the magnetosome, a membrane-bound, single crystal of biological magnetite (Fe₃O₄) or greigite (Fe₃S₄). Magnetosome chains orient single cells in the geomagnetic field and allow environmental positioning. Recent work on the genetics of biomineralization in magnetotactic bacteria (MB) has shown that both the greigite and magnetite genetic pathways share a common origin, supporting an earlier suggestion that the biochemistry of magnetotaxis is monophyletic.

Small subunit RNA sequence data from hundreds of extant MB are consistent with the origin of magnetotaxis within the Bacterial domain postdating the separation of stem-group cyanobacteria, but preceding radiation of the Proteobacteria (including the mitochondrial ancestor). We propose that the free-living mitochondrion ancestor was a magnetotactic bacterium. This hypothesis presents attractive implications for adopting cytoskeletal framework throughout the eukaryotic cell, for explaining the presence of magnetosomes throughout the Eukaryotes, and for justifying the demonstrated ability of the bacterial MagA protein to induce magnetite biomineralization in mammalian cells. The radiation of the Eukaryotic Domain of Life would therefore restore balance to the carbon cycle.

Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



BAO01-11

会場:301A

時間:5月21日 14:30-15:00

キーワード: Great Oxygenation Event, Cyanobacterial Evolution, Lomagundi Event, Mitochondrial Evolution, Magnetotactic Bacteria, Sedimentary Manganese

Keywords: Great Oxygenation Event, Cyanobacterial Evolution, Lomagundi Event, Mitochondrial Evolution, Magnetotactic Bacteria, Sedimentary Manganese

38億年前のグラファイトのSTEMによる観察 STEM observation of graphite from 3.8 Ga Isua Supracrustal Belt

掛川 武^{1*}

Takeshi Kakegawa^{1*}

¹ 東北大学院理

¹Tohoku University

Graphite is known to occur in the 3.8 billion years old Isua Supracrustal Belt (ISB) of western Greenland, and ¹³C-depleted graphite from sedimentary rocks of the ISB have been interpreted as traces of early life. The unequivocal documentation of biogenicity for graphite is complicated by the possibility of secondary graphite precipitation from metamorphic or igneous fluids and the difficulties in distinguishing biogenic from secondary graphite.

Graphite in meta-sedimentary rocks and carbonate veins in ISB is studied by STEM and Laser-Raman spectroscopy. Previous studies proposed that graphite in meta-sedimentary rocks are biogenic in origin, and others are chemical precipitates from CO₂-rich fluids. STEM observation of graphite in meta-sedimentary rocks indicated many unique textures, resemble to carbon nano-tube. On the other hand, graphite in veins show sheet-structure-dominated features. Laser-Raman analyses indicate that sedimentary graphite recorded the peak metamorphic temperature. Graphite in veins showed lower temperature than that of the peak metamorphism. Those data comprehensively provide more evidence of biogenic graphite that differ from non-biogenic graphite in ISB.

キーワード: STEM, ラマン

Keywords: STEM, Raman

Origin of life: Source rocks for the origin and evolution of life Origin of life: Source rocks for the origin and evolution of life

丸山 茂徳^{1*}, 戎崎 俊一²

Shigenori Maruyama^{1*}, Toshikazu Ebisuzaki²

¹ 東京工業大学 地球生命研究所, ² 理化学研究所

¹Earth-Life Science Institute, Tokyo Institute of Technology, ²RIKEN

One of the key factors for the origin and evolution of life is a nutrient supply which is derived from rocks. Life cannot be synthesized under nutrient-free conditions in an atmosphere as shown in the famous experiments of Miller (1953). For the beginning of life, three components are essential; (1) C (carbon)-centered sugar for fuel, (2) P (phosphorus)-centered metabolism, and (3) N (nitrogen)-centered information coded by basic pairs (DNA). Among them, P (negative ion), which is a centered nutrient coupled with K (positive ion), plays a critical role in metabolic activity. Can those nutrients be derived from any kinds of rocks on the Earth? The answer is, yes, but only three kinds of rocks as follows

(1) Granite: Granite can be formed by two-step extractions of nutrients: partial melting of mantle peridotite at a mid-oceanic ridge followed by partial melting at a subduction zone, either by slab melting or re-melting of lower mafic crust generated by partial melting of a mantle wedge. The major nutrient elements, such as P and K, are large ion lithophile elements (LILE), and hence, difficult to be bound into major mantle minerals. Plate tectonics increases the volume of calc-alkaline rocks such as TTG (tonalite-trondhjemite-granodiorite) or andesite and dacite at subduction zones through time.

Granitic rock was absent when a magma ocean was consolidated at 4.5Ga. In the Hadean, the major rock source for nutrient supply was not granite, but rather presumably the primordial continents.

(2) Primordial continents (Anorthosite with KREEP basalts): Primordial continental material is the second candidate for the source of nutrients. No remnants of those continents remain on the modern Earth (i.e., no Hadean rocks are left). Understanding of the primordial continent is developed through the geology of the Moon, and also from the concept of giant impact. The surface of Moon is covered by 50 to 70 km-thick anorthositic crust with local cover and dikes of KREEP (Potassium, Rare Earth Elements, and Phosphorus) basalt composition, and presumably underplated by KREEP-like rock types beneath the anorthositic continent. Both rock units have been interpreted as the final residue of magma ocean when the Moon was formed by the giant impact that led to the formation of the early Earth, where Mars-sized protoplanets collided with each other. If the giant impact theory is correct, the Earth must have been completely molten even up to the core. During the gradual cooling of the Moon and the Earth, the final liquid remained near the surface, forming the buoyant anorthositic crust, covered or underplated by KREEP magma similar to that observed on the Moon.

(3) Carbonatite: Carbonatite has only a single step of fractional melting of mantle peridotites under the cratons with an extremely small degree of melting. The selective removal of melt to form considerable amounts of nutrients under the sub-cratonic mantle creates carbonatite magma enriched in nutrients with highly volatile incompatible elements such as H₂O and CO₂ (more than 80% are volatiles). Nutrients concentrate into melt, depending on the degree of melting. Peridotite contains P=50 and K=240 (all values in ppm hereafter), but 100 times concentration of P=4495-3273 and 200-300 times of K=56118-68902 are seen in carbonatite. In general, the nutrient abundance is ideal for carbonatite, except for the U abundance. Carbonatite plays the role of milk-like materials to grow life. However, it may also function like an atomic-bomb magma to cause local mass extinctions, as well as resultant promotion of genome mutation by internal radiation through food chains.

火星生命探査計画：蛍光顕微鏡をもちいた火星表面での微生物探査 JAMP (Japan Astrobiology Mars Project): Search for Microbes on the Mars Surface with a Fluorescent Microscope.

山岸 明彦^{1*}, 吉村 義隆², 本多 元³, 宮川 厚夫¹, 佐藤 毅彦⁴, 石上 玄也⁴, 春山 純一⁴, 小林 憲正⁵, 長沼 毅⁶, 出村 裕英⁷, 佐々木 晶⁸, 宮本 英昭⁹

Akihiko Yamagishi^{1*}, Yoshitaka Yoshimura², Hajime Honda³, Atsuo Miyakawa¹, Takehiko Satoh⁴, Genya Ishigami⁴, Junichi Haruyama⁴, Kensei Kobayashi⁵, Takeshi Naganuma⁶, Hirohide Demura⁷, Sho Sasaki⁸, Hideaki Miyamoto⁹

¹ 東京薬科大学, ² 玉川大学, ³ 長岡科学技術大学, ⁴ 宇宙科学研究所, ⁵ 横浜国大, ⁶ 広島大学, ⁷ 会津大学, ⁸ 国立天文台, ⁹ 東京大学

¹Tokyo Univ. Pharm. Life Scie., ²Tamagawa University, ³Nagaoka University of Technology, ⁴ISAS/JAXA, ⁵Yokohama National Univ., ⁶Hiroshima Univ., ⁷Aizu Univ., ⁸NAOJ, ⁹The Univ. Tokyo

Introduction: Among the planets and giant satellites in our solar system, the characteristics of Mars are most similar to those of Earth. This suggests that the life similar to terrestrial life may arise and survive on Mars.

Here we propose a new life detection project on Mars within the scope of MELOS (Mars Exploration with Lander Orbiter Synergy), to search for methane-oxidizing microbes by fluorescence microscopy [1]. We propose to search for cells from a depth of about 5 - 10 cm below the surface, which is feasible with current technology. Microscopic observation has the potential to detect single cells. The subsequent analysis of amino acids will provide the information needed to define the origin of the cell.

Survivability of Life in the Mars Environment: Physical and chemical limits for terrestrial life have been major foci in astrobiology [2], and are summarized in ref. [1]. Combining the environmental factors, anywhere in the Martian environment where we can find the three components, water molecules, reducing compounds and oxidative compounds could be an environment where life can be sustained for long periods of time, if other factors such as temperature, pressure, UV and other radiations permit.

Methane Oxidizing Bacteria on Earth: Recently, a microbial consortium that is capable of using manganese (birnessite) and iron (ferrihydrite) to oxidize methane has been predicted in marine methane-seep sediments in the Eel River Basin in California [3]. Thus, there are several mechanisms of methane oxidation carried out by Bacteria and Archaea on Earth, and possibly on Mars.

Reference:

[1] Yamagishi, A, et al. (2010) Biol. Scie. Space, 24, 67-82. [2] Marion, G.M. et al. (2003) Astrobiol. 3, 785-811. [3] Beal, E. J., et al. (2009) Science, 325, 184-187.

キーワード: 火星, 微生物, 蛍光顕微鏡, 火星探査, アストロバイオロジー

Keywords: Mars, microbes, fluorescence microscope, Mars exploration, Astrobiology

Amazonis and Elysium basins and Their Link, Marte Vallis (AME), Tharsis/Elysium Corridor, Mars Amazonis and Elysium basins and Their Link, Marte Vallis (AME), Tharsis/Elysium Corridor, Mars

Dohm James^{1*}, 丸山 茂徳¹, 宮本 英昭²

James Dohm^{1*}, Shigenori Maruyama¹, Hideaki Miyamoto²

¹ 東京工業大学 地球生命研究所, ² 東京大学総合研究博物館

¹Earth-Life Science Institute, Tokyo Institute of Technology, ²The University Museum, The University of Tokyo

The geologic provinces of Mars, as identified through a synthesis of geologic, paleohydro-logic, topographic, geophysical, spectral, and elemental information [1], are windows into its evolution, with the youthful province, the Tharsis/Elysium corridor [2-4], recording hydrologic, tectonic, and volcanic activity, including fissure-fed eruptions and shield-volcano field development, within at least tens of millions of years.

Geologically recent activity in the Tharsis/Elysium corridor region is particularly high-lighted in the Amazonis and Elysium basins and their link, Marte Vallis (hereafter referred to as AME). A youthful shield-volcano field with at least seven gentle sloping shield volcanoes and fissure-fed vent structures have been identified, mapped, and characterized in the western part of the Elysium. In addition, the lava flows on and near the margins of the shield volcanoes display crenulated lava flow margins, possibly marking lava-water-ice interactions. Both the shield volcanoes and pristine lavas located within AME, which are marked by a paucity of relatively small (~few-km-diameter) superposed impact craters and modified by faults and fractures and valleys, point to geological and hydrological activity on Mars in recent geologic time, making AME a significant target for future reconnaissance, including testing the hypotheses of whether Mars is geologically, hydrologically, and biologically active.

Specifically, the following questions might be addressed through international reconnaissance missions to AME, which would include instrument suites with optimal geologic, geochemical, geophysical (including seismic), environmental, and biological capabilities, including whether Mars: (1) is geologically and hydrologically active, (2) contains salty groundwater and magma at relatively shallow depths, (3) has sustained elevated heat flow, (4) records seismic activity, and (5) comprises fossilized and/or extant life.

References

[1] Dohm, J.M. et al., (2013?in press), Nova Science Publishers, Inc.

[2] Dohm, J.M., et al., 2008, *Planetary and Space Science* 56, 985?1013, doi:10.1016/j.pss.2008.01.001

[3] Dohm, J.M., (2004) *Planetary and Space Science* 52, 189?198, doi:10.1016/j.pss.2003.08.013.

[4] Miyamoto, H., (2004) *Journ. of Geophy. Res.* 109, E06008, doi:10.1029/2003JE002234.

たんぽぽ計画の準備状況と微生物宇宙生存可能性の検討

Current status of preparation of TANPOPO mission and investigation of survivability of microbes in space

横堀 伸一^{1*}, 河口 優子¹, 橋本 博文⁷, 林 宣宏⁶, 東出 真澄⁷, 今井 栄一⁸, 河合 秀幸⁹, 小林 憲正¹⁰, 三田 肇¹¹, 中川 和道², 鳴海 一成³, 奥平 恭子¹², 清水 康之¹, 田端 誠⁹, 藪田 ひかる¹⁴, 太刀川 純孝⁴, 富田一横谷 香織¹³, 山下 雅道⁴, Yang Yinjie¹, 矢野 創⁴, 吉田 聡⁵, 山岸 明彦¹

Shin-ichi Yokobori^{1*}, Yuko Kawaguchi¹, Hirofumi Hashimoto⁷, HAYASHI, Nobuhiro⁶, HIGASHIDE, Masumi⁷, Eiichi Imai⁸, KAWAI, Hideyuki⁹, Kensei Kobayashi¹⁰, Hajime Mita¹¹, NAKAGAWA, Kazumichi², Issay Narumi³, Kyoko Okudaira¹², SHIMIZU, Yasuyuki¹, TABATA, Makoto⁹, Hikaru Yabuta¹⁴, TACHIKAWA, Sumitaka⁴, Kaori Tomita-Yokotani¹³, YAMASHITA, Masamichi⁴, Yinjie Yang¹, Hajime Yano⁴, YOSHIDA, Satoshi⁵, Akihiko Yamagishi¹

¹ 東京薬大・生命科学, ² 神戸大・院人間発達環境学, ³ 原子力機構・量子ビーム, ⁴ JAXA/ISAS, ⁵ 放医研・福島復興支援, ⁶ 東京工大・院生命理工, ⁷ JAXA・未踏技術研究セ, ⁸ 長岡科技大・生物, ⁹ 千葉大・院理, ¹⁰ 横浜国大・院工, ¹¹ 福岡工大・工, ¹² 会津大, ¹³ 筑波大・院生命環境, ¹⁴ 大阪大・院理

¹Sch. Life Sci., Tokyo Univ. Pharm. Life Sci., ²Grad. Sch. Human Develop. Environ., Kobe Univ., ³Quantum Beam, JAEA, ⁴JAXA/ISAS, ⁵NIRS, ⁶Grad. Sch. Biosci. Biotech., Tokyo Inst. Tech., ⁷JAXA, ⁸Nagaoka Univ. Tech., ⁹Grad. Sch. Sci., Chiba Univ., ¹⁰Grad. Sch. Eng. Yokohama Natl. Univ., ¹¹Fac. Eng., Fukuoka Inst. Tech., ¹²Univ. Aizu, ¹³Grad. Sch. Life Environ. Sci., Univ. Tsukuba, ¹⁴Grad. Sch. Sci., Osaka Univ.

Microbes have been collected from high altitude using balloons, aircraft and meteorological rockets since 1936, even it is not clear how those microbes could be ejected up to such high altitude. Spore forming fungi, spore-forming Bacilli, and Deinococci (e.g. *Deinococcus aerius* and *Deinococcus aetherius*) have been isolated in these experiments. If microbes could be found even at the higher altitude of low earth orbit (400 km), the fact would endorse the possibility of interplanetary migration of terrestrial life.

For the origin of life on Earth emerged within a short period after the end of heavy bombardment, Panspermia hypothesis was proposed. Recent the reports on the possible fossils of microbes in the Martian meteorite promote the debate on the possible existence of extraterrestrial life, and interplanetary migration of life as well.

On the other hand, it is the question where precursors of materials such as protein and nucleic acids came from in the era of "chemical evolution" on the Earth? Recent studies suggest that the some of such organic compounds were created in space. Then, they reached the surface of Earth via meteorites, cosmic dusts, and so on. Avoiding contamination of terrestrial materials from the extraterrestrial materials is quite important issues for the analysis of extraterrestrial materials. Capturing such extraterrestrial materials before falling down on the surface of Earth might be one of possible solutions.

We have proposed a mission, named TANPOPO, to examine possible interplanetary migration of microbes, and organic compounds at the Exposure Facility of Japan Experimental Module (JEM) of the International Space Station (ISS). The Tanpopo mission consists of six subthemes ? capture of microbes in space, exposure of microbes in space, capture of organic compounds in space, exposure of organic compounds in space, measurement of space debris at the ISS orbit, and evaluation of ultra low-density aerogel special for the TANPOPO mission.

Ultra low-density aerogel will capture micrometeoroid and space debris. Particles captured by aerogel will be analyzed from biological, chemical, and meteorological aspects.

In addition to particle-capture on ISS, we also proposed direct exposure experiments of microbial cell aggregates that might protect the microbes themselves from UV and cosmic rays. Deinococci (*Deinococcus radiodurans*, *D. aerius*, and *D. aetherius*), terrestrial cyanobacteria, and fungi are under consideration for space exposure. Amino acids and complex organic compounds that can be formed in space are also planned for space exposure.

In this paper, we overview the TANPOPO mission and discuss the current status of experiments related to the microbe existence/survival set for this mission.

キーワード: パンスペルミア, 宇宙曝露, 生命の起原

Keywords: Panspermia, Space exposure, Origin of life

サブミリメートルサイズの*Deinococcus*属の凝集細胞は宇宙空間でUVを遮蔽することが可能である

The sub-millimeter-sized aggregated deinococcal cells could be shield from solar UV

河口 優子^{1*}, Yinjie Yang¹, 川尻成俊¹, 白石啓祐¹, 高須昌子¹, 鳴海一成², 佐藤勝也², 橋本 博文³, 中川和道⁴, 谷川能章⁴, 桃木洋平⁴, 田邊真依子⁴, 杉野朋弘¹, 高橋勇太¹, 清水康之¹, 吉田聡⁵, 小林 憲正⁶, 横堀 伸一¹, 山岸 明彦¹
Yuko Kawaguchi^{1*}, Yinjie Yang¹, Narutoshi Kawashiri¹, Keisuke Shiraishi¹, Masako Takasu¹, Issay Narumi², Katsuya Satoh², Hirofumi Hashimoto³, Kazumichi Nakagawa⁴, Yoshiaki Tanigawa⁴, Yoh-hei Momoki⁴, Maiko Tanabe⁴, Tomohiro Sugino¹, Yuta Takahashi¹, Yasuyuki Shimizu¹, Satoshi Yoshida⁵, Kensei Kobayashi⁶, Shin-ichi Yokobori¹, Akihiko Yamagishi¹

¹ 東京薬科大学生命科学, ² 日本原子力機構量子ビーム, ³ JAXA/ISAS, ⁴ 神戸大学大学院人間発達科学, ⁵ 放射線医学研究所福島復興支援, ⁶ 横浜国立大学大学院工学府, ⁷ 東京都医学総合研究所

¹Department of Applied Molecular Biology, School of Life Sciences, Tokyo University of Pharmacy and L, ²Ion Beam Mutagenesis Research Group, Quantum Beam Science Directorate, ³Institute of Space and Astronautical Science, JAXA, ⁴Graduate School of Human Development and Environment, Kobe University, ⁵National Institute of Radiological Sciences, ⁶Graduate School of Engineering, Yokohama National University, ⁷Protein Metabolism Project, Tokyo Metropolitan Institute of Medical Science

To investigate the interplanetary transfer of life, numerous exposure experiments have been carried out on various microbes in space since 1960s. The results suggested that microbe spores might survive for a long period if the spores are shielded from intense solar radiation [1]. In the Tanpopo mission, we have proposed to carry out the experiments on capture and space exposure of microbes at International Space Station (ISS) [2]. Microbial candidates for the exposure experiments in space include *Deinococcus radiodurans*, *D. aerius* and *D. aetherius*. We have examined the survivability of *Deinococcus* spp. under the environmental conditions on ISS in orbit (i.e., long exposure to heavy-ion beams, temperature cycles, vacuum and UV irradiation). Among the space environmental factors, solar UV is most lethal to microbes, and damage is caused by the absorption of UV by DNA [3].

In this report, we examined the effect of solar UV radiation (172 nm, 254 nm and 280-315 nm respectively) on the deinococcal cell aggregates with different thicknesses to determine whether the size of the cell aggregate influences the cell survivability. Though the cells in thin layers of aggregates were killed by UV radiation, large number of cells survived the radiation when the cell layer was thick. The similar trend of survivability was observed for other UV range. Supposing that the aggregates are sphere, the diameter of the aggregate that is sufficient to shield the cells in the inner layer from solar UV radiation is 200 micrometer for *D. radiodurans*, 850 micrometer for *D. aerius*, and 700 micrometer for *D. aetherius*. We propose the microbial cell aggregate as an ark for the interplanetary transfer of microbes, and name it the 'masspanspermia' hypothesis.

[References]

[1] Horneck et al., (1994) Adv. Space Res. 14, 41-45 [2] Yamagishi et al., (2008) Int. Symp. Space Tech. Sci. (ISTS) Web Paper Archives 2008-k-05 [3] Horneck (1993) Orig. Life Evol. Biosph. 23, 37-52

キーワード: パンスペルミア仮説, 宇宙実験, *Deinococcus*, 微生物凝集体, マサパンスペルミア仮説

Keywords: Panspermia hypothesis, Space exposure experiments, *Deinococcus*, Cell aggregation, Masspanspermia hypothesis

復元した祖先型タンパク質を用いた単純型アミノ酸組成の検討 Restricting the amino acid usage of a resurrected ancestral protein

赤沼 哲史^{1*}, 島田真実¹, 中島慶樹¹, 横堀伸一¹, 山岸明彦¹
Satoshi Akanuma^{1*}, Masami Shimada¹, Yoshiki Nakajima¹, Shin-ichi Yokobori¹, Akihiko Yamagishi¹

¹ 東京薬科大学 生命科学部 応用生命科学科

¹Department of Applied Bioscience, Tokyo University of Pharmacy and Life Sciences

It is common among life on Earth that proteins are composed of 20 or nearly 20 kinds of amino acids. However, it is uncertain that, from the standpoint of astrobiology, proteins are always composed of the 20 kinds of amino acids. Even for the evolution of proteins on Earth, it has been argued that primordial proteins that existed before the last universal common ancestor involved less than 20 kinds of amino acids (Crick, 1968). Given simpler protein synthesis system, the primitive proteins, which might have comprised a reduced set of amino acids, must have had a sufficiently adequate structure for functional interactions and catalysis. To address this issue experimentally, we used the protein simplification engineering (Akanuma et al., 2002) to examine whether a protein composed of less than 20 types of amino acids can retain its stable structure and biological function. To this end, we first resurrected several ancestral proteins and then restricted the amino acid usage of a resurrected protein to a reduced amino acid set.

As the model protein, we chose a house keeping enzyme, nucleoside diphosphate kinase (NDK). Because extant genes are evolutionary descendants of ancient genes, ancestral sequences of a particular protein can be inferred by comparing extant homologous protein sequences. Along this line, we first inferred several ancestral amino acid sequences of NDK by phylogenetic analysis of the extant homologous sequences. The inferred sequences were then genetically reconstructed. One of the reconstructed protein, Arc1, is very thermally stable (unfolding temperature = 113°C) and shows catalytic efficiency similar to those of the modern NDKs.

It is currently impossible to infer amino acid sequences that existed before the last universal common ancestor. Therefore, using Arc1 as the starting molecule, we reconstructed a reduced amino acid set variant, Arc1-s2, in which Met, Gln, Lys, Tyr and Asn were replaced by other amino acids. Because cysteine is absent from Arc1, Arc1-s2 consists of only 14 amino acid letters. Arc1-s2 retains thermal stability similar to that of Arc1; whereas, no detectable level of catalytic activity was observed for Arc1-s2. Therefore, the fourteen amino acid types are sufficient to encode a thermally stable protein but more amino acid types would be required for its function. To reevaluate the individual contributions of the 20 amino acid types to the stability and activity of Arc1, we reconstructed 19 Arc1 variants in which one of the 19 amino acid types was all replaced by other amino acids. The result will be also present in the meeting.

(1) Crick, JMB 38, 367 (1968); (2) Akanuma et al., PNAS 99, 13549 (2002)