

Calcification mechanisms in foraminifera and proxy incorporation Calcification mechanisms in foraminifera and proxy incorporation

Bijma Jelle^{1*}, Nehrke Gernot¹, Raitsch Markus¹, ド・ノイエ レナート², Funcke Antje¹, ケウル ニイナ³
Jelle Bijma^{1*}, Gernot Nehrke¹, Markus Raitsch¹, Lennart Jan de Nooijer², Antje Funcke¹, Nina Keul³

¹ アルフレッドウェゲナー研究所, ²NIOZ, ³Lamont-Doherty Earth Observatory

¹AWI, ²NIOZ, ³Lamont-Doherty Earth Observatory

Calcifying organisms, such as pteropods, bivalves, corals and foraminifera provide a rich resource for pale-oceanographers and climatologists because their geochemical make-up (proxies) can be used to reconstruct past ocean history and evolution during and after natural carbon perturbations. However, it has been shown for all geochemical proxies that the main assumption of only one environmental variable controlling a target proxy is too simple. Empirical calibrations introduce a lot of uncertainty because the mechanisms of proxy incorporation are not well understood. The major problem is that the calcification mechanisms are still a black box. In this presentation I will review our current understanding of calcification and proxy incorporation in foraminifera.

キーワード: 石灰化

Keywords: calcification, foraminifera

有孔虫が石灰化する時の環境変動

Environmental variation during foraminiferal calcification

豊福 高志^{1*}

Takashi Toyofuku^{1*}

¹ 海洋研究開発機構

¹JAMSTEC

Foraminiferal calcification process will be indicated with pH and Calcium variations.

キーワード: 有孔虫, 石灰化

蛍光色素を用いた共生藻をもつ有孔虫の室形成過程の解明

Distribution of intracellular pH and algal endosymbionts during chamber formation in symbiont-bearing reef foraminifers

藤田 和彦^{1*}, 由地 蛍¹, 豊福 高志²

Kazuhiko Fujita^{1*}, YUJI, Kei¹, Takashi Toyofuku²

¹ 琉球大学 理学部, ² 海洋研究開発機構

¹Univ. Ryukyus, ²Biogeos, JAMSTEC

Calcification mechanisms of algal symbiont-bearing reef foraminifers have not yet been well understood. In particular, it is cytologically not clear how the photosynthesis by algal endosymbionts enhance the calcification of host foraminifers. We visualized the distribution of intracellular pH and algal endosymbionts using a fluorescence probe HPTS as well as chlorophyll autofluorescence. High pH vesicles were gradually stored around symbionts before chamber formation. Our observations suggest that the photosynthesis by symbionts enhances the production of calcite needles or carbonate pools in vesicles by removing CO₂ around them.

Autoclave aquaria allow for high-pressure culture experiments on deep-sea benthic foraminifera
Autoclave aquaria allow for high-pressure culture experiments on deep-sea benthic foraminifera

Wollenburg Jutta^{1*}
Jutta Wollenburg^{1*}

¹ アルフレッドウェゲナー研究所
¹AWI

Our understanding of palaeodeep-water circulation modes and deep-water renewal in the world's oceans is essentially based on isotopical and geochemical ratios recorded in tests of calcareous deep-water benthic foraminifera that precipitate their shells in a constant ratio to the surrounding water mass. However, as more field data are assembled it appears that the documentation of deep-water in benthic foraminiferal shells is not always straightforward. Therefore, culture experiments on deep-sea benthic foraminifera are needed to verify the established palaeodeep-water proxies. However, to our knowledge, barophilic species like *F. wuellerstorfi*, the most trusted species for reconstructions of palaeodeep-water circulation and ventilation, neither formed new chambers nor reproduced in mesocosms kept at 1 bar. To accommodate this problem we have developed facilities and procedures that allow long-lasting high-pressure culture experiments on undepressurized deep-sea sediments and associated fauna and flora. In this presentation I will describe the experimental set-up and present results from methane-seepage simulations.

キーワード: 圧力, 飼育, 有孔虫
Keywords: pressure, culture, foraminifera

飼育実験手法を基にした溶存酸素濃度の浮遊性有孔虫への影響評価
Laboratory culture experiments: effect of dissolved oxygen concentration on planktonic foraminifera

黒柳 あずみ^{1*}, da Rocha Regine E.², Bijma Jelle², Spero Howard³, 川幡 穂高¹
Azumi Kuroyanagi^{1*}, Regine E. da Rocha², Jelle Bijma², Howard Spero³, hodaka kawahata¹

¹ 東京大学 大気海洋研究所, ² アルフレッドウェゲナー極域海洋研究所, ³ カリフォルニア大学 デービス校

¹ Atmosphere and Ocean Research Institute, The University of Tokyo, ² Alfred-Wegener-Institut für Polar- und Meeresforschung,

³ University of California Davis

Planktonic foraminifera shows significant species turnover ratio related Cretaceous oceanic anoxic event (OAEs), however, the direct effect of dissolved oxygen (DO) concentration on planktonic foraminifera remain obscure. Although culture experiments can investigate quantitatively the relationship between foraminiferal ecology and environmental parameters, DO-controlling experiment has never been conducted because of the difficulty of observation and/or control of dissolved oxygen under the modern ocean condition. In the present study, we cultured two planktonic foraminifera species (*Orbulina universa* and *Globigerina bulloides*) under six different DO conditions. Both species have extremely high tolerance to low DO than we expected before, and it suggests that at least "dysoxic" condition (more than 0.7 mg O₂ l⁻¹) could not be a direct cause of the extinction of planktonic foraminifera at OAEs. Their high tolerance to extremely low DO would be caused by the evolutionary descendant of benthic foraminifera. Final shell weight increased with DO despite almost the same culture duration among treatments, thus it suggests foraminiferal fossil shell weight could reflect the DO conditions.

キーワード: 浮遊性有孔虫, 飼育実験, 溶存酸素濃度

Keywords: planktonic foraminifer, culture experiment, dissolved oxygen concentration

亜北極・北極 *Emiliana huxleyi* のココリス形態の培養温度に対する応答 Subarctic and Arctic *Emiliana huxleyi* coccolith morphological responses to the growth temperatures

猿渡 和子^{1*}, 佐藤真奈美¹, 原田尚美², 鈴木石根¹, 白岩善博¹

Kazuko Saruwatari^{1*}, Manami Satoh¹, Naomi Harada², Iwane Suzuki¹, Yoshihiro Shiraiwa¹

¹ 筑波大学生命環境科学, ² 海洋研究開発機構

¹ Faculty of Life and Environmental Sciences, University of Tsukuba, ² Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology

Coccolithophore is a key calcified phytoplankton for biogeoscience because of their specific products such as calcified scales, coccolith, and unsaturated ketones, alkenone. As a biomarker, the chemical compositions of alkenones are often used to estimate paleo ocean temperatures and to identify the changes of the marine environments (e.g., Harada et al., 2012). In particular, the polar region is an important area to recognize the changes of the global environments and the sea-ice melting is known to be accelerated by the global warming. MIRAI subarctic expeditions have been continued to investigate the subarctic to Arctic environment and concurrently the culture experiments of marine organisms including coccolithophores have been performed to predict the future changes of marine ecosystem. In this study, two *Emiliana huxleyi* strains, MR57N and MR70N, isolated by MIRAI subarctic expedition in 2010 were cultured at the different temperatures (5, 10, 15 and 20°C) and salinities (25, 32, and 35 per mill) and investigated from the viewpoint of the coccolith morphology. As the results, these strains showed the similar growth properties that the growth rate increased with raising the temperature. According to SEM observations and the image analyses, the size (length of distal shield: LDS) and the numbers of distal shield elements of the coccolith decreased with raising the temperature. The central area of the coccoliths was also changed from grill structures to completely calcified structures. Moreover, the cell size of *E. huxleyi* decreased with raising the temperature. Thus, the subarctic and Arctic *E. huxleyi* strains showed the correlations between cell sizes and coccolith morphometric parameters with variable central area morphology depending on the growth temperatures. For the salinity experiments, there is almost no variation on the growth properties and a little variation on the morphometrical parameters. These results imply that the subarctic and arctic *E. huxleyi* strains can maintain enough to grow in the temperature range between 5 to 20°C and the salinity range between 25 to 35 per mill, suggesting that these strains can ideally survive under the warmer and/or the less salinity environments. The coccolith morphologies and the parameters vary in response to the growth temperature but not to the salinity. This relationship was applied to the natural plankton samples reported by Bollmann et al. (2009) and the part of their data was correlated with sea water temperatures but the data shallower than 15 m depth was not. This may imply the importance of the light intensity and the further culture experiments with the different light intensity are required.

Harada et al. (2012) *Global Biogeochemical Cycles* 26, GB2036.

Bollmann et al. (2009) *EPSL* 284, 320.

放散虫 *Spongotrochus glacialis* にみられる 2 隠蔽種の形態的特徴及び好適環境要因
Morphological and ecological characters of two cryptic genetic types in the radiolarian
Spongotrochus glacialis.

石谷 佳之^{1*}, 氏家 由利香², 瀧下 清貴¹
Yoshiyuki Ishitani^{1*}, Yurika Ujii², Kiyotaka Takishita¹

¹ 海洋研究開発機構 深海生態系研究チーム, ² 高知大学海洋コア総合研究センター

¹Japan Agency for Marine-Earth Science and Technology, ²Center for Advanced Marine Core Research, Kochi University

The discovery of cryptic species often helps in grasping the true diversity and detailed ecological preferences of pelagic planktons. Radiolarians are a good environmental proxy for paleontology because of their worldwide oceanic distribution and the good preservation of their shells. However, cryptic species have not yet been recognized in this planktonic protist group. In the present study, we focused on the morphospecies *Spongotrochus glacialis* with specimens collected from different layers in the Equatorial-Subtropical Pacific. A molecular phylogeny based on the internal transcribed spacer of rDNA (ITS-rDNA) sequences recovered two clades of cryptic species (types I and II) from this single morphospecies. These two distinct types were separately distributed, either in the oligotrophic surface water (type I) or below the chlorophyll maximum layer (type II). Moreover, the types showed morphological differences in the shells. Our morphometric analyses established lengths of spines as a morphological criterion to distinguish between the two types: type I with longer spines and type II with shorter spines. The length of spines is apparently associated with the habitat of each type. Type I with longer spines could be suitable for extending flagella and floating on the surface, whereas type II with shorter spines are appropriate to protrude only short flagella and dwell in deep water. Such morphological and ecological features at the cryptic species level of Radiolaria could provide new proxies for paleoceanographic studies.

キーワード: 放散虫, 隠蔽種, ITS rDNA, 形態計測

Keywords: Radiolaria, cryptic species, ITS rDNA, morphometric

南シナ海北部の環境DNA中の放射虫類18S rDNAの系統解析

The Phylogenetic Affiliations of Radiolarian-like Environmental 18S rDNA Genes from the Northern South China Sea

李 緑硯^{1*}, 遠藤一佳¹

Luyan Li^{1*}, Endo Kazuyoshi¹

¹ 東京大学

¹The University of Tokyo

To determine the molecular taxonomic affiliations of radiolarian-like environmental 18S rDNA genes from the northern South China Sea, we constructed phylogenetic trees by the 18S rDNA sequences of identified radiolarian species, correlating environmental sequences from other sea areas with those full length 18S rDNA gene of representative clones from the northern South China Sea. In our previous work, a great number of our radiolarian-like sequences from the South China Sea were placed within the RAD-III, named by Not et al. (2007) of environmental sequences from the Sargasso Sea size-fractionated samples (< 2 micrometre). Now we found 18S rDNA sequences of two identified species *Arachnosphaera myriacantha* and *Astrosphaera hexagonalis*, collected from Okinawa Island by Yuasa et al. (2009) also belongs to RAD-III. There is a unique clade composed solely of environmental clone sequences formed basal to the Taxopodida. It does not belong to RAD IV or V, still remains unknown. Our *Nassellaria*-like environmental clone from the northern South China Sea was joined with the group of described *Plagoniid* species. And the clones from Cariaco Basin, Caribbean clustered with *Pterocorythid* and *Theoperid* groups. Comparison of our sequences with two recent survey of *Acantharia* 18S rDNA, the position of northern South China Sea clones was still difficult to distinguish. It belonged to the clade composed of subgroups *Arthracanthida*, *Sphaenacanthida* and RAD I (Unidentified Clade 1). These ambiguities may be because the genetic diversity of radiolarian in warm waters sea areas is still poorly known.

シアノバクテリア化石の顕生代海洋における pH プロキシとしての可能性 Fossil cyanobacteria as a potential pH proxy for Phanerozoic ocean

白石 史人^{1*}

Fumito Shiraishi^{1*}

¹ 白石史人

¹SHIRAISHI, Fumito

Fossil cyanobacteria appeared in the most part of Phanerozoic era, and its abundance changed significantly through time. Their fossilization related to calcification in the living state, which requires optimum chemistry in the ambient water. Therefore, the fossil record of cyanobacteria potentially reflects past ocean chemistry. Recent study revealed that cyanobacterial calcification depends mainly on pH, dissolved inorganic carbon, Ca^{2+} concentration (Shiraishi 2012, GCA). Using well-established proxies of Ca^{2+} and partial pressure of CO_2 , oceanic pH was estimated from the fossil record of calcified cyanobacteria. Estimated range and trend of pH are similar to those of previous studies, but exhibited more frequent changes. In a future study, it is necessary to understand the relationship between CaCO_3 saturation state achieved by photosynthesis and calcification amount, in order to increase the reliability of estimation.

Benthic foraminifera as Quaternary palaeoclimate proxies in the New Caledonia Basin Benthic foraminifera as Quaternary palaeoclimate proxies in the New Caledonia Basin

Briony Mamo^{1*}, Glenn Brock², Elsie Gretton³

Briony Mamo^{1*}, Glenn Brock², Elsie Gretton³

¹JAMSTEC, ²Macquarie University, ³Boggabri Coal

¹JAMSTEC, ²Macquarie University, ³Boggabri Coal

In 2001, gravity core GC4 was extracted from the New Caledonia Basin and the uppermost 141 cm, representing approximately the last 140 000 years, was investigated for chemical isotopic, carbonate, non-carbonate and trace element signatures. In this study, sub samples of the same uppermost 141 cm of GC4 were picked and sorted for all foraminiferal taxa. A total of 161 species of benthic foraminifera were obtained from the core and 46 species occurred in sufficient numbers to investigate changing patterns and trends in biodiversity and relative abundance. All changes can be related to broad scale oceanic and palaeoclimatic fluctuations during the last 140 000 years. Within GC4, two distinct foraminiferal assemblages were detected using Bray-Curtis cluster analysis and Multidimensional Scaling (MDS) analysis. Factors that influence the foraminiferal assemblages include environmental setting, SSP, oxygen levels at the sediment-water interface and transportation by oceanic current systems operating within the region.

キーワード: foraminifera

Keywords: foraminifera

南日本で採集した化石シャコガイ殻のSr/Ca比を用いた4.6kaの日射量の復元
Daily light cycle reconstructed by Sr/Ca in a fossil giant clam, *Tridacna gigas*, living in
4.6 ka, southern Japan

堀 真子^{1*}, 白井厚太郎¹, 高畑 直人¹, 佐野 有司¹, 小林夏子², 岡本 和明², 渡邊 剛³

Masako Hori^{1*}, Kotaro Shirai¹, Naoto Takahata¹, Yuji Sano¹, Natsuko Kobayashi², Kazuaki Okamoto², Tsuyoshi Watanabe³

¹ 東京大学大気海洋研究所, ² 埼玉大学教育学部, ³ 北海道大学理学部

¹Atmosphere and Ocean Research Institute, the University of Tokyo, ²Faculty of Education, Saitama University, ³Faculty of Science, Hokkaido University

Giant clams are long-lived bivalvia living in a shallow tropical ocean. Their aragonite shells are potential paleo-environmental archives, i.e., oxygen stable isotopic values record paleo-seawater temperatures. Generally, trace elements in calcium carbonates are also considered as a paleo-thermometer. However, the case of Tridacninae may not follow the traditional temperature dependent fractionation, but potentially reflects insolation. This is explained by "trans-calcification", which is one of photosynthetically induced calcification mechanisms. Enzymatic reaction pumping Ca^{2+} to host liquid of calcification results in relatively depleted Sr/Ca ratio in day time when photosynthesis activates. Following this hypothesis, we have reconstructed the past daily light cycle.

A fossil *T. gigas*, Stg04-b was collected at Ishigaki-jima Island in Okinawa Prefecture, Japan (N24 20'0.4" E124 09'22"). Two horizons of the specimen were dated by carbon-14 method at the accelerator mass spectrometry (AMS) center of Yamagata University. The micro analyses were performed for the outer layer of 1.4 cm thickness using an electron probe micro-analyzer (EPMA), JXA8900, and a micrometer-scale secondary ion mass spectrometer (NanoSIMS), CAMECA NS50 at the Atmosphere and Ocean Research Institute (AORI), the University of Tokyo. As a result, we found clear Sr lamination parallel to the daily increment of about 20 μm thickness. The hourly insolation (P) was calculated from the difference of Sr/Ca ratio ($\Delta\text{Sr}/\text{Ca}$) following the equation: $\Delta\text{Sr}/\text{Ca} = -a \times P$, where the parameter, a , is defined by analysis of modern *T. maxima* [1]. Using the technique, we distinguished the sunny and cloudy seasons in 4.6 ka.

Reference

[1] Sano, Y., Kobayashi, S., Shirai, K., Takahata, N., Matsumoto, K., Watanabe, T., Sowa, K., Iwai, K. 2012, Past daily light cycle recorded in the strontium/calcium ratios of giant clam shells. Nature Commun. 3, DOI: 10.1038/ncomms1763

Keywords: giant clam, Sr/Ca, NanoSIMS

「汚物だめ保存仮説」: 例外的に保存されたオルステンタイプ化石の化石化のメカニズムについて

The cesspool preservation hypothesis as a key to preservation of exceptionally well preserved orsten-type fossils

田中 源吾^{1*}, 前田晴良², 松島義章³

Gengo Tanaka^{1*}, Haruyoshi Maeda², Yoshiaki Matsushima³

¹ 群馬県立自然史博物館, ² 九州大学博物館, ³ 神奈川県立生命の星・地球博物館

¹Gunma Museum of Natural History, ²Kyushu University Museum, ³Kanagawa Prefectural Museum of Natural History

最近, 神奈川県追浜公園にて掘削された完新世のコア試料のうち, ペレットを大量に含む層準より, 例外的に軟体部が保存された介形虫(節足動物)化石が発見された(Tanaka et al., 2012)。

同様な軟体部が保存された節足動物の化石は, カンブリア紀後期のオルステン石灰岩中에서도発見されている(Maeda et al., 2011)。そのような「汚物だめ」は, 集積したペレットによって作りだされた局所的な磷酸塩の濃集による例外的な磷酸塩化によるものであると考えられる。この「汚物だめ保存仮説」はカンブリア紀後期以降の海成堆積物においてみられる, 例外的に保存の良い化石の化石化の解明につながると考えられる。

キーワード: 化石, カンブリア紀, 汚物だめ, 節足動物, 古生物学, 完新世

Keywords: Fossil, Cambrian, Cesspool, Arthropoda, Paleontology, Holocene

生鉱物化と古海洋代替指標の地球化学： シンポジウムの趣旨 Biocalcification and the geochemistry of proxies

北里 洋^{1*}

Hiroshi Kitazato^{1*}

¹ 独立行政法人海洋研究開発機構、海洋・極限環境生物圏領域

¹Institute of Biogeosciences, JAMSTEC

JpGU と EGU Biogeosciences Division が共同で行っている「生鉱物化と古海洋プロキシの地球化学」の科学的背景とセッションの趣旨について述べる。

キーワード: 地球生命科学, 生鉱物, 地球化学, プロキシ, EGU-JpGU 共同セッション

Keywords: Biogeosciences, biomineralisation, geochemistry, proxies, EGU-JpGU Joint Session

Assessing the environmental impact of T?hoku tsunami off Hachinohe (NE Japan): a multidisciplinary approach.

Assessing the environmental impact of T?hoku tsunami off Hachinohe (NE Japan): a multidisciplinary approach

フォンタニエ クリストフ², 豊福 高志^{1*}
Christophe Fontanier², Takashi Toyofuku^{1*}

¹ ボルドー大学, ² 海洋研究開発機構

¹Univeristy oif Bordeaux 1, ²JAMSTEC

On March 11th 2011 the Japanese East coast was hit by a tsunami, which killed more than 18.000 people, caused major devastation in the coastal zone and the meltdown of 3 nuclear reactors. A magnitude 9 on the Richter scale earthquake offshore Sendai resulted in Tsunami waves reaching heights of up to 40.5 meters, which travelled 10 kilometers inland. Whereas the devastation on land is clearly visible, underwater impact is more difficult to assess. Here we present an overview of the multidisciplinary approach used to describe the benthic ecosystems off Hachinohe (NE Japan), 5 months after the T?hoku earthquake. Middle height (~4m) of Tsunami also came to the coastal area of Shimokita Peninsula. An oceanographic cruise (cruise KT11-20? aboard R/V TANSEI MARU, AORI/JAMSTEC) took place in August 2011. An international group of Japanese, French and Dutch oceanographers, all specialists in marine ecology and marine biogeochemistry, joined this scientific mission in order to describe benthic ecosystems and fossilizing foraminiferal faunas. 4 scientific tasks were defined. The sedimentological investigation has consisted in the identification of all sedimentary evidences (physical structures and radionuclides) that illustrate hydrosedimentary processes at the seafloor (erosion, sediment gravity flow deposition). The geochemical investigation has consisted in the optimal characterization of geochemical conditions prevailing in the benthic ecosystems. A special attention has been addressed to the dissolved species (oxygen, nitrate...) in the bottom and pore water, the organics buried in the sediment and the nature of solid phases. The faunal investigation has consisted in the ecological study of benthic foraminifera (living and dead faunas). This study has given reliable information about the response of benthic life to environmental constraints related to tsunamis. The future investigation will consist in the geochemical study of trace elements in the foraminiferal shells (i.e. tests). Those overall observations should enlighten scientific community on the effect of the T?hoku tsunami on marine ecosystems off Hachinohe, and on the potential resilience of benthic communities.

キーワード: 津波, 有孔虫

Keywords: foraminifera

Using microComputedTomography to study the impact of environmental stressors on benthic foraminifera: initial results

Using microComputedTomography to study the impact of environmental stressors on benthic foraminifera: initial results

Filipsson Helena¹, 豊福 高志^{2*}, 佐々木 理³
Helena Filipsson¹, Takashi Toyofuku^{2*}, Osamu Sasaki³

¹ ルンド大学, ² 海洋研究開発機構, ³ 東北大学総合博物館

¹Lund University, ²JAMSTEC, ³Tohoku University

We performed microComputedTomography (μ CT) scans of benthic foraminifera, in order to better understand how multiple environmental stressors are affecting biomineralization as well as preservation of benthic foraminifera. Both live (Cell Tracker Green labeled) and dead foraminiferal specimens from the Skagerrak and Kattegat, NE Atlantic were scanned. The samples originate from 330m and 130m of water depth, where salinity ranged between 35.2 (Skagerrak) and 34.7 (Kattegat) and dissolved oxygen content varied from full oxygenated in the Skagerrak to hypoxic conditions ($<2\text{mlO}_2/\text{L}$) at the Kattegat station. Substantial differences were noted in test (shell) preservation and morphology between fossil and modern samples, where pre-industrial samples were less affected by dissolution processes.

キーワード: 有孔虫

Keywords: foraminifera

Ecological impact of the Tōhoku Tsunami on shallow-water marine biodiversity off Hachinohe (NE JAPAN)

Ecological impact of the Tōhoku Tsunami on shallow-water marine biodiversity off Hachinohe (NE JAPAN)

Duros Pauline^{1*}

Pauline Duros^{1*}

¹ 海洋研究開発機構

¹JAMSTEC

The 11th of March 2011, Japan was struck by one of the most powerful known earthquakes, the so-called Tōhoku earthquake. This earthquake presented a magnitude of 9.0 and an epicenter located 70 kilometers east of the Oshika Peninsula of Tōhoku. It triggered extremely destructive tsunami waves of up to 10 meters that struck Japanese coasts. Both earthquake and tsunami caused extensive and severe structural damage in Japan. More than 15,000 people died; 8,000 are still missing. This aim of the present study is to evaluate the post-crisis environmental health of the marine biosphere from the NE Japan. In order to assess the impact of this terrible disaster on marine ecosystems, and more precisely, to assess the impact of tsunami on coastal marine ecosystems, an oceanographic cruise occurred in August 2011 and sediments off Iwate prefectures (NE JAPAN) were sampled. Living benthic foraminifera collected in these sediments were used as bio-indicators of sedimentary disturbance. Indeed, after a sediment gravity event (e.g. turbidite) triggered for instance by an earthquake, high amount of organic and inorganic detritus may be supply by lateral advection to the ocean. There, foraminiferal faunas are characterised either by recolonisation stages occurring after physical disturbance (e.g. turbidite related to tsunami) or by equilibrium phases related to gradual organic matter focussing (e.g. eutrophication). Biotic recovery after benthic crisis consists in the dominance of opportunistic pioneer species. The foraminiferal biodiversity is low. When the resilience of an ecosystem is surpassed (after weeks, months or years), opportunistic taxa are generally replaced by highly specialised communities. Then, the foraminiferal diversity increases. In case of the Tōhoku Tsunami, the inherent question is: Do benthic Foraminifera indicate environmental alteration/resilience of marine biodiversity in relation to tsunami?

キーワード: 有孔虫

Keywords: foraminifera

First-order estimate of the planktic foraminifer biomass in the modern global ocean

Movellan Aurore¹, シーベル ラルフ^{1*}, Howa Helene¹
Aurore Movellan¹, Ralf Schiebel^{1*}, Helene Howa¹

¹ アンジェ大学

¹Laboratoire des Bio-Indicateurs Actuels et Fossiles (BIAF), CNRS UMR 6112 LPGN, Universite d'Angers

Planktic foraminifers are heterotrophic mesozooplankton of global marine abundance ubiquitously used in paleoecology, paleoceanography, and paleoclimate reconstruction. However, the biomass and trophic role of planktic foraminifers was largely unknown. To better understand the position of planktic foraminifers within the regional and global plankton, we have developed a new analytical method and quantified the individual and species specific planktic foraminifer biomass. With a new non-destructive protocol developed from the bichinchoninic acid (BCA) method and nano-photospectrometry, we have analysed the foraminifer protein-biomass, along with test morphometry. From additional CHN analysis, it can be assumed that protein biomass equals carbon-biomass. The foraminifer cytoplasm is exposed to the analytical reagents without breaking the test by applying an osmotic shock. The new method is quick and easy to apply, and we have so far produced a data set of the protein-biomass in function of test size of 21 planktic foraminifer species from Atlantic, Pacific, and Southern Ocean waters.

Our data include a wide range of oligotrophic to eutrophic conditions covering six orders of magnitude of assemblage biomass. Samples include symbiont bearing and symbiont-barren species from the sea surface down to 2500 m water depth. Being secondary producers with an omnivorous diet, which ranges from algae to small metazoans, planktic foraminifers are not limited to a single food source, and are assumed to occur at a balanced abundance displaying the overall marine biological productivity at a regional scale. Accordingly, the average individual planktic foraminifer protein- and carbon biomass amounts to 0.845 μ g. Conversion factors between individual biomass and assemblage-biomass are calculated for test sizes between 72 and 845 μ m (minimum test diameter). Assemblage-biomass data presented here include 1128 sites and water depth intervals. The regional coverage of data includes the North Atlantic, Arabian Sea, Red Sea, Caribbean, as well as literature data from the eastern and western North Pacific off Japan, and covers a wide range of oligotrophic to eutrophic waters over six orders of magnitude of planktic foraminifer assemblage-biomass (PFAB). A first order estimate of the average global planktic foraminifer biomass production ($>125 \mu$ m) ranges from 8.2?32.7 Tg C yr⁻¹ (i.e. 0.008?0.033 Gt C yr⁻¹), and might be more than three times as high including neanic and juvenile individuals adding up to 25?100 Tg C yr⁻¹. However, this is a first estimate of regional PFAB extrapolated to the global scale, and future estimates based on larger data sets might considerably deviate from the one presented here. This paper is supported by, and a contribution to the Marine Ecosystem Data project (MAREDAT). Data are available from <http://www.pangaea.de> (<http://doi.pangaea.de/10.1594/PANGAEA.777386>).

キーワード: 有孔虫

Keywords: foraminifera

Vertical migration, nitrate uptake and denitrification: survival mechanisms of foraminifers under low DO conditions

Vertical migration, nitrate uptake and denitrification: survival mechanisms of foraminifers under low DO conditions

Koho Karolina^{1*}, Pina-Ochoa E², Geslin Emmanuelle³, Risgaard-Petersen N²
Karolina Koho^{1*}, E Pina-Ochoa², Emmanuelle Geslin³, N. Risgaard-Petersen²

¹Faculty of Geosciences, Utrecht University, ²Center for Geomicrobiology, Institute of Biological Sciences, Aarhus University,

³Recent and Fossil Bio-Indicators, Angers University

¹Faculty of Geosciences, Utrecht University, ²Center for Geomicrobiology, Institute of Biological Sciences, Aarhus University,

³Recent and Fossil Bio-Indicators, Angers University

¹⁵N₂O₃ - isotope labelling experiments were carried out to study foraminiferal nitrate uptake strategies and the role of pseudopodial networks in nitrate uptake. *Globobulimina turgida* were placed below the nitrate penetration depth in homogenised sediment cores, which were subsequently incubated in artificial seawater containing the label. The physical migration of foraminifera to strata containing nitrate and oxygen was prevented by a nylon net, however, potential access to such strata by extension of pseudopods was still possible. As no ¹⁵N₂O₃ - was found in *G. turgida* in the experimental cores, we concluded that foraminifera cannot extend their pseudopods for nitrate uptake through several millimetres of sediment. In stead they must physically migrate upwards closer to nitratecontaining strata. The foraminiferal migration patterns in the control cores (with no nylon net) were observed to be erratic, suggesting that individuals move in random orientations until they find favourable conditions (i.e. free nitrate or oxygen).

A second experiment showed that foraminifera actively collect nitrate in both the presence and absence of oxygen, although uptake was initiated faster if oxygen was absent from the environment. However, no systematic influence of the size of the intracellular nitrate pool on nitrate uptake was observed, as specimens containing a large range of intracellular nitrate (636-19992 pmol/cell) were measured to take up ¹⁵N₂O₃ - at comparable rates.

キーワード: 有孔虫

Keywords: foraminifera

Seawater Mg/Ca variability during the Middle Miocene Climatic Optimum Seawater Mg/Ca variability during the Middle Miocene Climatic Optimum

ド・ノイエ レナート^{1*}, van Dijk IEY¹, 豊福 高志³, Sluijs A², Reichart GJ¹
Lennart Jan de Nooijer^{1*}, IEY van Dijk¹, Takashi Toyofuku³, A Sluijs², GJ Reichart¹

¹NIOZ, ²ユトレヒト大学, ³海洋研究開発機構

¹Royal NIOZ, ²Utrecht University, ³JAMSTEC

Variability in seawater [Ca²⁺] and [Mg²⁺] over timescales >1 Ma challenges the use of foraminiferal Mg/Ca as a temperature proxy. Since temperature and seawater Mg/Ca both determine foraminiferal Mg/Ca, reconstructed temperatures need to be corrected for past seawater Mg/Ca when applied to long timescales. Currently, such corrections are based on models with a low temporal resolution and relatively large uncertainty in past seawater Mg/Ca. Moreover, when applying correction factors it is assumed that the sensitivity of the Mg/Ca-temperature calibration is not affected by seawater Mg/Ca. To quantify the combined impact of seawater Mg/Ca and temperature on foraminiferal Mg/Ca, we conducted a set of culturing experiments in which these parameters were manipulated independently. The combined effect of seawater Mg/Ca and temperature on calcite Mg/Ca in a hyaline (*Elphidium crispum*) and a miliolid (*Quinqueloculina* sp.) species was determined by laser ablation-ICP-MS.

The dependencies of calcite Mg/Ca on these two parameters for both species were used to reconstruct seawater Mg/Ca over the Middle Miocene Climatic Optimum (MMCO) from the Equatorial Pacific using IODP core 1338. Using the different Mg-incorporation mechanisms of hyaline and miliolid foraminifera reveals that seawater Mg/Ca for this interval is on average different and more variable than previous studies suggested. The accompanying deep sea temperatures for this interval are on average lower than previously reported. This new reconstruction also shows that variability in seawater Mg/Ca warrants high resolution reconstructions when correcting temperatures based on foraminiferal Mg/Ca.

キーワード: 有孔虫

Keywords: foraminifera

Seasonality in the Arabian Sea over glacial-interglacial cycles Seasonality in the Arabian Sea over glacial-interglacial cycles

ド・ノイエ レナート^{1*}, Tjallingii R¹, Brummer GJ¹, Reichart GJ¹
Lennart Jan de Nooijer^{1*}, R Tjallingii¹, GJ Brummer¹, GJ Reichart¹

¹Royal Netherlands Institute for Sea Research

¹Royal Netherlands Institute for Sea Research

The Indian monsoon system controls seasonal precipitation alterations over the Indian continent and upwelling of nutrient-rich waters to the surface in the northern Arabian Sea. Functioning and strength of this weather system due to climate change is one of the important issues in predicting the effects of global warming on the region's economy, agriculture and social welfare. The strength of the Indian monsoon system through time can be studied by changes in seawater temperature and chemistry from single-specimen analysis of planktic foraminiferal calcite. Temperature reconstructions based on many single specimens allow reconstruction of past seasonal sea water temperatures ranges and thus seasonal temperature variability.

Here we present seawater reconstructions based on single-specimen Mg/Ca of the surface dweller *Globigerinoides ruber* and the deeper-living *G. dutertrei* of two sediment cores of the western equatorial Indian Ocean off Tanzania and the northern Arabian Sea. From both cores, specimens are analyzed for calcitic Mg/Ca using laser ablation-ICP-MS of time-intervals representing the Holocene optimum, Last Glacial Maximum, Marine Isotope Stage 4, MIS 5 and MIS6. The resulting temperature ranges allow reconstruction of variability in the strength of the Indian Monsoon as well as cross-equatorial heat transport during glacial and interglacials.

キーワード: 有孔虫

Keywords: foraminifera

アカガイの微細構造形成の水温飼育実験

Temperature-controlled experiments for the shell microstructural formation of *S. broughtonii* (Mollusca: Bivalvia)

西田 梢^{1*}, 鈴木 淳², 磯野良介³, 渡邊裕介³, 野尻 幸宏⁴, 森千晴¹, 佐藤瑞穂², 佐藤圭¹, 佐々木猛智¹

Kozue Nishida^{1*}, Atsushi Suzuki², Ryosuke Isono³, Yusuke Watanabe³, Yukihiro Nojiri⁴, Chiharu Mori¹, Mizuho Sato², Kei Sato¹, Takenori Sasaki¹

¹ 東京大学大学院理学系研究科, ² 産総研, ³ 海洋生物研究所, ⁴ 国立環境研

¹The University of Tokyo, ²AIST, ³MERI, ⁴National Institute for Environmental Studies

A temperature experiment was performed to corroborate the thermal control of microstructural formation, and the cultured specimens were found to exhibit differences in shell formation by temperature. It has been suggested that the cyclical changes in the shell microstructures of *S. broughtonii* might be affected by temperature seasonality based on a study of field specimens (Nishida et al., 2012). This is the first report of temperature experiments in relation to the microstructural formation of shells. Additionally, this experiment contributes to the reconstruction of the paleoenvironments using shell microstructures and to our understanding of the mechanisms of shell microstructural formation.

We cultured specimens of *S. broughtonii* under five different temperature conditions at the Demonstration Laboratory, Marine Ecology Research Institute (MERI) in Kashiwazaki City, Niigata Prefecture, Japan. The investigated temperatures were 13 C, 17 C, 21 C, 25 C, and 29 C, and the specimens were cultured for approximately 58 days. We placed 5 aquariums (12 liters) in the laboratory with 5-7 specimens placed in each aquarium. We removed part of the marginal periostracum to determine shell growth during the experiment.

The shell sizes and increment of the shell deposition during the experiment show that the most rapid growth occurs at 17 C. Based on the $\delta^{18}O$ data, the specimens at 17 C, 21 C, 25 C, and 29 C formed shell material at each temperature condition. The thickness of the composite prismatic structure increases at higher water temperatures, and this trend is same as that of the field specimens. The specimen at 17 C showed the sharpest edge in the marginal part of the outer layer in comparison to the specimens cultivated at 21 C, 25 C, and 29 C. Accounting for the outer layer, the area of the composite prismatic structure increases as the water temperature is reduced. The growth increment of the crossed lamellar structure was relatively constant, whereas that of the composite prismatic structure increased rapidly as the thickness of the composite prismatic structure increased at cooler temperatures. This finding suggests that the optimum temperature for *S. broughtonii* growth as determined experimentally is consistent with the shell growth in the temperate area and that the formation of the composite prismatic structure increases the shell growth, especially the expansion of the growth increments in the outermost part of the outer layer.

Keywords: Bivalve, shell microstructure, Stable oxygen isotopes, temperature experiment, shell growth

東地中海クレタ島沖の海底塩湖堆積物の地球化学 (KH06-04 航海): リンの形態別存在量から探る過去5~21万年前の酸化還元状態 Phosphorus speciation analysis of sediments in the hypersaline Meedee Lake, off Crete Island, Eastern Mediterranean Sea

山口 友理恵^{1*}, 山口 耕生², 村山 雅史³, 池原 実³

Yurie Yamaguchi^{1*}, Kosei E. Yamaguchi², Masafumi MURAYAMA³, Minoru Ikehara³

¹ 東邦大学理学研究科化学専攻, ² 東邦大学理学研究科化学専攻, Nasa Astrobiology Institute, ³ 高知大学海洋コア総合研究センター

¹Department of Chemistry, Graduate School of Science, Toho University, ²Department of Chemistry, Graduate School of Science, Toho University, Nasa Astrobiology Institute, ³Center for Advanced Marine Core Research, Kochi University

現代の地中海には、約 5.6-5.3Ma の Messinian Salinity Crisis 時に堆積した海底下の岩塩層からしみ出した高塩分濃度の流体が起源となる、海底塩水湖が複数存在する。学術研究船白鳳丸の KH06-04 航海で採取された東地中海クレタ島沖の海底塩水湖 (Meedee Lake; 塩分濃度は海水の 10 倍) 堆積物は、明・暗色層が数 cm ~ 数 10cm 間隔で互層する。この色調変化は、有孔虫種の解析等により、当時の堆積環境の酸化還元の変遷に起因するとされる (泉谷, 2010)。本研究では、この酸化還元状態の変化の定量化を行うため、リンの存在形態別の定量分析を行った。海洋中では主にリン酸として存在するリンは、(微)生物活動・酸化還元状態・続性作用の影響を受けて、堆積物中ではアパタイトを形成したり、鉄酸化物に吸着したり、有機態として保存される。逆に、堆積物中のリンの存在形態別の良は、堆積環境の推定を行う手がかりとなる。Ruttenberg (1992) と Schenau and de Lange (2000) を改良した橋本 (2010) の方法に従って、KH06-04 堆積物の粉末試料中のリンの 5 種の存在形態別 ([1] 吸着性リン + 生体由来アパタイト + 炭酸塩結合態リン、[2] 鉄結合態リン、[3] 炭酸フッ素アパタイト (CFAP)、[4] 碎屑性リン (FAP)、[5] 有機態リン) の連続抽出分析を行った。リン濃度の測定はモリブデン青法により吸光度計で行った。

形態別では、CFAP 相に含まれるリン濃度が他の形態と比較し大きく、明色層中では全体の約 76%、暗色層中では全体の約 67% であった。有機態リン・碎屑性リンの濃度は予想値よりも遥かに小さかった。

明/暗色層でのリン濃度差が最も大きい形態は、鉄結合態リンであり、明色を示すほど、鉄結合態リン濃度は低くなる傾向にあった。これは、リン酸は酸化的環境では鉄酸化物の表面に吸着するが、還元的環境では鉄酸化物の還元的溶解により表面吸着していたリン酸が海水中に放出されたためと考えられる。明色層は還元的・暗色層は酸化的な堆積環境であったことが示唆されるが、これは、泉谷 (2010) の結果と整合的である。

有機物は、酸化的環境下で好氣的分解反応が進むために存在量は減少し、有機態リンも存在量が減少すると考えられるが、今回得られた結果はその逆で、酸化的環境を示唆する暗色層の有機態リンより多かった。これは、高塩分濃度水からなる Meedee Lake の湖面付近では密度差があるために、有機物等の沈降粒子が塩水湖中に沈殿しにくく、かつ暗色層に底生有孔虫が多く生息していた (泉谷, 2010) ことから、暗色層の有機態リンの起源は底生有孔虫であることが示唆される。あるいは、堆積環境の影響ではなく、気候変動による影響を受けたとも考えられる。有孔虫殻に含まれる酸素同位体の解析より明色層・暗色層はそれぞれ温暖期・寒冷期に属していたことが明らかにされている (泉谷, 2010)。寒冷期には、海水準が低下し露出した陸棚の堆積物からリンが供給され富栄養価して生物生産が高くなったので、その影響により生産された有機物が多く堆積した可能性も考えられる。

CaCO₃ 量の多い有孔虫殻が起源となり、炭酸塩結合態のリン濃度が最も高いと予想されたが、実際は CFAP 相のリン濃度が一番高かった。その要因として 2 つ考えられる。まず、地中海塩水湖におけるリンは分解速度が大きく、分解後には海水にリン酸として回帰したのち再び塩水湖中の堆積物中に戻り、続成作用によって CFAP 相へと変化した可能性が挙げられる。通常、鉄結合態リン、有機態リンが初期続成作用を受け CFAP 相へと変化するため、それらの濃度が小さかったことから調和する。もしくは、本試料の CaCO₃ 量 (%) と CFAP 相のリン濃度が類似した深度分布を示すことから、続成作用ではなく、蒸発岩由来の鉱物等を起源とする非生物性 CaCO₃ の濃度を反映していることも考えられる。

以上より、KH06-04 航海で採取された堆積物は、酸化還元状態の変遷に加え、塩水湖自体の影響も大きく受けていることが予想される。