(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

HSC04-01

会場:102A



時間:5月23日14:15-14:30

## Future Earth, Great East Japan Disaster, and IHDP-Japan Future Earth, Great East Japan Disaster, and IHDP-Japan

氷見山 幸夫 <sup>1\*</sup> Yukio Himiyama<sup>1\*</sup>

<sup>1</sup> 北海道教育大学 <sup>1</sup>Hokkaido University of Education

Global environmental research is currently undergoing major re-structuring under the framework of Future Earth, or a platform for sustainability science initiated by ICSU, ISSC (International Social Science Council), and Belmont Forum. IHDP-Japan is uniquely positioned to implement Future Earth in a broader interdisciplinary perspective incorporating the various lessons of the Great East Japan Disaster. The paper discusses the roles IHDP-Japan can, and should, play for the increment of sustainability of the world which Future Earth aims at.

キーワード: IHDP, ICSU, Future Earth, 持続可能性科学, 東日本大震災, 地球環境変化の人間的側面研究計画 Keywords: IHDP, ICSU, Future Earth, sustainability science, Great East Japan Disaster, International Human Dimensions Programme

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



会場:102A

時間:5月23日14:30-14:45

# Relationship Between Precipitation and Carbon Dioxide (CO2) Over Indian Sub-Continent Relationship Between Precipitation and Carbon Dioxide (CO2) Over Indian Sub-Continent

R.B. Singh<sup>1\*</sup>, Mansi Janmaijaya<sup>1</sup> R.B. Singh<sup>1\*</sup>, Mansi Janmaijaya<sup>1</sup>

The association between carbon dioxide and rainfall is investigated over the Indian region. The study is performed on fine scale (2 degree, 2.5 degree) grid map for the period of 9 years from 2003 to 2011. Carbon dioxide data from Atmospheric infrared Sounder (AIRS) and rainfall data from Tropical Rainfall Measuring Mission (TRMM) are used in this study. Data are mapped for both carbon dioxide and rainfall across all the months/ seasons of the study period. The concentration of carbon dioxide is found to be highly variable spatially as well as seasonally. It varies from ~3.68,10-4 (July-August, 2003) to ~4,10-4 (January-February,2009) over land and which is larger than over oceans. It is observed that in the winter months (November to February) the concentration is relatively higher, while during the monsoon season the concentration is low. It seems to be the wash out effect of rainfall over carbon dioxide. Over the oceans the Carbon Dioxide concentration show the slight seasonal changes and the range is less as compared to the land. The rainfall varies from no rainfall to 102 cm/month (July,2007), on average monthly basis.

The correlation coefficient (Rxy) between Carbon Dioxide and rainfall is calculated during all the months from 2003 to 2011, and the values have been mapped and shown using MATLAB. The correlation coefficient is highly variable during different months over the study region, with a wide range. The computed minimum value was in January (-.8859) and the maximum value in the month of August (+.8863). The computed correlation coefficients show positive values for the summer months of the sub-continent over land with rise in values of correlation from the month of April, reaching uniformly high value in August over 9 years, and then the coefficient decreases as the autumn sets in. However, over the oceans the fluctuations are not high with the change in seasons. It is discerned that the percentage increase in rainfall is not in direct proportion to the increased carbon dioxide concentration i.e. the relationship is not linear. Moreover, the increased rainfall in response to increasing carbon dioxide concentration in a region is limited which may be attributed to certain regional parameters such as topography of the area, vegetation and emissions of carbon dioxide due to various anthropogenic activities. The study would be extended in future by taking large area and other parameters into account.

 $\neq - \nabla - F$ : Carbon dioxide concentration, rainfall, correlation coefficient, TRMM, AIRS, India Keywords: Carbon dioxide concentration, rainfall, correlation coefficient, TRMM, AIRS, India

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

HSC04-03



時間:5月23日14:45-15:00

#### Impact of Liberalized Border Trade on Rural Households and Their Land Use: The Case of Laos-China Border Area Impact of Liberalized Border Trade on Rural Households and Their Land Use: The Case of Laos-China Border Area

Takeshi Sakurai<sup>1\*</sup>, CHIEN, Hsiaoping<sup>1</sup>, SAYAVONG, Somboune <sup>3</sup>, XUAN, Yi<sup>4</sup>, KANG, Yunhai<sup>4</sup> Takeshi Sakurai<sup>1\*</sup>, CHIEN, Hsiaoping<sup>1</sup>, SAYAVONG, Somboune <sup>3</sup>, XUAN, Yi<sup>4</sup>, KANG, Yunhai<sup>4</sup>

<sup>1</sup>Hitotsubashi University, <sup>2</sup>JIRCAS, <sup>3</sup>Policy Research Center, Lao PDR, <sup>4</sup>Yunnan Academy of Social Sciences <sup>1</sup>Hitotsubashi University, <sup>2</sup>JIRCAS, <sup>3</sup>Policy Research Center, Lao PDR, <sup>4</sup>Yunnan Academy of Social Sciences

In Asia trade liberalization has been rapidly going-on. At the macro-level, liberalized trade should enhance efficiency and promote economic growth. However, it is relatively unknown what kind of impacts the liberalization has on rural households and their land use in remote, border area. In order to answer these questions, this paper utilized household data collected by household survey conducted in Laos-China border area in 2010. The sample size is 120 households in Louang Namtha Province, Laos and 120 households in Xishuangbanna autonomous prefecture, Yunnan, China.

In Laos, rubber planted area of sample households is increasing since 2005. Due to the increasing rubber plantation on upland, forest and fallow land are decreasing. In the Lao study site, rice is grown in lowland, and hence there is no conflict in land use between rubber and rice. However, rice producing area is decreasing slightly since 2005 probably due to labor allocation. In Yunnan, land for staple food crops (rice and maize) is decreasing while that for rubber is increasing in recent years. Farmers now purchase rice including rice imported from Laos. Most of the paddy fields are now leased out to banana growers.

Thus, the border trader between Laos and Yunnan has impacts on agriculture and land use of both sides. In Laos, farmers export rice to Yunnan. In Yunnan, farmers grow rubber on upland and rent-out lowland to banana growers. This is a more efficient resource allocation compared with the situation before the trade liberalization, and households on both sides seem to have become better off. However, concerns are the reduction of biodiversity due to rubber plantation on Lao side and a threat to food security caused by the specialization on Yunnan side.

 $\neq - \nabla - F$ : Land use, Trade liberalization, Laos, Yunnan, Border trade, Market integration Keywords: Land use, Trade liberalization, Laos, Yunnan, Border trade, Market integration

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



HSC04-04

#### 会場:102A

#### 時間:5月23日15:00-15:15

#### 中国蘭州市における都市開発と土地利用変化における意味 Urban development and its implication on land use change of Lanzhou City, China

土居 晴洋<sup>1</sup>\*, 柴彦威<sup>2</sup> Haruhiro Doi<sup>1</sup>\*, Chai Yanwei<sup>2</sup>

<sup>1</sup> 大分大学, <sup>2</sup> 北京大学 <sup>1</sup>Oita University, <sup>2</sup>Beijing University

シルクロードの歴史的都市である蘭州市は黄河流域にあり,甘粛省の省都であるとともに,中国内陸地域の最も西に ある大都市である。蘭州市の土地利用は内陸地域の中心都市としての発展を目指す国家的政策のもとで,工業や住宅開 発を通して変化してきた。

人口は計画経済に入って増加した。三線建設という国家政策のもとで,石油化学や重工業が設立された。1978年の 改革開放政策導入以後,重化学工業都市としての性格は総合的中心都市へと変化した。1990年代以降市街地東部の低位 河岸段丘上でスプロール的都市化が進展した。また,古い市街地再開発を目的として新しい住宅開発が行われている。中 心部では商業住宅として高層建築が作られている。中心部から高速道路で1時間の距離のところに新しい都市開発が進 行している。これは蘭州新区と言い,その広さは40平方キロメートルに及ぶ。

キーワード: 地域開発, 地形条件, 内陸地域 Keywords: regional development, topographical condition, inland area

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



HSC04-05

会場:102A

時間:5月23日15:15-15:30

### 守られる森林と放置される森林周辺部 ナガラホレ国立公園(インド)の事例 Protected Forest and its periphery - a Case of Nagarahole National Park, Karnataka, India

木本 浩一<sup>1\*</sup> Koichi Kimoto<sup>1\*</sup>

<sup>1</sup> 広島女学院大学 <sup>1</sup>Hiroshima Jogakuin University

In India, we can divided the trend of forest into two periods, the period of deforestation until the 1980s and the period of well forest protection, i.e. Joint Forest Management, since the 1990s. As a result, we can see the institutional development and a lot of effort in various areas, and the rate of forest has been improved from 10% level to 20.6% (2001). As for the forest and its periphery as forest region, the former has been under strict control as a national park, and the latter has neglected and left the chaotic region consisting of miscellaneous residents. In this presentation, I would like to investigate the possibility of governance in the region spread in a donut shape, with understanding the both region as a forest "region".

キーワード: インド, 森林, 森林地域, カルナータカ Keywords: India, Forest, Forest region, Karnataka

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

HSC04-06



時間:5月23日15:30-15:45

### Recent Flood Disaster in Jabodetabek Region: A question toward resilience Megapolitan Recent Flood Disaster in Jabodetabek Region: A question toward resilience Megapolitan

Ernan Rustiadi<sup>1\*</sup>, Andrea Emma Pravitasari<sup>3</sup>, La Ode Syamsul Iman<sup>1</sup>, Tutuk Lufitayanti<sup>2</sup> Ernan Rustiadi<sup>1\*</sup>, Andrea Emma Pravitasari<sup>3</sup>, La Ode Syamsul Iman<sup>1</sup>, Tutuk Lufitayanti<sup>2</sup>

<sup>1</sup>Center for Regional Systems Analysis, Planning and Development, Bogor Agricultural University, <sup>2</sup>Faculty of Agriculture, Bogor Agricultural University, <sup>3</sup>Laboratory of Regional Planning, Graduate School of Global Environmental Studies, Kyoto University

<sup>1</sup>Center for Regional Systems Analysis, Planning and Development, Bogor Agricultural University, <sup>2</sup>Faculty of Agriculture, Bogor Agricultural University, <sup>3</sup>Laboratory of Regional Planning, Graduate School of Global Environmental Studies, Kyoto University

Various environmental deterioration in Indonesia has increased the potential for anthropogenic disaster hazards in different regions. Floods in urbanized area such Jakarta City has impact a huge economic loss. Jakarta City and its surroundings (Megapolitan Jabodetabek) is the nation's largest urban agglomeration where more than 11 percent of the population and more than 26% of the national GDP is located. Flood in the city of Jakarta is an annual phenomenon with repeated each year and mainly occurred at the peak rainy season (between the months of December to January). Current flood disaster in Jakarta (January and February 2013) is considered a 5-year flood cycle. Various indications show that an increase in flood potential flood hazard in the region. This Increasing on the frequency flooding hazard raises the question whether the increased frequency of flooding is a result of climate change or by the deterioration of the local environmental conditions that cause the decrease in regional resiliency. This study examines the facts of climate change locally (especially changes in precipitation) as well as a variety of local environmental changes especially due to land use/cover changes that impact of the environmental carrying capacity for over the last 20 years.

 $\neq - \nabla - F$ : Flood disaster, Jabodetabek, recilience megapolitan, climate change, land use/cover change Keywords: Flood disaster, Jabodetabek, recilience megapolitan, climate change, land use/cover change

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



HSC04-P01 会場:コンベンションホール

#### Relationship between Farm Income with Migration Rate of Rural Communities (Case study: Ayah Village, Kebumen Regency) Relationship between Farm Income with Migration Rate of Rural Communities (Case study: Ayah Village, Kebumen Regency)

Agam Rafsanjani<sup>1\*</sup>, Muhammad Iqnaul Haq<sup>2</sup>, Sudrajat<sup>3</sup> Agam Rafsanjani<sup>1\*</sup>, Muhammad Iqnaul Haq<sup>2</sup>, Sudrajat<sup>3</sup>

<sup>1</sup>Undergraduate Student, Faculty of Geography, Universitas Gadjah Mada, <sup>2</sup>Undergraduate Student, Faculty of Geography, Universitas Gadjah Mada, <sup>3</sup>Academic Staff, Faculty of Geography, Universitas Gadjah Mada

<sup>1</sup>Undergraduate Student, Faculty of Geography, Universitas Gadjah Mada, <sup>2</sup>Undergraduate Student, Faculty of Geography, Universitas Gadjah Mada, <sup>3</sup>Academic Staff, Faculty of Geography, Universitas Gadjah Mada

Agriculture is the dominant sector in Kebumen. 64% of the population depends on agriculture. However, since the monetary crisis in 1998 weakened this sector. Decline in the agricultural sector is spurring people to look for other alternatives to meet their basic needs. One alternative is to migrate to other areas in search of better jobs. Research on this issue is very important because rural migration greatly affects the economy of the village.

The purpose of this study is (1) Determine the amount of net income of farm households in Ayah Village (2) Knowing the rate of labor migration in each household in Ayah Village (3) Knowing the relationship between farm income with labor migration rate in each household of Ayah Village. This research is experimental research with survey method. Data collected in the form of primary and secondary data. The primary data obtained through structured interviews. respondents were selected using purposive sampling method. Secondary data needed such as agricultural yield data from farmers' groups and data on the number of migrants from the government and local NGOs.

The results showed there is an association between the agricultural income on migration rate of rural communities. The lower income makes the higher migration.

 $\neq - \neg - arkappa$ : Farm Income, Rural Migration, Cost and Benefit of Farming, Migration Rate, Rural Migration Keywords: Farm Income, Rural Migration, Cost and Benefit of Farming, Migration Rate, Rural Migration

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



HSC04-P02

会場:コンベンションホール

Geoscience Union

# 天然資源管理における保険制度の潜在的役割-ザンビアにおける天候インデックス保険の事例-The Potential Role of Formal Insurance in Natural Resources Management: Evidence from Weather Index Insurance in Zambia

三浦 憲<sup>1\*</sup>, 櫻井 武司<sup>2</sup> Ken Miura<sup>1\*</sup>, Takeshi Sakurai<sup>2</sup>

#### 1一橋大学大学院経済学研究科,2一橋大学経済研究所

<sup>1</sup>Graduate School of Economics, Hitotsubashi University, <sup>2</sup>Institute of Economic Research, Hitotsubashi University

Poor people in rural areas of developing countries depend on natural/environmental resources for their livelihood. Particularly, once any negative shock happens, their dependence on such resources increases to cope with it. Such behavior creates concerns about the degradation and exhaustion of such resources. The concerns may be mitigated, if formal insurance works well. This is because farmers with an insurance payout will reduce their use of natural resources as an ex post coping strategy in the aftermath of a shock. In order to explore this possibility, we investigate demand for formal weather index insurance in a rural area of Zambia, where people depend heavily on natural resources, and there are no formal financial institutions.

Index insurance, basing insurance payouts on an officially observable index highly correlated with crop yield, has been expected a promising way to insulate the vulnerable against weather shocks. However, previous studies have reported lower take-up rates than expected. Although they also have pointed out impediments to insurance take-ups such as liquidity constraints, further investigation is required to understand them fully. One of potential barriers, which have not been tackled well, is ex ante self-insurance mechanisms employed by farmers for precautionary purpose.

To examine the relationship between the mechanism and insurance demand, this paper utilizes data for two years from a pilot scheme selling index insurance based on rainfall amounts in rural Zambia. Most of the farmers purchased too small amounts of the rainfall index insurance contract to completely shield them from weather risk, although the take-up rates are extremely high (more than 90% in both years). First, this paper presents descriptive evidence on the determinants of rural farmers' demand for weather index insurance. We show that impediments to insurance purchases in the scheme are (1) risk aversion, (2) limited trust in insurance provider, and (3) poor understanding of the insurance contract, all of which are consistent with findings from previous literature.

Then, we discuss the relationship with small livestock saving, one of farmers' essential self-insurance mechanisms in the study site. Even after controlling wealth levels, we find a concave relationship among them, suggesting that farmers in the site have strong demand for further insulating them against weather risks. Given the empirical result, we speculate that only traditional self-insurance mechanisms are not enough to reach the optimal level of precautionary savings. Overall, this paper provides suggestive evidence on the potential role of weather index insurance, allowing farmers to cope with weather shocks without an ex post excess dependence on natural resources. Future research will offer direct evidence on the causal impact of formal insurance provision on farmers' use of natural resources after negative shocks.

キーワード: 天然資源管理, 天候リスク, 天候インデックス保険, サブサハラ・アフリカ Keywords: natural resources management, weather risk, weather index insurance, Sub-Saharan Africa

# Japan Geoscience Union Meeting 2013 (May 19-24 2013 at Makuhari, Chiba, Japan)

(May 19-24 2013 at Makuhari, Chiba, Japan) ©2013. Japan Geoscience Union. All Rights Reserved. Japan Geoscience Union

HSC04-P02

会場:コンベンションホール

時間:5月23日18:15-19:30

