

Lithostratigraphy and radiolarian age of Late Cretaceous pelagic sedimentary rocks within the Oman Ophiolite

Toshiyuki Kurihara^{1*}, Kousuke Hara¹

¹Graduate School of Science and Technology, Niigata University

Radiolarites and micritic limestone within the northern Oman Ophiolite contain essentially continuous pelagic sedimentary records of the central Tethyan Ocean from Cenomanian to Coniacian or Santonian (Tippit et al., 1981; Hara and Kurihara, 2012). We are now reinvestigating the lithostratigraphy and radiolarian biostratigraphy of Late Cretaceous pelagic sedimentary rocks distributed in the Wadi Jizzi and surrounding areas, 40 km west of Sohar, northern Oman. We here present lithofacies of the pelagic sediments and radiolarian faunal change from late Cenomanian to Turonian.

Pelagic sediments associated with metalliferous sediments (umber) in the ophiolite are overlying basaltic extrusive rocks. Based on the geochemical feature, the basaltic rocks have been subdivided into three volcanic units: the V1 lava with the N-MORB signature, the V2 lava formed by intra-oceanic volcanism, and the V3 lava generated by intra-plate seamount magmatism (Ernewein et al., 1998). We investigated the stratigraphic distribution of radiolarians in pelagic sedimentary sequences overlying the V1 lava at Suhaylah section. The sequence of the analyzed section consists of lower metalliferous and siliceous rocks (umber and red mudstone with chert intercalations) and upper micritic limestone. In these lithologies, numerous radiolarian tests are loosely packed within the matrix, and well-preserved specimens can be extracted by a combined HF-HCl etching technique. In this section, a profound faunal change was detected; *Thanarla pulchra* and *Guttacapsa biacuta* were recovered from lower chert, indicating late Cenomanian in age, and *Rhopalosyringium scissum* and *Dictyomitra formosa* occur abundantly in upper chert and micrite. *R. scissum* first appeared near the base of Turonian (Musavu-Moussavou et al., 2007). Radiolarian occurrences indicate that the Cenomanian/Turonian boundary is present in the lower siliceous sequence. At Lasail and South Zabyan sections, pelagic sedimentary sequences overlying the V2 lava are well developed. The sequence at Lasail section consists of highly altered metalliferous sediments, red mudstone with chert intercalations, and micritic limestone, in stratigraphic ascending order. Hara and Kurihara (this volume) presents the result of detailed lithologic observation and radiolarian biostratigraphy of this section, and reports the occurrence of *Dictyomitra koslovae*, indicating late Coniacian to Santonian. In the Hilti area, we also have observed several good sections of chert and micrite overlying the V2 lava (Kurihara and Hara, 2012). A preliminary biostratigraphic result of this area has been presented by Kurihara and Hara (2012), and additional detailed lithostratigraphic descriptions are presented by Hara et al. (this volume).

At this time, we have obtained late Cenomanian to Coniacian (or Santonian) radiolarians from several sections of different tectonic settings deduced from the volcanostratigraphy and geochemistry of basaltic extrusive rocks. Time scale given by the radiolarian biostratigraphy shows potential usefulness to give age constraints for change in the tectonic setting of basaltic rocks. In addition, faunal transition of radiolarians will provide information on marine environmental changes in the central Tethys during Late Cretaceous.

Upper Cretaceous radiolarian biostratigraphy of the Suhaylah Formation in the Wadi Jizzi area of the Oman Ophiolite

Kousuke Hara^{1*}, Toshiyuki Kurihara¹

¹Graduate School of Science and Technology, Niigata University

The Oman Ophiolite, a member of the Tethyan ophiolites, crops out over 600 km long and up to 150 km wide in the Oman Mountains. The ophiolite suite consists of mantle peridotites, gabbros, a sheeted dyke complex, and extrusive lavas overlain by pelagic sediments. The pelagic sediments, called the Suhaylah Formation, have been studied by Fleet and Robertson (1980) in terms of depositional environments. Tippit et al. (1981) reported Late Cretaceous radiolarians from the Suhaylah Formation and estimated their age as early Cenomanian to Santonian; however detailed stratigraphic distributions of radiolarians were not shown in their study. In this study, the lithostratigraphy and radiolarian biostratigraphy of pelagic sediments overlying extrusive lavas of the Oman Ophiolite are reinvestigated in order to understand the history of the pelagic sedimentation and the faunal transition process and evolution of radiolarians during Late Cretaceous.

Radiolarian study has been conducted at Lasail section, about 35 km west of Sohar. The pelagic sediments of this section are subdivided into three lithologies: hydrothermal sediments (5 m), red shale with chert intercalations (13 m), and micritic limestone (18 m). Basal lava could not be observed at this section. The hydrothermal sediments are highly altered and intercalated with dark red to orange chert. The red shale is dark red purple in color in the lower part and dark red in the upper part. Several chert layers are interbedded with the red shale. The micritic limestone is muddy and red purple in color in the lower part. The upper part consists of red purple and greenish gray micritic limestone with fine lamination.

We obtained moderately preserved radiolarians by a combined HF-HCL etching technique. We recovered poorly preserved *Pseudodictyomitra pseudomacrocephala* (Squinabol) from the hydrothermal chert of the lowermost part. *Rhopalosyringium scissum* O' Dogherty was obtained from chert in the lower red shale. Based on these occurrences, the hydrothermal chert in the lowermost part and the lower red shale are Turonian in age. In addition, *Dictyomitra koslovae* Foreman is recovered from muddy micritic limestone in the lower part of micritic limestone. The range of *D. koslovae* is late Coniacian to Maastrichtian based on Hollis and Kimura (2001). Thus, age of the lower part of micritic limestone is probably younger than late Coniacian.

Keywords: Oman Ophiolite, Suhaylah Formation, radiolarians, Upper Cretaceous

Sulfur isotope profiles in the pelagic Panthalassic deep sea during the Permian-Triassic transition

Satoshi Takahashi^{1*}, Kunio Kaiho², Rie Hori³, Paul Gorjan⁴, Takahiro Watanabe², Satoshi Yamakita⁵, Yoshiaki Aita⁶, Atsushi Takemura⁷, Bernhard K. Sporli⁸, Takeshi Kakegawa², Masahiro Oba²

¹Department of Earth and Planetary Science, University of Tokyo, ²Graduate school of Science, Tohoku University, ³Graduate school of Science and Engineering, Ehime University, ⁴Department of Earth and Planetary Science, Washington University, ⁵Faculty of Education and Culture, University of Miyazaki, ⁶Faculty of Agriculture, Utsunomiya University, ⁷Geoscience Institute, Hyogo University of Teacher Education, ⁸Geology, School of Environment, The University of Auckland

Mesozoic accretionary complexes in Japan and New Zealand contain Panthalassic low latitude and southern mid-latitude deep-water sedimentary rock respectively. These sedimentary rocks record environmental changes in the pelagic Panthalassic Ocean during the transition associated with the severe Permian-Triassic mass extinction. This study presents sulfur isotope records of sulfide from continuous deep-sea Permian-Triassic boundary sections located in northeast Japan (the Akkamori section-2, the most continuous section amongst other previously reported deep-sea sections) and North Island of New Zealand (the Waiheke-1 section, providing the first sulfur isotopic record from a southern hemisphere deep-sea section). Both sections show sharp minus 15 permil drops of the sulfur isotope ratio coupled with a negative shift of organic carbon isotope ratio. Similar decreases in sulfur isotope ratio of carbonate-associated sulfates by minus 10 permil accompanied with a negative shift of inorganic carbon isotope ratio at the end-Permian mass extinction horizon have been reported in some shallow water Paleotethyan sections. These sulfur isotope changes suggest that a massive release of ³²S-enriched sulfur from the H₂S-rich water to the oxic surface-waters coincided with the end-Permian mass extinction.

Keywords: Mass extinction, Panthalassa, Pelagic deep sea, Permian, Triassic, Sulfur Isotope

Towards an integrated Triassic magneto-biostratigraphic time scale for the pelagic Panthalassa Ocean

Tetsuji Onoue^{1*}, Hikaru Kato², Koji Uno³, Noritoshi Suzuki², Honami Sato⁴, Satoshi Takahashi⁵, Satoshi Yamakita⁶

¹Kagoshima University, ²Tohoku University, ³Okayama University, ⁴Kyushu University, ⁵University of Tokyo, ⁶Miyazaki University

The aim of this study is to contribute to the development of an integrated geologic time scale for the Middle and Upper Triassic Panthalassic sites. The chronology for the Triassic pelagic deposits in the Panthalassa Ocean is based on the radiolarian zonation, which is well established in the Middle and Upper Triassic bedded chert successions in the Japanese accretionary complex. Although accurate calibration for the chronostratigraphic stages and substages are established basically by means of ammonites and conodonts, most of the Japanese radiolarian zones were calibrated through correlation with zonal schemes established in other regions, and have not been calibrated with magnetostratigraphy and conodont biostratigraphy. Here we present the results of magneto- and biostratigraphic study of two Triassic sections in Japan: (i) Middle Triassic Ajiro section in the Tsukumi area, Chichibu Terrane and (ii) Upper Triassic Sakahogi section in the Inuyama area, Mino Terrane. The study sections are correlated with the Triassic geomagnetic polarity timescale by means of radiolarian and conodont biostratigraphy.

Keywords: Triassic, radiolaria, conodont, magnetostratigraphy, chert

Quantitative Evaluation of Morphology for the Planktic Foraminifer *Globigerinoides ruber*

Naoko Kishimoto^{1*}, Katsunori Kimoto²

¹Setsunan University, ²JAMSTEC

Recently we can obtain three dimensional details of skeleton of foraminifer and radiolaria using micro X-ray CT. *Globigerinoides ruber* is an abundant planktic foraminiferal species often used in reconstruction of sea surface conditions in the global ocean. Within the *G.ruber*, two variations in shell color are recognized, *G.ruber* "white" and *G.ruber* "pink" with pink colored chambers. Results from molecular phylogenetic analysis supported the separate treatment of the two chromotypes. In this study, we present results from morphometric measurements on three dimensional information of *G. ruber* white and pink using micro X-ray CT.

Creating exact 3D models for scientific research

Yoshinori Teshima^{1*}

¹Chiba Institute of Technology

We involved with 'Research on the recognition of 3D objects by visually handicapped persons and development of 3D geometrical teaching materials' project in Japan (FY2006 - FY2009). One of our main aims is to develop teaching materials to enrich the world of observation of blind people by touch. We have been creating models that blind people have never touched. The models have innumerable topics, but can be divided into two categories. The first includes models with objects that we cannot actually touch because of their size-gigantic or microscopic objects. As concrete examples, we are developing models of the planets in the solar system (Earth, Mars, Venus, and Moon). Enlarged models of microorganisms like radiolarian and foraminifera are also examples of such models. The second category consists of models with abstract objects. We created models related to mathematics. Models include ring torus, horn torus, spindle torus, hula-hoop surface, Bohemian dome, Klein bottle, regular polyhedra, semi-regular polyhedra and crystallographic structures. Touchable models are significant for not only visually impaired people but also sighted people. The models are useful for systematic and intuitive learning of mathematics and sciences.

Keywords: 3D models

Os isotope stratigraphy of ferromanganese crust in the Pacific Ocean and Philippine Sea

Tatsuo Nozaki^{1*}, Kosuke Goto², Ayaka Tokumaru³, Yutaro Takaya², Katsuhiko Suzuki¹, Qing Chang¹, Jun-Ichi Kimura¹, Yasuhiro Kato⁴, Gen Shimoda², Akira Usui⁵, Tetsuro Urabe³

¹JAMSTEC/IFREE, ²AIST/GSJ, ³Univ. of Tokyo, ⁴Univ. of Tokyo, ⁵Kochi Univ.

We report the Os isotope stratigraphy of Fe-Mn crusts collected from the Takuyo #5 and MC10 Seamounts in northwestern Pacific Ocean and Ryusei Seamount in the Philippine Sea. Based on the depositional age of Fe-Mn crusts dated by Os isotopes together with major and trace element compositions determined by ICP-QMS analyses, we will discuss growth rate, geochemical signature and genesis of Fe-Mn crusts.

Keywords: ferromanganese crust, geochemistry, Re-Os isotope, Pacific Ocean, Philippine Sea

Geology and radiolarian ages of the Kamiaso unit in the Mino-Seki area, Gifu Prefecture, central Japan

Yusuke Kitagawa^{1*}, Atsushi Matsuoka²

¹Graduate School of Science and Technology, Environmental Science and Technology, Earth Science, Niiga, ²Department of Geology, Faculty of Science, Niigata University

The Mino terrane, one of the disrupted terranes in southwest Japan, is divided into several tectonostratigraphic units on the basis of composition, fabric and age. However, there is a problem that these data are biased, because detailed studies have been conducted only in limited areas. The Mino-Seki area of the central part in Gifu Prefecture is one of such areas. According to Wakita (1988b), this area is occupied by the Kamiaso unit that is characterized by repeating coherent chert-clastic sequences.

As a result of a detailed field work, accretionary complexes in the Mino-Seki area are divided into an upper or coherent unit and lower or melange unit, based on differences in lithology and geological structure. The upper unit is characterized by a tectonic pile composed of chert-clastic sequences that retain the oceanic plate stratigraphy. Middle to Late Triassic radiolarians were obtained from chert. The lower unit includes conglomerate, melange and alternating beds of chert and limestone unlike the upper unit. There are also differences in lithology of the chert. Black chert with weathered red surface is widely distributed along the Nagara River. These lithofacies generally are not recognized the Kamiaso unit. Chert samples yield Middle Triassic to Early Jurassic radiolarians, while siliceous mudstone samples yield Middle Jurassic radiolarians. A chert sample in alternating beds of chert and limestone yields of Late Triassic radiolarian. Igo and Koike (1975) reported late Norian conodonts from a limestone sample in alternating beds of chert and limestone.

The lower unit may be correlated to a tectonostratigraphic unit other than the Kamiaso Unit on the basis of composition, fabric and radiolarian ages.

Keywords: Mino terrane, Kamiaso unit, Mino-Seki area, chert, siliceous mudstone, radiolaria

Benthic foraminifers on the observational buoy: Evidence of the meroplankton stage in their life

Katsunori Kimoto^{1*}, Shiro Hasegawa², Hiroshi Namikawa³, Minoru Kitamura¹, Hajime Kawakami¹, Makio Honda¹

¹Japan Agency for Marine-Earth Science and Technology, ²Kumamoto University, ³National Museum of Nature and Science

Colonization of new habitat of benthic foraminifers is related to their diversion, survival strategies and evolutions. However their dispersal mechanisms are not well documented and still poorly understood. Here we report a new evidence of floating benthic foraminifers communities in the open ocean. They had lived on the stems of hydrozoan attaching to observational moorings in the Pacific Ocean.

Physical and biogeochemical observational mooring systems were deployed at the Station S1 (30N, 145E, water depth: 5,900m). Moored periods were from August 2011 to July 2012. Hydrozoan attaching on the mooring systems were observed on the surface of sediment trap and float at shallower depths (~320 m) and we could not observed hydrozoan at the 550 m water sediment trap. More than 300 individuals of benthic foraminifers attached of the surface of hydrozoan body. Fourteen living benthic foraminifers were identified under the microscope. Some of them were sessile and shallow water species. It is noteworthy that some agglutinated specimens were also identified: They made their shells by using other calcareous plankton (i.e. planktic foraminifers, coccolithophores, and calcareous dinoflagellata). Although it is known that only 5 species has meroplanktic (temporary planktic) stage in their life, there are no reports for above benthic foraminifers since now. Our finding suggests that benthic foraminifers which has meroplanktic life stage exist much further in the nature.

Keywords: Benthic foraminifera, agglutinated foraminifera, meroplankton, Hydrozoan, Pacific Ocean

Mesozoic radiolarian biostratigraphy in pelagic sediments in the Kermanshah area, west Iran

Atsushi Matsuoka^{1*}, Seyed Hamid Vaziri²

¹Niigata University, ²Islamic Azad University

The ophiolite belts in Iran are important regions to elucidate paleoenvironmental reconstruction of the entire Neo-Tethys. Pelagic sequences associated with ophiolitic rocks are well exposed in the Kermanshah area, west Iran. In our field survey in 2011, we worked at several localities of the pelagic sequences. Radiolarian analysis revealed that the pelagic sequences are categorized into two groups: Upper Triassic sequence represented by the Gohareh section and Middle Jurassic-Lower Cretaceous sequence represented by the Bisetun section.

The Gohareh section is composed mainly of red bedded chert with alternating beds of chert and limestone. Some micritic limestone beds contain nodular cherts. Several samples of red chert yield moderately preserved Late Triassic radiolarians including *Tritortis(?)* sp. and *Capnucosphaera* sp. The Bisetun section consists of red and green chert. Limestone-dominated intervals are also recognized in the section. Middle Jurassic to Early Cretaceous radiolarians were obtained from red and green chert samples. Identified radiolarian zones include the *Striatojaponocapsa conexa* Zone (middle Bathonian-late Callovian), *Kilinora spiralis* Zone (Oxfordian), *Hsuum maxwelli* Zone (Kimmeridgian), and *Pseudodictyomitra carpatica* Zone (Tithonian-early Valanginian).

Gharib and De Wever (2010) reported Mesozoic radiolarians ranging in age from early Pliensbachian to Turonian in the Kermanshah area for the first time. Our research adds the occurrences of Late Triassic radiolarians from pelagic sequences together with Middle Jurassic-Early Cretaceous radiolarians. The pelagic sequences in the study area were accumulated at different depositional sites of the Neo-Tethys. Our research clarified that the depositional history of the ophiolitic belts in west Iran, part of the Neo-Tethys, can be traceable to the Late Triassic.

Keywords: Radiolaria, siliceous sediment, pelagic, Mesozoic, Kermanshah, Iran

Lithofacies and Late Cretaceous radiolarians of pelagic sediments on the V2 lava of the Oman Ophiolite

Kousuke Hara^{1*}, Yumi Agui², Rina Hayashi², Tatsuya Hinohara¹, Toshiyuki Kurihara¹

¹Graduate School of Science and Technology, Niigata University, ²Department of Geology, Faculty of Science, Niigata University

The Oman Ophiolite consists of mantle peridotites, gabbros, a sheeted dyke complex, and extrusive lavas overlain by pelagic sediments. The basaltic rocks have been subdivided into three volcanic units: the V1 lava with the N-MORB signature, the V2 lava formed by intra-oceanic volcanism, and the V3 lava generated by intra-plate seamount magmatism (Ernewein et al., 1998). Pelagic sediments commonly occur at the boundaries between these volcanic units and consist of metalliferous sediments, red shale, chert, and micritic limestone. The V2 and V3 lavas are widely distributed in the Wadi Hilti area, about 25 km west of Sohar, northern Oman Mountains. Very recently, the emplacement mechanism of the V3 lava was studied by Umino (2012). Pelagic sediments, about 50 m thick at a maximum, overlie V2 effusive rocks and are covered by the V3 lava. The sediments also occur within the V3 lava. We observed lithostratigraphies of pelagic sediments for the following sections. 073 section: This section is composed of lower dark red purple metalliferous sediments and upper red shale intercalated with micritic limestone. The total thickness of these sediments is 12 m. 254 section: This section consists of red siliceous shale (3.5 m), black siliceous shale (1 m), and brown chert (0.5 m), in stratigraphic ascending order. 029 section: we measured light gray and red micritic limestone of ca. 7 m thick. The following species have been obtained from red micrite of 029 section (Kurihara and Hara 2012): *Dictyomitra formosa* Squinabol, *Pseudotheocampe urna* (Foreman), and *Rhopalosyringium scissum* O'Dogherty. Based on O'Dogherty (1994) and Musavu-Moussavou et al. (2007), *R. scissum* first appeared near the base of Turonian. Thus, these radiolarians are assignable to early Turonian or slightly younger age. Future detailed work for radiolarian biostratigraphy of these pelagic sediments will provide information about the age constraints for igneous activities of the V2 and V3 lavas.