

ISS 暴露部搭載 極端紫外光撮像装置 (EUVI) の初期結果 Initial Results from the Extreme Ultraviolet Imager (EUVI) of ISS-IMAP

伊佐敷 一裕^{1*}, 吉川 一郎¹, 宇治 賢太郎¹, 吉岡 和夫², 村上 豪², 山崎 敦², 穂積 裕太³, 齊藤 昭則³
Kazuhiro Isashiki^{1*}, Ichiro Yoshikawa¹, Kentaro Uji¹, Kazuo Yoshioka², Go Murakami², Atsushi Yamazaki², Yuta Hozumi³,
Akinori Saito³

¹ 東大, ² 宇宙研, ³ 京大

¹Univ. of Tokyo, ²ISAS/JAXA, ³Kyoto univ.

The extreme ultraviolet imager (EUVI) on the international space station (ISS) was launched in July 2012 and started observations in December 2012. The EUVI detects resonantly scattered EUV emissions from He⁺ (30.4 nm) and O⁺ (83.4 nm). It directs toward the Earth's limb and provides images of He⁺ and O⁺ distribution in the ionosphere and plasmasphere on the night side with 15 degrees of field of view. The maximum spatial resolution is 0.1 degree and the exposure time is 1 minute. In this presentation, we show images taken by EUVI at various latitudes, longitudes and local times. We will discuss temporal and spatial variations of He⁺ and O⁺ in the ionosphere and plasmasphere.

キーワード: 電離圏, プラズマ圏, 国際宇宙ステーション, 極端紫外光
Keywords: ionosphere, plasmasphere, ISS, EUV

ISS-IMAP 搭載極端紫外光撮像装置 (EUVI) の地上較正試験 Performance of the Extreme Ultraviolet Imager (EUVI) of ISS-IMAP mission

宇治 賢太郎^{1*}, 吉川 一郎¹, 伊佐敷 一裕¹, 吉岡 和夫², 村上 豪², 山崎 敦², 齊藤 昭則³, 穂積 裕太³
Kentaro Uji^{1*}, Ichiro Yoshikawa¹, Kazuhiro Isashiki¹, Kazuo Yoshioka², Go Murakami², Atsushi Yamazaki², Akinori Saito³,
Yuta Hozumi³

¹ 東大, ² 宇宙研, ³ 京大

¹Univ. of Tokyo, ²ISAS/JAXA, ³Kyoto Univ.

The Extreme Ultraviolet Imager (EUVI), which is the part of ISS-IMAP (Ionosphere, Mesosphere, upper Atmosphere and Plasmasphere mapping) mission, was successfully launched in July 2012 and began its on-orbit operations. EUVI has two prime-focus telescopes that are designed to detect the resonantly scattered emissions from He⁺ (at the wavelength of 30.4 nm) and O⁺ (83.4 nm) respectively. In order to clarify the plasma transport process in Earth's upper atmosphere, EUVI observes distributions of He⁺ and O⁺ in the ionosphere and plasmasphere.

The optical instruments consist of multilayer coated mirrors, metallic thin filters and microchannel plate detectors. The EUVI ground calibration has been carried out and the total sensitivities are estimated to be 0.060 cps/R for He⁺ and 0.0023 cps/R for O⁺.

In this poster, the instrumental design and performance of EUVI are presented.

キーワード: 国際宇宙ステーション, 極端紫外光, 電離圏, プラズマ圏

Keywords: ISS, EUV, ionosphere, plasmasphere

GPS PWV Estimation during Dust Storm Season GPS PWV Estimation during Dust Storm Season

DongSeob Song^{1*}

DongSeob Song^{1*}

¹Dept. of Ocean Construction Engineering, The Kangwon National University

¹Dept. of Ocean Construction Engineering, The Kangwon National University

Dust storms originate from the arid area are of concern to microwave propagation and satellite telemetry due to they can be influenced to carrier depolarization and phase delay. As technology advances, and new methods of making measurements are developed or made more economical, it becomes feasible to make comparative measurements of the same parameters using independent techniques. One such technique is the ground-based GPS meteorology, which is able to quickly and inexpensively expand the number of global upper-air moisture observations for radiosonde and satellite verification. On 31 March-02 April 2007, heavy Asian dust storm hit the Korean peninsula in East Asia causing extensive damages. In this paper, the Korean GPS permanent stations were used to estimate precipitable water vapor (PWV) during dust storm season. The GPS PWV estimates have monitored the density variations of dust storm associated with anomalous atmospheric wet/dry conditions. In order to monitor the GPS PWV variations to establish the relationship with the level of atmospheric concentration of aerosol particulates during an Asian dust storm, GPS PWV estimations were analyzed during March 30-April 04, 2007. In this period, the maximum value of PM10 (2,019 microgram/cubicmeter) was recorded in Daegu station at 07hr April 01 2007 (UTC) and at Seoul is 1,233 microgram/cubicmeter (12hr April 01) and at Cheonan is 1,261 microgram/cubicmeter (15hr April 01). It can be found that the actual GPS PWV variations are correlated with the spatio-temporal variability of the PM10 data.

Acknowledgement

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012-0007973).

キーワード: GPS PWV, Dust Storm, PM10

Keywords: GPS PWV, Dust Storm, PM10