

## LLR simulation study for future observations LLR simulation study for future observations

野田 寛大<sup>1\*</sup>, Dale H. Boggs<sup>2</sup>, James G. Williams<sup>2</sup>  
Hiroto Noda<sup>1\*</sup>, Dale H. Boggs<sup>2</sup>, James G. Williams<sup>2</sup>

<sup>1</sup>National Astronomical Observatory of Japan, <sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

<sup>1</sup>National Astronomical Observatory of Japan, <sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

**Introduction:** Lunar Laser Ranging (LLR) measures the distance between laser link stations on the Earth and retroreflectors on the Moon by detecting the time of flight of photons from a high-powered laser pulse emitted from the ground stations. Since the Earth-Moon distance contains information on lunar orbit, lunar solid-body tides, and lunar orientation and rotation, we can estimate the inner structure of the Moon by constraining relevant physical parameters. Several lunar landing missions which will carry new retroreflectors to the lunar surface are under study in several countries. Furthermore, retroreflectors with a larger single aperture are under development for more precise ranging, apart from the conventional array-type retroreflectors that were realized in the Apollo and Luna mission from 1969-1973. It is not obvious how lunar physical parameters such as Love numbers will be better constrained by using range data with higher accuracy. Therefore we have conducted a simulation study of the LLR observations by using the LLR analysis software of JPL.

**Method:** Simulated data were created by adding noise to the predicted distances between the Earth surface and retroreflectors on the Moon using the lunar ephemeris and range model. The simulated data were fit using a least-square solution, and then the uncertainties of the fit were evaluated. There are high degrees of freedom for the creation of simulated data in terms of the number and locations of retroreflectors, the amount of data from each retroreflector, etc. In this study we set the condition as follows:

- the number of yearly range data was set to be about 600, monthly about 50
- the numbers of range data toward the new retroreflectors are at the same level as for the Apollo 11 and 14 sites
- range accuracy for new retroreflectors was set one order-of-magnitude higher than that of the existing retroreflectors
- locations of the new retroreflectors are near the north pole, near the south pole, and the mid-latitude.

**Results and remarks:** After about 35 years of observation, the uncertainty of some parameters is reduced by about 1/2 by adding one high-accuracy retroreflector compared to the case in which only existing retroreflectors are ranged. It is suggested that the uncertainty could be reduced by observing high-accuracy retroreflectors more than existing ones, or changing the weighting of data.

キーワード: LLR, rotation, tide, Moon, retro-reflector, simulation  
Keywords: LLR, rotation, tide, Moon, retro-reflector, simulation

## Determination of the normal modes of the Moon's libration Determination of the normal modes of the Moon's libration

Nicolas Rambaux<sup>1\*</sup>, James Williams<sup>2</sup>, Dale Boggs<sup>2</sup>

Nicolas Rambaux<sup>1\*</sup>, James Williams<sup>2</sup>, Dale Boggs<sup>2</sup>

<sup>1</sup>University Paris 06, IMCCE, Observatory of Paris, <sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology

<sup>1</sup>University Paris 06, IMCCE, Observatory of Paris, <sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology

The Lunar Laser Ranging experiment has been active since 1969 when Apollo astronauts placed the first retroreflector on the Moon. The data accuracy of a few centimeters, over a time-span of several decades, along with the numerically integrated ephemeris, DE421, encourages analysis of the lunar physical librations, and especially the detection of three modes of free physical librations (longitude, latitude, and wobble modes). This analysis was performed by using iteratively a frequency analysis and linear least-squares fit of the wide spectrum of DE421 Moon's physical librations. From this analysis we identified and estimated about 130 terms in the angular series for latitude librations and about 70 terms in the longitude angle and polar coordinates. In this determination, we found the non-negligible amplitude of the three modes of free physical libration. The determined amplitudes become 1.296'' in longitude (after correction of two close forcing terms), 0.032'' in latitude and 8.196'' X 3.312'' for the wobble, with the respective periods of 1056.13 days, 8822.88 days (referred to the moving node), and 27257.27 days. The presence of such terms, despite short damping times of 104 to 106 yr, suggests the existence of some source of stimulation acting in geologically recent times.

キーワード: Libration, Moon

Keywords: Libration, Moon

## THE THEORY OF THE MOON ROTATION AND DETERMINATION OF FOURTH MODE ITS FREE PHYSICAL LIBRATION THE THEORY OF THE MOON ROTATION AND DETERMINATION OF FOURTH MODE ITS FREE PHYSICAL LIBRATION

Yury Barkin<sup>1\*</sup>, Hideo Hanada<sup>2</sup>, Koji Matsumoto<sup>2</sup>, Sho Sasaki<sup>2</sup>, Jose Manuel Ferrandiz Leal<sup>3</sup>, Mikhail Barkin<sup>4</sup>  
Yury Barkin<sup>1\*</sup>, Hideo Hanada<sup>2</sup>, Koji Matsumoto<sup>2</sup>, Sho Sasaki<sup>2</sup>, Jose Manuel Ferrandiz Leal<sup>3</sup>, Mikhail Barkin<sup>4</sup>

<sup>1</sup>Sternberg Astronomical Institute, Moscow, Russia, <sup>2</sup>National Astronomical Observatory of Japan, Mizusawa, Japan, <sup>3</sup>Alicante University, Alicante, Spain, <sup>4</sup>Moscow Aviation Institute, Moscow, Russia

<sup>1</sup>Sternberg Astronomical Institute, Moscow, Russia, <sup>2</sup>National Astronomical Observatory of Japan, Mizusawa, Japan, <sup>3</sup>Alicante University, Alicante, Spain, <sup>4</sup>Moscow Aviation Institute, Moscow, Russia

**Resume.** In result of the comparison of our analytical theory of lunar physical libration, in particular the free librations, and available now, the empirical theory of the Moon's rotation (Rambaux, Williams, 2011), we have identified period, amplitude, and the initial phase of the forth mode of free libration of the Moon, caused by liquid core.

**Analytical theory.** The main study here is a construction and development of a highly accurate analytical theory of physical libration of the two-layer Moon (with uniform ellipsoidal liquid core and non-spherical elastic mantle). The core of the Moon is modeled by ellipsoid with an ideal homogeneous fluid. The mantle is considered as non-spherical solid body. The theory is developed on the basis of the canonical equations in Andoyer - Poincare variables and by special methods of the perturbation theory on construction of quasi-periodic solutions and investigation of their vicinity (based on the relevant equations in variations). The tables of values of the amplitudes and periods of forced and free librations for Andoyer ? Poincare variables describing the libration of the Moon and the core, for the variations of the components of the angular velocity of rotation of the Moon and the angular velocity of rotation of the coordinate system of Poincare (with respect to which a simple fluid motion is determined) have been obtained and studied. In first we have studied contributions in librations of the Moon of the second harmonic of selenopotential in accordance with the modern Selena model of gravitational field of the Moon (Matsumoto et al., 2010). The novelty of the theory and its practical significance are determined by the following principal provisions:

1.New forms of equations of physical libration of the two-layer model of the Moon (in particular in Andoyer ? Poincare variables) and new methods for their study; 2.Highly accurate description of the developments of spherical functions of the coordinates of the Moon, in the expression of the force function; 3.The new two-layer Mizusawa model of the Moon and Selena model of the gravitational field of the Moon; 4.Cassini rotation of the Moon, forced and free librations of the Moon in analytical form and their tables; 5.Dynamical effects in forced and in free librations caused by a liquid core; 6.Dynamical effects in forced and free librations of the Moon caused by its elasticity; 7.Determination of the forth mode of free libration caused by the liquid core; 8.Identification of some terms of modern Rambaux-Williams empirical theory.

**Determination of the period, amplitude and phase of the fourth mode of the free libration of the Moon caused by the liquid core.** We have been compare free libration terms from our analytical theory with some unidentified terms from empirical theory (Rambaux, Williams, 2011). In results 8 unidentified terms for classical variables in empirical theory were explained and amplitude, initial phase of the Moon free libration have been determined. The period of free libration of the pole of the Moon with liquid ellipsoidal core appreciated by us in 205.7 yr. The amplitude and initial phase of Poincare long-periodic argument of the free libration in pole motion due to liquid core have been determined in 0°0395 and -134 degrees (for initial epoch 2000.0 JD). In accordance with developed analytical theory this period corresponds to the sum of dynamic compressions of the core in  $7.24 \times 10^{-4}$ , that is in agree with seismographic data and data of laser observations (Barkin, Hanada et al., 2012). In assumption about similarity of ellipsoidal core and the entire Moon we have obtained the estimations of oblatenesses of the liquid core:  $4.42 \times 10^{-4}$  and  $2.83 \times 10^{-4}$ .

References:

Rambaux N., Williams J.G. (2011) *Cel. Mech. and Dyn. Astron.*, Vol. 109, Issue 1, pp.85-10.

Matsumoto K. et al. (2010) *J. Geophys. Res.*, doi:10.1029/2009JE003499.

キーワード: Moon, rotation, resonance, free librations, LOD

Keywords: Moon, rotation, resonance, free librations, LOD

## 測地および月震データによる月内部構造制約のシミュレーション研究

### A simulation study for constraining the lunar internal structure by geodetic and seismic data

松本 晃治<sup>1\*</sup>, 山田 竜平<sup>1</sup>, 菊池 冬彦<sup>1</sup>, ホーセンス サンダー<sup>2</sup>, 鎌田 俊一<sup>5</sup>, 岩田 隆浩<sup>3</sup>, 花田 英夫<sup>1</sup>, 石原 吉明<sup>4</sup>, 佐々木 晶<sup>1</sup>  
Koji Matsumoto<sup>1\*</sup>, Ryuhei Yamada<sup>1</sup>, Fuyuhiko Kikuchi<sup>1</sup>, Sander Goossens<sup>2</sup>, Shunichi Kamata<sup>5</sup>, Takahiro Iwata<sup>3</sup>, Hideo Hanada<sup>1</sup>,  
Yoshiaki Ishihara<sup>4</sup>, Sho Sasaki<sup>1</sup>

<sup>1</sup> 国立天文台, <sup>2</sup>NASA ゴダード宇宙飛行センター, <sup>3</sup> 宇宙航空研究開発機構, <sup>4</sup> 産業技術総合研究所, <sup>5</sup> 東京大学  
<sup>1</sup>National Astronomical Observatory of Japan, <sup>2</sup>NASA Goddard Space Flight Center, <sup>3</sup>JAXA, <sup>4</sup>AIST, <sup>5</sup>University of Tokyo

Internal structure and composition of the Moon provide important clue and constraints on theories for how the Moon formed and evolved. The Apollo seismic network has contributed to the internal structure modeling. Efforts have been made to detect the lunar core from the noisy Apollo data (e.g., [1],[2]), but there is scant information about the structure below the deepest moonquakes at about 1000 km depth. On the other hand, there have been geodetic studies to infer the deep structure of the Moon. For example, LLR (Lunar Laser Ranging) data analyses detected a displacement of the lunar pole of rotation, indicating that dissipation is acting on the rotation arising from a fluid core [3]. Bayesian inversion using geodetic data (such as mass, moments of inertia, tidal Love numbers  $k_2$  and  $h_2$ , and quality factor  $Q$ ) also suggests a fluid core and partial melt in the lower mantle region [4]. Further improvements in determining the second-degree gravity coefficients (which will lead to better estimates of moments of inertia) and the Love number  $k_2$  will help us to better constrain the lunar internal structure. Such improvements will be made by future lunar missions including Japanese SELENE-2. A preliminary simulation study shows that the  $k_2$  accuracy of better than 1% is anticipated by the SELENE-2 differential VLBI mission for which one of the radio sources is fixed on the moon serving as the reference to determine the orbiter's trajectory.

We carried out a feasibility study using Bayesian inversion on how well we can constrain the lunar internal structure when such improvements are made on the geodetic data. It is difficult to tightly constrain the internal structure from the geodetic data only because there are trade-offs among crust, mantle, and core structures. However, when combined with the existing Apollo seismic data which constrain the structures of crust and mantle, such geodetic data will contribute to narrow the range of the core structure models. We will discuss the impact of the crustal structure uncertainties on the estimation of the core structure, and also the assumption we have to place on the mantle structure in order to recover the core structure.

#### References

- [1] Weber et al. (2011), *Science*, 331, 309-312, doi:10.1126/science.1199375
- [2] Garcia et al. (2011), *PEPI*, doi:10.1016/j.pepi.2011.06.015
- [3] Williams et al. (2001), *JGR*, 106, E11, 27,933-27,968
- [4] Khan and Mosegaard (2005), *GRL*, 32, L22203, doi:10.1029/2005GL023985

キーワード: 月, 重力場, 潮汐ラブ数, 内部構造, VLBI, SELENE-2

Keywords: Moon, gravity field, tidal Love number, internal structure, VLBI, SELENE-2

## 月深部状態から揮発性成分を探る：SELENE-2 VLBI-LLR 計画

### Investigation of lunar interior volatile from the state of the core and the lower mantle: SELENE-2 VLBI-LLR proposals

佐々木 晶<sup>1\*</sup>, 菊池 冬彦<sup>1</sup>, 松本 晃治<sup>1</sup>, 野田 寛大<sup>1</sup>, 荒木 博志<sup>1</sup>, 花田 英夫<sup>1</sup>, 山田 竜平<sup>1</sup>, 國森 裕生<sup>2</sup>, 岩田 隆浩<sup>3</sup>

Sho Sasaki<sup>1\*</sup>, Fuyuhiko Kikuchi<sup>1</sup>, Koji Matsumoto<sup>1</sup>, Hirotomoto Noda<sup>1</sup>, Hiroshi Araki<sup>1</sup>, Hideo Hanada<sup>1</sup>, Ryuhei Yamada<sup>1</sup>, Hiroo Kunimori<sup>2</sup>, Takahiro Iwata<sup>3</sup>

<sup>1</sup> 国立天文台, <sup>2</sup> 情報通信研究機構, <sup>3</sup> 宇宙科学研究所/JAXA

<sup>1</sup>National Astronomical Observatory of Japan, <sup>2</sup>NICT, <sup>3</sup>ISAS/JAXA

現在有力な月の起源説では、46億年前の原始地球に火星サイズの天体が衝突したときに放出された物質から形成されたという、ジャイアントインパクト（巨大衝突）説が有力である。形成時の月は、高温熔融状態で、マグマオーシャンが形成され、その中で金属核が分離した。月内部は揮発性物質に乏しくなるはずである。

ルナプロスペクタ、かぐやの磁気探査により、月表層の磁気異常が確実となり、過去に熔融核でダイナモ作用が起きた可能性が高い。月レーザ測距による自轉變動の解析から、誤差が大きいが現在も核が融けていることが示唆されている。月下部マントルでは月震源が無いことから、剛性率が低く、融けているか、柔らかい可能性がある。しかし、アポロ着陸計画で設置された熱流量計のデータの再解析は、従来の見積よりも低い熱流量値を示す。高温月内部モデルは考えにくい。月の核が融けているならば、硫黄が融点を下げている可能性が高い。月の下部マントルは、地球のアセノスフェアの圧力に相当するので、月の深部に水が存在すれば剛性率は低くなる。月内部の揮発性成分の存在は、巨大衝突による月の高温起源説に修正を求めることになる。

月の深部状態を明らかにするため、我々は低次重力場係数と潮汐ポテンシャルラブ数  $k_2$  の精度改善を目的とした新たな月重力場計測ミッションを SELENE-2 へ提案している。着陸機と周回衛星の両方に VLBI 用電波源を搭載し、同一ビーム VLBI 観測を用いて重力場を計測する。かぐやでは 2 機の小型衛星の軌道を同時に求めたのに対して、SELENE-2 では、電波源の一方が月面に固定されており、これを基準としてもう一方の電波源（周回衛星）の軌道を決定するという手法をとる。また、電波源間の離角が常時同一ビームの条件を満たすように周回衛星の軌道を選定すること、また 2 つの電波源の離角が大きいときに別々の受信機で観測すること（2 ビーム）により、高精度で大量の VLBI 観測データを取得し、低次重力場係数と  $k_2$  の精度を効率的に向上させることを目指す。

また、月レーザ測距では、反射板が月面の南半球に無いこと、秤動により反射板アレイの両端の間で時間差が生じるため、月内部のエネルギー消散過程に関係する微小変動を求めるには精度がこれまで不十分であった。SELENE-2 では、既存の反射板から離れた南半球に新たな単一型の反射鏡を設定して、月回轉變動を高精度に測定することを提案している。

キーワード: 月の起源・進化, 月下部マントル, 月コア, 潮汐ラブ数, 月回転, 揮発性成分

Keywords: origin and evolution of the moon, lunar lower mantle, lunar core, tidal love number, lunar rotation, lunar volatiles

## METHOD OF ANALYSIS OF EXISTANCE OF THE ZONES OF LOWER SEISMIC VELOCITIES OF PLANETS AND SATELLITES METHOD OF ANALYSIS OF EXISTANCE OF THE ZONES OF LOWER SEISMIC VELOCITIES OF PLANETS AND SATELLITES

Yury Barkin<sup>1\*</sup>, Hideo Hanada<sup>2</sup>, Mikhail Barkin<sup>3</sup>

Yury Barkin<sup>1\*</sup>, Hideo Hanada<sup>2</sup>, Mikhail Barkin<sup>3</sup>

<sup>1</sup>Sternberg Astronomical Institute, Moscow, Russia, <sup>2</sup>Astronomical Observatory of Japan, Mizusawa, Japan, <sup>3</sup>Moscow Aviation Institute, Moscow, Russia

<sup>1</sup>Sternberg Astronomical Institute, Moscow, Russia, <sup>2</sup>Astronomical Observatory of Japan, Mizusawa, Japan, <sup>3</sup>Moscow Aviation Institute, Moscow, Russia

In connection with the planned space missions to the Moon and Mars to study the internal structure of these celestial bodies by seismic methods seem to be very relevant theoretical studies of possible features or internal structures of these celestial bodies. The mechanism of forced oscillations of the core and mantle of the celestial bodies and study of their geodynamic and geophysical consequences gives us opportunity to study some inner structures of the Earth, the Moon and Mars and some others celestial bodies (in particular the zones of lower seismic velocity).

The mechanism of forced relative oscillations, displacements and rotations of shells of given celestial body under the action of the gravitational attraction of external celestial bodies [1] in the last decade has attracted wide attention of specialists in various geosciences and planetary science. On the base of this geodynamical mechanism and model some fundamental problems of geodynamics and celestial mechanics, geology, geodesy and geophysics have been solved in last 10-15 years. Here we study the possible role of forced relative oscillations of the core and mantle of the Earth and Mars, some satellites of Jupiter and Saturn in the formation of the shell structures (layers) of these celestial bodies. First and foremost, the existence and nature of the zones of low seismic velocities (LVZ), as well as the zones of the extreme radial deformation of the spherical layer of the mantle. For the Earth and Mars the zones of low seismic velocity correspond to the spherical zones of the mantle for which the displacements of the particles due to gravitational action of displaced core are either small or absent and the change in directions of radial displacements is observed.

Our results suggest that for the Earth, the origin of the low-velocity zone and its position (with mean depth about 144 km) is determined by the mechanism of forced displacements of the Earth's core. A similar low-velocity zone at about 300 km depth we have been predicted for Mars. The style of deformations of the mantle layers of the Mars and the Earth (and probably for Venus) have much in common. Therefore the existence of the low-velocity zone of the planet Mars, at a depth of 300 km, seems quite real. Similar studies we fulfill now for some synchronous satellites of Jupiter and Saturn (Io, Ganymede, Europa, Titan, Enceladus, etc.) to identify zones of extreme radial deformations that occur during forced relative radial oscillations of the shells. In the report the preliminary findings on the positions situation of the zones with extreme largest deformation and the zone in which the deformations are small or absent are discussed.

### References:

[1] Barkin Yu.V. (2002) Proc. of the Section of Earth Sciences, Russian Academy of Natural Sciences. Moscow, VINITI. No. 9, C. 45-97. In Russian.

キーワード: Core, motion, mantle, deformation, existence, LVZ

Keywords: Core, motion, mantle, deformation, existence, LVZ

## 測月学および測地学のための小型望遠鏡の開発におけるいくつかの技術的課題 Some technological problems in development of a small Telescope for selenodesy and geodesy

花田 英夫<sup>1\*</sup>, 鶴田 誠逸<sup>1</sup>, 荒木 博志<sup>1</sup>, 浅利 一善<sup>1</sup>, 鹿島 伸悟<sup>1</sup>, 田澤 誠一<sup>1</sup>, 野田 寛大<sup>1</sup>, 松本 晃治<sup>1</sup>, 佐々木 晶<sup>1</sup>, 船崎 健一<sup>2</sup>, 佐藤 淳<sup>2</sup>, 谷口 英夫<sup>2</sup>, 加藤 大雅<sup>2</sup>, 菊池 護<sup>2</sup>, 伊藤 陽介<sup>2</sup>, 千葉 皓太<sup>2</sup>, 稲葉 健太<sup>2</sup>, 横川 琳吾<sup>2</sup>, 宇都宮 真<sup>3</sup>, 郷田 直輝<sup>4</sup>, 矢野 太平<sup>4</sup>, 山田 良透<sup>5</sup>, 丹羽 佳人<sup>4</sup>, 國森 裕生<sup>6</sup>, 平 勳松<sup>7</sup>, 岩田 隆浩<sup>8</sup>, 日置 幸介<sup>9</sup>  
Hideo Hanada<sup>1\*</sup>, Seiitsu Tsuruta<sup>1</sup>, Hiroshi Araki<sup>1</sup>, Kazuyoshi Asari<sup>1</sup>, Shingo Kashima<sup>1</sup>, Seiichi Tazawa<sup>1</sup>, Hiroto Noda<sup>1</sup>, Koji Matsumoto<sup>1</sup>, Sho Sasaki<sup>1</sup>, Ken-ichi Funazaki<sup>2</sup>, Atsushi Satoh<sup>2</sup>, Hideo Taniguchi<sup>2</sup>, Hiromasa Kato<sup>2</sup>, mamoru.kikuchi@iwate-u.ac.jp<sup>2</sup>, Yosuke Itoh<sup>2</sup>, Kouta Chiba<sup>2</sup>, Kenta Inaba<sup>2</sup>, Ringo Yokokawa<sup>2</sup>, Shin Utsunomiya<sup>3</sup>, Naoteru Gouda<sup>4</sup>, Taihei Yano<sup>4</sup>, Yoshiyuki Yamada<sup>5</sup>, Yoshito Niwa<sup>4</sup>, Hiroo Kunimori<sup>6</sup>, Jinsong Ping<sup>7</sup>, Takahiro Iwata<sup>8</sup>, Kosuke Heki<sup>9</sup>

<sup>1</sup> 国立天文台 RISE, <sup>2</sup> 岩手大学, <sup>3</sup> 宇宙航空研究開発機構, <sup>4</sup> 国立天文台 JASMINE, <sup>5</sup> 京都大学, <sup>6</sup> 情報通信研究機構, <sup>7</sup> 北京天文台, <sup>8</sup> 宇宙科学研究所, <sup>9</sup> 北海道大学

<sup>1</sup>RISE, NAOJ, <sup>2</sup>Iwate University, <sup>3</sup>JAXA, <sup>4</sup>JASMINE, NAOJ, <sup>5</sup>Kyoto University, <sup>6</sup>NICT, <sup>7</sup>Beijing Astronomical Observatory, CAS, <sup>8</sup>ISAS, JAXA, <sup>9</sup>Hokkaido University

Observations of the lunar rotation are one of the essential and basic selenodetic observations for investigation of the interior of the Moon as well as those of gravity fields, and high accuracy of the observations have a potential to detect signals related to the structure of lunar deep interior including the core. Technological development dedicated to highly accurate observations for the Moon, on the other hand, can return to the application to the Earth again. We are developing a small telescope for observations of Lunar rotation with a target accuracy of 1 milli arc second, and it has a potential to observe deflection of the vertical on the ground with high accuracy. This accuracy is epochal also for the observation of the deflection of the vertical even if it may be deteriorated to some extent by atmospheric fluctuations etc.

Major problems in the observations on the Moon are large temperature change and the difficulty of adjustment. We developed an objective using a diffractive lens in order to loosen the condition of the temperature change, and we adopted PZT (Photographic Zenith Tube) having a horizontal reference plane of a mercury surface in order to realize an adjustment-free system. Observations on the ground, on the other hand, are mostly affected by ground vibrations and atmospheric fluctuations. The effect of temperature change is not very large and it is relatively easy to control the temperature around the tube.

As the results of laboratory experiments, it is possible that the vibration of the mercury surface caused by the ground vibrations lead to fluctuations of star positions on CCD as large as 1 second of arc. The amplitude of the fluctuations depend on the amplitude of the ground vibrations and the depth of mercury pool. We can reduce the effect of the vibrations by making the mercury pool shallow down to the minimum depth. In the case of the mercury pool of 64mm diameter, the depth of 0.5mm is the minimum depth judging from our experience. Shallower pool will shorten the life time and will be affected more easily by the tilt.

It is important to keep the proper period of the mercury pool away from the period of ground vibrations in order to avoid the resonance. It is also effective to lengthen the integration time, and it can improve the reliability of the mean value of the center of a star image by statistical procedure. Adaptive optics is widely used for compensating the effects of atmospheric fluctuations and for obtaining sharper images approaching the diffraction limit. The adaptive optics, however, is not always effective for the astrometric telescope like PZT because it is possible to shift the center of star image by deformable and tip-tilt mirrors.

We investigate the cause of fluctuations which can affect the observations on the ground, and we explore the possibility for a new effective observations with the telescope like PZT.

キーワード: 月回転, 鉛直線偏差, 写真天頂筒, 水銀面, 地盤振動, 熱変形

Keywords: lunar rotation, deflection of the vertical, PZT, mercury surface, ground vibration, thermal deformation

## New findings in Earth's sciences New findings in Earth's sciences

Yury Barkin<sup>1\*</sup>, Stanislav Perov<sup>2</sup>, Nikolay Sidorenkov<sup>3</sup>, Hideo Hanada<sup>4</sup>  
Yury Barkin<sup>1\*</sup>, Stanislav Perov<sup>2</sup>, Nikolay Sidorenkov<sup>3</sup>, Hideo Hanada<sup>4</sup>

<sup>1</sup>Astronomical Institute of Lomonosov State University, Moscow, Russia, <sup>2</sup>Central Aerological Observatory, Moscow, Russia, <sup>3</sup>Hydrometeorological Centre of Russia, Moscow, Russia, <sup>4</sup>National Astronomical Observatory of Japan, Mizusawa, Japan  
<sup>1</sup>Astronomical Institute of Lomonosov State University, Moscow, Russia, <sup>2</sup>Central Aerological Observatory, Moscow, Russia, <sup>3</sup>Hydrometeorological Centre of Russia, Moscow, Russia, <sup>4</sup>National Astronomical Observatory of Japan, Mizusawa, Japan

An excitation mechanism and forced relative translational oscillations, swing and rotation of the Earth shells (and other planets and moons) under the gravitational action of external celestial bodies is a powerful source of endogenous activity with pronounced cyclical manifestations at different time scales endogenous energy of the planet as an open thermodynamic system (represented by the scheme with threads of different types of energy, the entropy and others) made it through gravitational interaction of external celestial bodies. The energetic of this mechanism allows to specify the energy budget of the Earth in comparison with contributions of another acting mechanisms. The estimates of the power dissipation for a viscous-elastic deformation of the Earth's mantle caused by the relative displacements of the center of mass of the mantle and core have been obtained. The inversion phenomenon of the Earth's climate changes with respect to Northern and Southern hemispheres its contemporary manifestations are analyzed and discussed.

For analysis the general global astrophysical factor - Earth's rotation rate (ERR) was used. The important role of the lunar-solar gravitational tides for weather, climate and geophysical processes in the atmosphere, oceans, and other geospheres and in the biosphere has been shown. The role of tides by using the ERR has been demonstrated on the base of observational data on precipitation in the Indian monsoon period, in the formation of tropical depressions and typhoons, in the perturbations of Earth's magnetosphere and during strong earthquakes, as well as in medical terms cardiovascular system patients and other diseases. Earth's rotation rate is so high compared to the speed of the proper motion of the tidal waves in the solar system of reference, we are dealing only with quasi-diurnal waves and their sub-harmonics. At spectral or harmonic analysis measuring low frequency waves of gravitational tides also merge with the harmonics of daily or annual thermal tides and become virtually invisible to learn. To low-frequency tidal waves are not lost in the spectral analysis, it is necessary to exclude the effects of rotation and revolution of the Earth, that is demodulated time series measurements. It is enough to fix the period of measurement (one measurement: for a day, to prevent rotation of the Earth, or the year to avoid the annual revolution of the Earth). Even before it was established that the weather changes during the lunar month is synchronized to within 0-2 days from the extremes of the EER, describing the motion of the Moon and the Earth around the barycenter. We have detected and studied a week and semi-month lunar tidal waves in the spectrum of the angular momentum of the atmosphere. Estimates have shown that classical gravitational tidal forces can not explain the tremendous energy associated with variations of the angular momentum of the atmosphere. We discuss possible role of the resonance effects in origin observed cyclicities in natural processes. Big prospects here open in dynamical and empirical studies of new tides the nature of which is connected with action of mechanism of forced oscillations of the Earth shells in gravitational field of the Moon, the Sun and all another bodies of solar system. In particular it was shown that mechanism of new tides for discussed in report frequencies and periods of oscillations generates phenomenal power of dissipations (and variations of planetary heat flow) of the order 10 (15) Wt. This energetic is sufficient for explanation of observed activity and cyclicities of all planetary processes.

キーワード: angular momentum, atmosphere, magnetosphere, tidal waves, climate changes  
Keywords: angular momentum, atmosphere, magnetosphere, tidal waves, climate changes



## Geodetic consequences of the northern drift of the Earth's core and their confirmations in the space geodesy data

## Geodetic consequences of the northern drift of the Earth's core and their confirmations in the space geodesy data

Yury Barkin<sup>1\*</sup>, Wen-Bin Shen<sup>2</sup>, Shuanggen Jin<sup>3</sup>

Yury Barkin<sup>1\*</sup>, Wen-Bin Shen<sup>2</sup>, Shuanggen Jin<sup>3</sup>

<sup>1</sup>Sternberg Astronomical Institute at Moscow State University, Moscow, Russia, <sup>2</sup>Dept of Geophysics, School of Geodesy and Geomatics, Wuhan University, China, <sup>3</sup>Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai, China

<sup>1</sup>Sternberg Astronomical Institute at Moscow State University, Moscow, Russia, <sup>2</sup>Dept of Geophysics, School of Geodesy and Geomatics, Wuhan University, China, <sup>3</sup>Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai, China

In the last 10 - 15 years the Russian and Chinese scientists have been fulfilled geodetic studies of contrast changes in the shape of the Earth, in the northern and southern hemispheres. In the works of Chinese scientists were executed empirical studies of secular variations of volumes of the northern and southern hemispheres, secular changes in the lengths of the circles of latitude in the southern and the northern hemispheres, the secular changes in the mean radius of the northern and southern hemispheres on the basis of current data space geodesy and VLBI data. For what was made a careful selection of monitoring stations and analyzed long series of high-precision measurements of the radii of stations and their displacements in the basically Earth's reference frames. In the works of Prof. Y. Barkin and colleagues (starting from 1995 -1996) the modern geodynamics of the forced librations of the core and mantle of the Earth by the gravitational attraction of external celestial bodies has been developed and has been given wide applications in geosciences, in particular in geodesy.

The most important result is the prediction and justification of the existence of the secular trend of the center of mass of the Earth as a consequence of the secular northern drift of the core of the Earth relative to the mantle. The wide geodynamic, geophysical, geodetic studies and their role in climatic change, seismic and volcanic activity, and in many other natural processes have been fulfilled in the last 15 years. Modern DORIS satellite observations (in space geodesy) indicate the existence of the secular polar drift of the center of mass of the Earth (to the North) at 5.29 mm / year. This drift reflects the Earth's core drift (drift of the center of mass of the core relatively to the center of mass of the mantle) at a rate of  $27.4 \pm 0.8$  mm / year (Barkin, 2005). Gravitational effect of the shifting core causes deformations of all layers of the mantle and various offsets of its points (both on the surface and inside the Earth). As a result of these deformations the mean radius of the northern hemisphere increases with secular velocity about 0.17 mm / year, and the mean radius of the southern hemisphere on the contrary, decreases with the same magnitude of velocity - 0.17 mm / year (Barkin, 2005, 2011). As a result of careful processing of satellite data and VLBI observations at 845 stations in a recent paper (Wenbin Shen et al., 2012, private communication) were obtained related values of  $0.46 \pm 0.01$  mm / year and  $-0.19 \pm 0.01$  mm / year, respectively. The theoretical value of the greatest secular velocity of lengthening of latitudinal circles in the southern hemisphere at latitude 45° S is  $4.17 \pm 0.12$  mm / year, and in the northern hemisphere the secular velocity of shortening of latitudinal circle (for latitude 45° N) is  $-4.17 \pm 0.12$  mm / year. According to the processing data of GPS observations Jin Shuanggen in 2005 has obtained related values of secular velocity in  $4.2 \pm 0.5$  mm / year and  $10.0 \pm 1.0$  mm / year for the corresponding hemispheres (Jin, 2005; Barkin, Jin, 2006, 2007). Thus the main trends of geodetic changes of the northern and southern hemispheres has obtained a nice explanation. Revealed an asymmetry in the shape of the Earth changes are apparently related to the formation of zones of rifting and subduction zones in their asymmetrical arrangement in the northern and southern hemispheres. Also in our joint work we have fulfilled studies of the dependence of the mean radius of the Earth and the mean velocity of secular change in the length of the latitudinal belts from the latitude (Barkin, Jin, 2007; Wenbin Shen, 2012). Theoretical results and data of satellite observations are in good agreement. The report also discusses the expected related phenomena on the Moon and Mars and the possibility of theoretical prediction and detection by high-precision observations in the planned space missions.

キーワード: drift of center of mass, secular geodesy variations, mean radiuses of hemispheres, the Earth, Mars

Keywords: drift of center of mass, secular geodesy variations, mean radiuses of hemispheres, the Earth, Mars

## Manifestations of tides in geospheres and in the biosphere Manifestations of tides in geospheres and in the biosphere

Stanislav Perov<sup>1\*</sup>

Stanislav Perov<sup>1\*</sup>

<sup>1</sup>Central Aerological Observatory, Russia

<sup>1</sup>Central Aerological Observatory, Russia

The gravitational tides in the atmosphere are recorded as the waves with the periods close to one day and its sub harmonics. Some of them are usually interpreted as the proper atmospheric modes. They commonly have either the amplitude or the frequency modulations. A new explanation of the quasi-diurnal and quasi-semidiurnal tides lines in the spectrum of the atmospheric angular momentum and other atmospheric characteristics is proposed. The role of gravity tides in the dynamics of the atmosphere and the ocean is underestimated. The reasons of a wrong estimation of a role of the tidal phenomena in geophysics are explained.

It's shown here that zonal tidal forcings with periods of ~3.5 and ~7 days determine processes in the atmosphere (weather change, formation of tropical depressions, cyclones (including typhoons), monsoon precipitation periodicity, in the magnetosphere (increasing geomagnetic activity), in the lithosphere (seismic-disturbances, including earthquakes) and increase the medical statistics in cardio-vascular illnesses and in other ones as well (see below). Luni-solar gravitational tides correlates absolutely with Earth rotation rate (ERR) and can be calculated for any time in advance. The correlation of 3,5 and 7 days periods of weather change with the same periods of human health parameters has been established. That has been demonstrated during International Conference "Space Weather Effects on Humans: in Space and on Earth" (Moscow, from 4-8 June 2012) and II Russian Conference on chronobiology and chronomedicine (in the frames of XIV World congress "Health & Education millennium" (14-17 November 2012, Moscow, Russia).

We discuss some new physical processes which are very important for Earth's climate. An example of our analysis of a number of annual rings of Japanese cypress, 800 years (data for 1100-1920 taken from the book Selected papers on Climatic Change written by well known Japanese climatologist H.Arakava) is given. This analysis shows a clear link of QBO periods with periods of El Nino - Southern Oscillation and the Chandler period (CP) first mentioned by Sidorenkov. Major and decisive role for the Earth as an open system, are external to it gravitationally interacting system of "oscillators" (Moon, Earth, Sun and planets) and gravity (as well as for the thermal atmosphere of the Earth) tidal forces. The interaction of the tides of all types and multiple binding modes with phase transitions of water vapor forms quasi-three-dimensional structures. Such structures have been discovered on space-time sections, constructed from orbital observations from satellites. We suggest new concept of tidal phenomena in the atmosphere, which is in conflict with the existing classical theory of Chapmen and Lindzen

After conducting field experiments in southern India, and at the equator in the Indian Ocean in March and June 1990 within the framework of the international program DYANA, observing a wide range of fluctuations of the ozone layer and the middle atmosphere of the author formulated a working hypothesis - the atmosphere is a system of oscillators interacting with each other and with oscillators in other geospheres and in space.

A new approach to the problem of long- and short-term prediction of atmospheric phenomena should be done, taking into account all types of tides and their multiple modes. Results of successful experiments on modification of meteorological processes are given.

So, the 1st main conclusion must be done that atmosphere/ocean phenomena are deterministic processes rather than stochastic ones. That leads inevitably to the 2nd important inference: so called the time limit of forecasting (f.e. for weather change) is not exist! 3d one is: we deal with open physical system that is resonant one. Mentions above will open a way to high technology of weather and climate modification.

キーワード: Luni-solar gravitational, Manifestations of tides, Earth's climate., weather modification

Keywords: Luni-solar gravitational, Manifestations of tides, Earth's climate., weather modification

## 宇宙測地学的手法による火星回転変動計測 Measurements of Martian rotational variations by space geodetic techniques

岩田 隆浩<sup>1\*</sup>, 松本 晃治<sup>2</sup>, 石原 吉明<sup>3</sup>, 原田 雄司<sup>4</sup>, 菊池 冬彦<sup>2</sup>, 佐々木 晶<sup>2</sup>

Takahiro Iwata<sup>1\*</sup>, Koji Matsumoto<sup>2</sup>, Yoshiaki Ishihara<sup>3</sup>, Yuji Harada<sup>4</sup>, Fuyuhiko Kikuchi<sup>2</sup>, Sho Sasaki<sup>2</sup>

<sup>1</sup>JAXA 宇宙研, <sup>2</sup>国立天文台, <sup>3</sup>産総研, <sup>4</sup>東大地震研

<sup>1</sup>ISAS/JAXA, <sup>2</sup>NAOJ, <sup>3</sup>AIST, <sup>4</sup>ERI, Univ. Tokyo

一般に固体天体の表層環境の変化や内部構造は、その天体の回転（自転）変動に影響を及ぼし、その変動の振幅や位相は電波航法や VLBI などの宇宙測地学的手法により観測可能である。特に火星は、地球と類似の構成要素を有しながらも、過去から現在に至る進化やダイナミクスの様相は大いに異なっており、その相違を表層環境・内部構造から解き明かすことは比較惑星学に重要である。

火星の回転変動は、従来は着陸機（ランダー）の追跡データ（2 ウェイ測距・距離変化率計測：RARR など）を利用して計測されており、例えば Viking 1 および 2, Mars Pathfinder によって、歳差や自転速度変動が得られてきた。これらの計測手法には、地球・火星間の経路上および地上局の位相安定度等に起因すると考えられる測定精度の限界がある。そこで欧州の研究グループは、周回探査機（オービター）とランダー間の RARR を利用した電波航法による精度向上を提案している。

一方我が国では、オービターとランダー等の複数探査機を用いた火星複合探査（MELOS: Mars Exploration with Lander-Orbiter Synergy）の検討が開始された。そこで我々は、4 ウェイドブラ計測（FWD）、逆 VLBI（iVLBI）等の新たな測地学的手法による、火星回転変動の高精度計測を提案する。4 ウェイドブラ計測（FWD）は、宇宙機の位置・軌道の中継機を経由した 4 経路から決定する測地手法であり、我々は月周回衛星「かぐや」での月裏側の重力場観測という実績を有している。MELOS 計画では、宇宙機の構成は未定であるが、オービターとランダーという構成が一案である。この時、追跡局 オービター ランダー オービター 追跡局の経路でドブラ計測（距離変化率計測）を実施する。同時に、オービターとランダーの 2 ウェイ RARR を実施する。本方式で期待される測定精度は従来の電波航法より向上し、オービター・ランダー間 RARR を利用した場合の精度に相当すると推定される。逆 VLBI（iVLBI）は、複数の宇宙機からの同期した信号の位相比較を行って、各々の相対位置を測定する手法であり、測定精度は周波数の逆数に比例し、地上と対象宇宙機の距離に依存しないで高精度で決定できるという特徴を持つ（Kawano et al., 1999）。MELOS 計画では、ランダーから地上への直接送信波と、オービターでコヒーレントに中継された信号を、追跡局で同時またはスイッチングにより計測する。ランダーの位置変化に対する原理的な感度は、想定される最も低い周波数である X 帯を用いた場合でも 0.3mm であり、FWD や RARR 等の従来手法と比べて著しく高精度である。一方、通信機経路内の高精度の位相補償を要することが、現状では技術課題である。

キーワード: 火星, 自転, 宇宙測地, 衛星間測距, VLBI

Keywords: Mars, rotation, space geodesy, satellite-to-satellite tracking, VLBI

## A preliminary observation of 531-day period in wobble of the polar motion A preliminary observation of 531-day period in wobble of the polar motion

Hao Ding<sup>1</sup>, Wen-Bin Shen<sup>1\*</sup>

Hao Ding<sup>1</sup>, Wen-Bin Shen<sup>1\*</sup>

<sup>1</sup>Wuhan University

<sup>1</sup>Wuhan University

Previous studies show that the polar motion contains two dominant components, namely the annual wobble (AW) with a 12-month period and the Chandler wobble (CW) with a 14-month period. Some scholars consider that the frequency of the CW varies with its amplitude; some scholars consider that CW has double or multiple frequencies; some scholars consider that the frequency of CW is invariant. In 180s, a 530-day-period wobble in polar motion was marginally detected. Since then, we did not find literatures addressing this wobble period from any kind of observations. In this preliminary study, we apply the ensemble empirical mode decomposition (EEMD) method to analyses of two kinds of observations. First, applying EEMD to two polar motion time series, the EOP C04-05 series with one-day sampling interval spanning 1962 to 2012 and the POLE2010 series with one-month sampling interval spanning 1900 to 2011, we observed a 531-day-period (about 0.68683 cpy) wobble and a 530-day-period (0.68913) wobble, respectively. Noting that the estimated amplitudes and frequencies of this wobble from the two series are different from each other, we consider that the difference is caused by the relative poor quality of the POLE2010 series during 1900-1961. Deleting the poor quality data sets, we obtain the 531-day-period wobble from both time series. Our results show that the frequency modulation of the CW may greatly suppress the 531-day-period wobble so that it cannot be observed in conventional direct power spectra of the polar motion series. Second, applying EEMD to two superconducting gravimeter records with a length of about 15 years and one-day sampling interval, we also observed the 531-day-period wobble. If the 531-day-period wobble really exists, it might be caused by the fluctuations of global atmospheric and oceanic angular momentums. Further investigations are still in progress. This study is supported by NSFC (grant No. 41174011), National 973 Project China (grant No. 2013CB733305) and NSFC (grant No. 41210006, 41128003, 41021061, 40974015).

キーワード: 531-day-period, wobble, polar motion, ensemble empirical mode decomposition, EOP and POLE series, superconducting gravimeter records

Keywords: 531-day-period, wobble, polar motion, ensemble empirical mode decomposition, EOP and POLE series, superconducting gravimeter records