

風化の反応・輸送モデリングの最近の研究動向 Recent trend of reactive transport modeling of rock weathering

横山 正^{1*}Tadashi Yokoyama^{1*}¹ 大阪大学大学院理学研究科宇宙地球科学専攻¹Department of Earth and Space Science, Osaka University

岩石の風化は、岩石表面および内部における溶解・沈殿等の反応と、間隙中の物質移動により進行する。これらの過程を定量的に理解するために、反応・輸送方程式を用いた解析（反応・輸送モデリング）が行われる。以下は、一次元の反応・輸送方程式の一例である：

$$\frac{dc}{dt} = D \frac{d^2c}{dx^2} - v \frac{dc}{dx} + Ar_0 f(c)$$

ここで、 c は溶液中の元素濃度、 t は時間、 x は距離、 θ は間隙率、 D は有効拡散係数、 v は間隙中の流速、 A は岩石の単位体積あたりの表面積、 r_0 は速度定数、 $f(c)$ は溶解速度の濃度依存性を表す関数である。反応・輸送方程式を解くことにより、岩石内部の元素濃度や溶解速度の分布、およびその時間変化が分かる。計算に必要な各パラメーターを求める方法としては、例えば D については、直接実験により測定する方法 (Yokoyama and Nakashima, 2005) や、試料の間隙率から経験式 (Archie's law) により推定する方法などがある。反応速度の濃度依存性に関しては、[1] 遷移状態理論に基づき平衡からのずれに線形依存の形で表すもの (Aagaard and Helgeson, 1982; Lasaga, 1984), [2] Al の阻害効果を考慮するもの (Oelkers et al., 1994), [3] 平衡濃度に近い場合と遠い場合とで溶解速度の濃度依存性を分けて考えるもの (Hellmann and Tisserand, 2006) などがある。石英の反応を扱う場合は、 $f(c) = (1 - c/c_{eq})$ (c_{eq} は平衡濃度) となる (Lasaga, 1998)。反応・輸送方程式を用いた計算結果と、実際の天然における風化状態とを比較することで、風化のメカニズムの議論や、天然と室内実験とで得られる風化速度が最大 5 桁食い違う (White and Brantley, 2003) 原因の検討などが行われる。

近年の反応・輸送モデリングに関する研究では、例えば、Maher et al. (2009) は、実験結果から見積もった常温での速度定数と類似した r_0 値を用いてモデル計算を行い、上述 [2] および [3] の速度則を適用した場合に土壤中で観察された風化プロファイルをよく再現できることを示すと共に、二次鉱物の沈殿が一次鉱物の溶解を支配する重要な因子であることを指摘した。Maher (2010) は、風化速度が流体の滞留時間または流速に強く依存することを指摘した。Navarre-Sitchler et al. (2011) は、玄武岩の風化のモデリングにおいて、 θ , D , A の経時変化を計算に組み込むことによって、実際に観察される風化殻の厚さ、生成速度、鉱物組成を再現できることを示した。Moore et al. (2012) は、花崗岩の風化プロファイルを計算で再現するためには、平均流速や反応面積を測定値よりも小さく設定する必要があることを指摘した。

現状の風化の反応・輸送モデリングでは、常温より高い温度 (150 °C など) の実験で得られた反応速度の濃度依存性をそのまま常温での反応の解析に用いている場合が多いが、この妥当性は自明ではない。また、二次鉱物の沈殿速度はモデリングの結果に大きく影響するが、対象とする二次鉱物 (低結晶性のアルミノケイ酸塩など) の沈殿速度の知見が不足している場合が多い。さらに、一般に地表付近では、岩石間隙が完全には水で満たされていない不飽和状態になるが、不飽和状態の反応面積がどの程度飽和状態と異なるかについてもよく分かっていない。これらの点を改善していくことが、反応・輸送モデリングの精度の向上において重要と考えられる。

キーワード: 風化, 反応・輸送モデリング

Keywords: Weathering, Reactive transport modeling

フェリハイドライトへの硫酸イオンの吸着：表面錯体モデリングとその場赤外分光観察

Adsorption of sulfate to ferrihydrite: surface complexation modeling and in-situ infrared spectroscopy

福士 圭介^{1*}, 青山和樹², 楊晨², 北台紀夫¹, 中嶋悟³

Keisuke Fukushi^{1*}, Kazuki Aoyama², Yang Chen², Norio Kitadai¹, Satoru Nakashima³

¹ 金沢大学環日本海域環境研究センター, ² 金沢大学大学院自然科学研究科自然システム学専攻, ³ 大阪大学大学院理学研究科宇宙地球科学専攻

¹Institute of Nature and Environmental Technology, Kanazawa University, ²Division of Earth and Environmental Sciences, Kanazawa University, ³Department of Earth and Space science, Graduate School of Science, Osaka University

Sulfate, a major anion in nature, can affect the mobility of coexisting dissolved toxic trace elements by influencing the adsorption properties of the species on mineral surfaces. Ferrihydrite is an important scavenger for trace elements in natural water because of its adsorption capacity and its ubiquitous formation in surface conditions. To elucidate the adsorption behavior of the trace elements in natural water, it is important to construct a predictive model of sulfate adsorption on ferrihydrite that can predict the adsorption behavior of sulfate quantitatively under widely various environmental conditions based on the spectroscopic information of sulfate adsorption.

In-situ infrared spectra for sulfate adsorption on ferrihydrite as a function of pH (3-7), ionic strength ($I = 0.01$ and 0.1) and sulfate loading ($[\text{SO}_4^{2-}] = 0.1$ and 0.2 mM) were obtained to constrain the surface speciation of sulfate on ferrihydrite. The shape of the spectra was pH-dependent. The degree of *nu*3 band splitting decreases with pH. Little difference of the spectra was found between different ionic strengths and sulfate loadings for the same pH. The little influence of ionic strength on the IR spectra indicates that the inner sphere and outer sphere species most likely do not exist simultaneously at the same pH condition. Based on the IR spectra obtained from lower pH showing the splitting of *nu*3 band to two peaks with activation of *nu*1 band, the surface species is identified to be single inner sphere monodentate sulfate. The changes of spectra with pH are most likely attributable to the changes of the electric field strength posed to the sulfate on ferrihydrite surface, which is strongly pH-dependent.

The predictive model for sulfate adsorption was constructed using an extended triple layer model (ETLM). The pH adsorption edges and proton surface charges in the presence of sulfate as a function of ionic strength and sulfate concentration were obtained, respectively, from batch adsorption and acid-base titration experiments. The sulfate adsorption on ferrihydrite increases continuously with decreasing pH and ionic strength. These macroscopic adsorption data were analyzed using ETLM to retrieve the sulfate adsorption reaction and the equilibrium constant. Results of ETLM analyses showed that adsorption of sulfate on ferrihydrite is a single monodentate inner sphere process that is consistent with in-situ infrared spectroscopic observation.

Batch adsorption data from earlier studies of sulfates on ferrihydrite were reasonably reproduced using ETLM with the same adsorption reaction and equilibrium constant.

キーワード: フェリハイドライト, 硫酸, 表面錯体モデリング, その場赤外分光

Keywords: ferrihydrite, sulfate, surface complexation modeling, in-situ infrared spectroscopy

フェリハイドライト・ナノ粒子の生成・溶解・移動・Pu吸着 Formation, dissolution, migration and Pu-sorption of ferrihydrite nanoparticles

月村 勝宏^{1*}, 鈴木 正哉¹, 鈴木 庸平², 村上 隆²

Katsuhiko Tsukimura^{1*}, Masaya Suzuki¹, Yohey Suzuki², Takashi Murakami²

¹ 産業技術総合研究所, ² 東京大学

¹AIST, ²The University of Tokyo

This paper describes the behavior of Pu and ferrihydrite nanoparticles in the Lake Karachai area. The Pu in the Lake Karachai area is sorbed to colloidal ferrihydrite and moves with the ferrihydrite in the groundwater. The Pu is incorporated in the inner part of ferrihydrite because the partition coefficients of Pu between solution and ferrihydrite have a wide variation. The ferrihydrite is formed from Fe(III), which is supplied through the dissolution of Fe(II) from metallic iron in Lake Karachai and iron-bearing minerals in rock and the oxidation of the Fe(II) by oxygen or nitrate. The formation of hematite or goethite, which is more stable than ferrihydrite, decrease the concentration of ferrihydrite. When the supply of Fe(III) exceeds the formation of hematite or goethite, the concentration increases, and when the supply of Fe(III) is less than the formation of hematite or goethite, the concentration decreases. The formation rate of hematite or goethite is proportional to the concentration of ferrihydrite nanoparticles, the formation rate decreases by more than one order of magnitude in groundwater because of the present of trace amount of dissolved silica. The migration of ferrihydrite nano-particles depends on the concentration and the size of nanoparticles, and the pH of the groundwater. When the concentration of nano-particles exceeds the critical aggregation concentration (around 10⁻³ mol Fe /L), the nano-particles aggregate and precipitate. Large nano-particles are trapped in the dimples of fractures or filtrated in constrictions. When the size of ferrihydrite nano-particles is 70 nm, which is the same as that in the Lake Karachai area, the 99 % of the particles are trapped in a dimple of 3 mm. The gravity also significantly affects the downward movement of nano-particles. The ferrihydrite nano-particles with the diameter of 70 nm move downward with the velocity of 0.12 m/year, which is significant because the assessment period of radioactive nuclear wastes is more than 10,000 years.

キーワード: 吸着, フェリハイドライト, ナノ粒子, 生成, 溶解, 移動

Keywords: sorption, ferrihydrite, nanoparticles, formation, dissolution, migration

地球表層での元素濃度、同位体比、物質循環を支配するナノ鉱物 Importance of nanominerals on the concentrations, isotopic abundances, and element cycles at earth's surfaces

高橋 嘉夫^{1*}

Yoshio Takahashi^{1*}

¹ 広島大学

¹Hiroshima University

Nano-sized minerals (nanominerals) have substantial influences on the abundances of elements, isotopic ratios, and cycles of various elements at earth's surface. In this paper, I will give examples of essential roles played by nanominerals in various phases in environment.

In the atmosphere, aerosols are very important for the accurate prediction of global warming. Aerosols smaller than 1 micron are generated by physical and chemical processes in the atmosphere. Sulfate and oxalate aerosols are main species produced by such processes, which are considered to have a cooling effect of the earth due to their high hygroscopicity to assist cloud formation. If sulfate and oxalate form insoluble metal complexes, however, the species do not absorb water, which results in the mitigation of the cooling effect. We employed X-ray absorption fine structure spectroscopy (XAFS) to distinguish the sulfate and oxalate species. As a result, it was found that most of oxalate species was actually insoluble metal complexes, showing that cooling effect of the organic acids can be smaller than expected so far. For the analysis of sulfate in such fine particles, scanning transmission X-ray microscope (STXM), which is at present constructed in Photon Factory by our group, can give distribution of various chemical species with 50 nm spatial resolution.

In soil, various nanominerals are playing an important role to control the migration of various ions. Related to the Fukushima Daiichi Nuclear Power Plant accident, it has been indicated that 2:1 type clay minerals and micas strongly adsorb (radio)cesium within the soil. The affinity of Cs and availability of Cs to plants have been evaluated by Radiocesium Interception Potential (RIP) for various minerals and soils. On the other hand, we can determine the ratio of inner- and outer-sphere complexes of Cs adsorbed on the minerals and soils. We found that the amount of inner-sphere complex has a close linear-correlation to RIP value, showing that the mobility of Cs is decided by the amount of inner-sphere complex to the solid media.

In seawater, abundances and isotopic ratios of oxyanions such as chromate, molybdate, and tungstate are governed by the adsorption reactions on the nanominerals such as iron oxyhydroxides and manganese oxides. In particular, XAFS analysis and quantum chemical calculation (QC) revealed that chromate and molybdate form almost outer-sphere complexes, whereas tungstate inner-sphere complex on ferrihydrite. In particular, it was shown that inner-sphere complex is mainly a bidentate-binuclear complex. It has been indicated that affinity of the oxyanion to Fe (oxyhydr) oxides and the attachment mode (inner- or outer- sphere complex) is interpreted by linear free energy relationship (LFER) using pKa of conjugate acids. We here found that average pKa value, or $(pKa_1 + pKa_2)/2$, is better to explain the systematics, because contributions of two acid dissociation groups are important in terms of the structure of the surface complex.

Isotopic fractionation between hydrated molybdate and that adsorbed on manganese oxide was calculated from vibrational frequencies. The largest isotopic fractionation was observed when Mo formed inner-sphere complex with octahedral coordination. In our study, it was suggested that (i) taking account of the adsorbent in the calculation and (ii) calculation of hydration including actual water molecules are needed to quantify the large isotope fractionation by QC. Structural information at atomic level coupled with QC calculation is essential to study the interaction of various chemical species with nanominerals.

As seen in this paper, nanominerals have great impacts on the migration and cycle of any elements at earth's surface. Understanding of the interaction is essential to elucidate geochemical information from geological materials and to predict behaviors and cycles of any elements at earth's surfaces and their effects on the environment.

惑星物質科学におけるナノ鉱物 Significance of nano minerals in the planetary materials science

野口 高明^{1*}

Takaaki Noguchi^{1*}

¹ 茨城大学

¹Ibaraki University

Introduction: Nowadays we can investigate diverse planetary materials in laboratories: meteorites, interplanetary dust particles, micrometeorites, returned samples from the Moon, the Wild 2 comet, and the asteroid Itokawa, and so on. These planetary materials often contain submicron minerals (hereafter nano minerals).

Nano minerals in planetary materials: Why nano minerals are common among planetary materials? Their small sizes are probably related to their formation conditions and/or mechanisms. Now we look out several examples of nano minerals in planetary materials.

Nano minerals in hydrated chondritic meteorites: CIs, CMs, CVoxBs, CRs, CHs, CBs, and some UOCs contain various amounts of hydrated minerals [e. g. 1]. Phyllosilicates in them are saponite and/or serpentine. Their crystallinity is poor and only a few to at best a few tens repetitions of (001) lattice fringes can be seen in TEM images. When they coexist, they often form mixed layers. After accretion of anhydrous materials, water ice and organics, water ice was melted probably by ²⁶Al decay heat and water was reacted with anhydrous materials within their parent bodies to form hydrated minerals and the other minerals such as carbonates. The alteration temperatures and water/rock ratios are estimated to have been different between CIs and CMs: >100 °C and ~1 and <20 °C and 0.3-0.6, respectively [e. g. 1]. Low water rock ratios and possibly stagnant fluids might be related to their poor crystallinity. Because no remarkable elemental fractionation by aqueous alteration is not observed in these meteorites, fluid may have been lost by freeze-dry on their parent bodies.

Aqueous alteration products in Martian meteorites: Nakhilites are a kind of Martian meteorites formed in lava flows or shallow intrusions [e. g. 2]. The meteorites are unique because they contain aqueous alteration products formed on Mars, which typically appear as reddish brown veinlets in olivine phenocrysts. The alteration product of Lafayette meteorite is composed of siderite, poorly crystalline saponite and serpentine, and silicate gel. The hydrothermal assemblage suggest that a CO₂-rich hydrothermal fluid reacted at 150-200 °C, pH 6-8 with a water/rock ratio of <300 [3]. Because elemental fractionation is remarkable in the alteration product, the fluid obviously moved along clacks. Another intriguing example is ALH84001 orthopyroxenite containing bacteria-like objects [4]. Although it contains truncated nanomagnetite, debates on its origin have not finished even today [5].

Nano minerals in chondritic porous interplanetary dust particles: Chondritic porous interplanetary dust particles (CP IDPs) are typically ~10 micron-sized planetary materials, which have been regarded as cometary dust and the most primitive planetary materials that can be investigated in laboratories even after the Stardust mission [6]. They are loose aggregates of nano minerals with some isolated minerals larger than 1 micron. Because CP IDPs experienced almost no modification on their parent bodies, each constituent has its own history. The most curious object in CP IDPs are GEMS (glass with embedded metal and sulfide), which is <500 nm spheroidal amorphous silicate containing nanocrystals of Fe-Ni metal and Fe sulfide [7]. Its origin is still in dispute [8].

Summary: As described above, nano minerals in planetary materials hold the key to understand their formation conditions and origins although some are still in debate.

References: [1] Brearley and Jones (1998) and the references therein. In: Planetary materials. [2] Treiman (2005) Chem. Erde 65, 203-270. [3] Bridges and Schwenzer (2012) EPSL 359-360, 117-123. [4] McKay et al. (1996) Science 273, 924-930. [5] Weiss et al. (2004) PNAS 101, 8281-8284. [6] Busemann et al. (2009) EPSL 288, 44-57. [7] Bradley (2006) In: Meteorites, Comets, and Planets. [8] Keller and Messenger (2011) GCA 75, 5336-5365.

キーワード: ナノ鉱物, 惑星物質

Keywords: Nano mineral, Planetary materials

フェリハイドライトの相変化がアンチモンの環境挙動に及ぼす影響 Rigid immobilization of antimony(V) with hydrous ferric oxide (HFO) aging

光延 聖^{1*}, 村松 千尋¹, 坂田 昌弘¹
Satoshi Mitsunobu^{1*}, Muramatsu Chihiro¹, Sakata Masahiro¹

¹ 静岡県立大学環境科学研究所

¹University of Shizuoka, Institute for Environmental Sciences

In this study, we investigated the behavior of Sb(V) during the transformation of poorly crystalline Fe(III) oxyhydroxides (two-line ferrihydrite) with various Sb/Fe molar ratios at pH 6.0. Both XRD and Fe EXAFS analyses confirmed that goethite and hematite are the primary transformation products of the ferrihydrite in the presence of Sb(V). The crystallization kinetics showed that the transformation rate with Sb(V) was approximately the same as that of the control (without Sb(V)), which indicates that the presence of Sb(V) does not influence the transformation rate to a noticeable extent. Throughout the transformation, Sb(V) dominantly partitioned in the solid phase and no desorption of Sb(V) was observed. Furthermore, Sb EXAFS analyses suggested that Sb(V) in the solid phase is structurally incorporated into crystalline goethite and/or hematite generated by the ferrihydrite transformation. Hence, Sb(V) transfers into the thermodynamically stable solids from the metastable ferrihydrite with aging, indicating a rigid immobilization of Sb(V). These findings are valuable for making predictions on the long-term fate of Sb associated with ferrihydrite in the environments.

キーワード: フェリハイドライト, XAFS, アンチモン
Keywords: ferrihydrite, XAFS, antimony

福島県土壌での非晶質物質におけるCs吸着量について The amount of Cs adsorption to the amorphous materials in the Fukushima soil

鈴木 正哉^{1*}, 月村 勝宏¹
Masaya Suzuki^{1*}, Katsuhiko Tsukimura¹

¹産総研 地質調査総合センター

¹Geological Survey of Japan, AIST

この研究では、放射性Csを含む福島県の土壌での、非晶質物質におけるCs吸着量について述べる。放射性Csは雲母などの粘土鉱物に強く吸着しているとされているが、福島県の土壌の表層は、火山灰起源による非晶質物質が多く存在する。それゆえ、粘土鉱物だけでなく、非晶質物質にもCsが吸着されている可能性がある。本研究では、福島県の放射性Csを含む土壌を水ひにより分級を行い、それぞれの画分を試料とした。各試料においては、過酸化水素にて有機物を、シュウ酸にて非晶質物質を溶解させ、溶解前後の質量を測定するとともに、放射線量の測定を行った。8~16 μm、16~32 μm、32~64 μmの試料において、重量減少率はそれぞれ47wt%、40wt%、49wt%であり、放射線量の減少率は52%、58%、52%であった。この結果により、細かい土壌各分には、非晶質物質および有機物におけるCs吸着が相当量あることが明らかとなった。

キーワード: セシウム, 非晶質, 吸着, 土壌

Keywords: cesium, amorphous, adsorption, soil

モノハイドロカルサイトの相転移におけるPbとZnの移動のモデリング：予備結果 Modelling of Pb and Zn transport during phase transformation of monohydrocalcite: Preliminary results

宗本 隆志^{1*}, 菅崎 良貴¹, 村上 隆¹
Takashi Munemoto^{1*}, Yoshiaki Kanzaki¹, Takashi Murakami¹

¹ 東京大学大学院理学系研究科地球惑星科学専攻

¹Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo

A simple model has been developed to examine Pb and Zn transport during phase transformation of monohydrocalcite. Monohydrocalcite is a nano-mineral with a diameter of ~100 nm and is metastable, being transformed to aragonite (several micrometer) in a few days. Accordingly, metals initially adsorbed on monohydrocalcite are redistributed between solution and solid during transformation, which affects metal transport. The model was made based on the results of Pb and Zn sorption experiments during transformation of monohydrocalcite (unpublished study). The model calculates changes in the concentrations of monohydrocalcite and aragonite and those of Pb and Zn sorbed to the solids and dissolved in solution with time at a given point in a water flow.

Major factors that the model considers are (i) transformation rate, (ii) change in metal distribution during transformation, (iii) change in particle size during transformation, (iv) sedimentation rate, and (v) water flow rate. The model calculations are made for the following cases: (1) with different transformation rate (e.g., retardation and inhibition) and without transformation, (2) with change in water flow rate, (3) with change in particle size of monohydrocalcite, and (4) with difference between metal redistribution mechanisms.

Metal transport is slower with transformation than without transformation because of the transformation and the subsequent sedimentation of aragonite with much larger size than monohydrocalcite. When the transformation rate is retarded, metal transport becomes faster than that with transformation, indicating important effects of transformation rate on metal transport. With slower water flow rate, metal transport is slower depending on transformation rate. The difference in particle size of hypothesized monohydrocalcite (10 and 100 nm) makes little difference when the water flow is fast. The difference in metal redistribution mechanisms (i.e., between only sorption and combination of sorption and metal precipitation) affects metal transport.

キーワード: モノハイドロカルサイト, 相転移, 移動モデリング
Keywords: monohydrocalcite, phase transformation, transport modelling

状態変化におけるナノ鉱物：炭素、レアアースと塩素含有物 Nano-Minerals Formed by State-Changes: Carbon-, Rare-Earth REE- and Chlorine-Bearing Materials

三浦 保範^{1*}

Yasunori Miura^{1*}

¹ 客員 (国内外大学)

¹ Visiting (Universities)

地球および太陽系天体において、ナノ鉱物相の形成は、状態変化を示す残存固体相と考えられる。

1) 地球惑星など三圏の気圏・海水圏・固体圏を変化する状態変化において、その過程変化の証拠として常温常圧においてナノ鉱物固体相を残存形成する。

2) 鉱物は従来マクロな(目視できる)固体相であるが、水銀など液体相も鉱物に含めており、またナノ固体を示す液体・気体からの残存相が存在するので、鉱物の定義を「広く状態変化をする一過程の状態である常温の固体相(マクロからマイクロナノ相)」と考える(AMS e-mail, Miura, 2012)。

3) 衝突爆発時の高温高圧条件での迅速反応において、気体・液体・固体(VLS)間で状態変化する時に極限状態では安定な状態相がナノ鉱物相である。例えば、炭素含有物がナノ鉱物相で表面・反応固体相・高圧構造相などを形成する。これを現在話題で科学的に未解明な高温温暖化ガスの固定に利用している(Miura, 2007; 特願・特許)。火山爆発などもこのナノ炭素含有相(下関市金雲母・浜田市萩市火山などの晶洞)が観察され、別途報告している(山口大・卒論論文; 2009)。

4) 月面のCa斜長石は、構造式と化学式の不一致があり、これを急冷時のEuや炭素の含有により説明可能である。衝突破砕岩にレアースREEが(Ca, Cと共に)多くなるのが月アポロ試料の解析で分かり、隕石衝突時にREEが多くなるのが地球上ではサドベリー破砕岩で観察できている(月資源NASA報告; Miura, 2011; Miura, 2009)。

5) 炭素と同じ衝突反応時に鉄と塩素がナノ鉱物的に形成されて、マイクロな赤金石鉱物相を形成することが分かり、人工的にもナノ鉱物相の生成が確認出来ている(山口大・卒論論文; 2009)。

6) 海水圏が豊富にある地球(地表の約7割の海水圏)での衝撃波反応時だけでなく、無水条件での地球外太陽系においても、衝突の瞬間の状態変化で、ナノ鉱物個体が生成残存するので、ナノ鉱物相に広く利用できる重要性があると考えられる。

キーワード: ナノ鉱物, 定義, 状態変化, 衝撃波反応, 炭素と塩素, レアアース

Keywords: Nano-minerals, Definition, State-changes, Shock-wave reaction, Carbon and Chlorine, Rare-Earth