

西部北太平洋における溶存有機物の光学的性質 Spectral characteristics of chromophoric dissolved organic matter in the western North Pacific

山下 洋平^{1*}, 野坂裕一², 鈴木光次¹, 小川浩史³, 高橋一生⁴, 齊藤宏明⁵

Youhei Yamashita^{1*}, Yuichi Nosaka², Koji Suzuki¹, Hiroshi Ogawa³, Kazutaka Takahashi⁴, Hiroaki Saito⁵

¹北海道大学大学院地球環境科学研究院, ²北海道大学大学院環境科学院, ³東京大学大気海洋研究所, ⁴東京大学大学院農学生命科学研究科, ⁵独立行政法人水産総合研究センター 東北区水産研究所

¹Faculty of Environmental Earth Science, Hokkaido University, ²Graduate School of Environmental Science, Hokkaido University, ³Atmosphere and Ocean Research Institute, The University of Tokyo, ⁴Graduate School of Agricultural and Life Sciences, The University of Tokyo, ⁵Tohoku National Fisheries Research Institute, Fisheries Research Agency

Chromophoric dissolved organic matter (CDOM) ubiquitously occurs in marine environments and plays a significant role in the marine biogeochemical cycles. Basin scale distributions of CDOM have recently been surveyed in the global ocean and indicate that quantity and quality of oceanic CDOM are mainly controlled by in situ production and photo-degradation. However, factors controlling the spectral parameters of CDOM at UV region (i.e., S275-295 and SR) have not been well documented. To evaluate the factor controlling the spectral characteristics of CDOM at UV region in open ocean, we determined the quantitative and qualitative characteristics of CDOM in the subarctic and subtropical surface waters (5-300 m) of the western North Pacific. Absorption coefficients at 320 nm in the subarctic region were significantly higher than those in the subtropical region throughout surface waters, suggesting that magnitudes of photobleaching were different between two regions. The values of S275-295 and SR were also significantly higher in the subtropical region compared with the subarctic region. The dark microbial incubation showed biodegradation of DOM little effected on S275-295, but slightly decreased SR. On the other hand, increases and unchanging was observed for S275-295 and SR during photo-irradiation incubations respectively. These experimental results indicated that photobleaching of CDOM mainly produced qualitative differences in CDOM at UV region between the subarctic and subtropical surface waters. The results of this study imply that S275-295 can be used as a tracer of photochemical history of CDOM in open ocean.

キーワード: 海洋, 溶存有機物, 光学的性質, 光分解

Keywords: Marine biogeochemistry, Dissolved organic matter, Spectral characteristics, Photobleaching